



**FACULTY OF MEDICINE AND
HEALTH SCIENCES**

**DEPARTMENT OF
BIOCHEMISTRY**



FACULTY OF SCIENCES



**DEPARTMENT OF
MEDICAL PROTEIN SCIENCES**

The role of the glyoxalase system and its substrate methylglyoxal in the induction of growth factor-withdrawal induced apoptosis and the regulation of the cell cycle

RUBEN AUDIFFRET

**Promotor: Prof. Dr. Katia Vancompernelle
Co-promotor: Prof. Dr. Guy Haegeman**

**Thesis submitted to fulfill the requirements for achievement of the degree of
Doctor (PhD.) in Science (Biotechnology)
Academic year 2007-2008**

Exam commission

Promotor

Prof. Dr. Katia Vancompernelle
(secretary)

Department of Biochemistry - UGent
Departement of Medical Protein Research - VIB

Co-promotor

Prof. Dr. Guy Haegeman

Department of Molecular Biology - UGent

Promotion commission

Prof. Dr. Johan Grooten
(chairman)

Department for Molecular Biology - UGent

Prof. Dr. Peter Brouckaert

Department for Molecular Biology - UGent
Department for Molecular Biomedical Research -
VIB

Prof. Dr. Emmanuel Dejjardin

Department of Virology and Immunology -
University of Liège

Prof. Dr. Patrick Van Oostveldt

Department of Molecular Biotechnology - UGent

Dr. Wim Vanden Berghe

Department of Molecular Biology - UGent

The research presented in this thesis was performed at the department of Biochemistry, Faculty of Medicine and Health Sciences, University of Ghent/ Departement of Medical Protein Research, VIB.

Ruben Audiffret was supported by IWT Vlaanderen.

Title page	1
Exam Commission	3
Table of Contents	4
Abbreviations	6
Summary	9
Samenvatting	10
 Part I:	
General introduction	11
 Chapter 1 The glyoxalase system	12
1.1 Introduction	12
1.2 Detoxification of α -oxoaldehydes	12
1.3 Glyoxalase I	13
1.4 Posttranslational modification of glyoxalase I	15
1.5 Glyoxalase II	16
1.6 Glyoxalase III	17
1.7 Possible functions of the glyoxalase system	17
 Chapter 2 Methylglyoxal	20
2.1 Introduction	20
2.2 Modification of proteins	21
2.3 Detoxification of methylglyoxal	23
2.4 Methylglyoxal as a cytostatic and cytotoxic molecule	23
2.5 Methylglyoxal in other signaling pathways	24
 Chapter 3 RAGE, the receptor for advanced glycation endproducts	27
3.1 Introduction	27
3.2 The receptor for AGEs	27
3.3 Signaling through RAGE	28
3.4 Involvement of RAGE in proliferation and survival	30
 Chapter 4 Growth factors	32
4.1 Introduction	32
4.2 Growth factors inhibit intrinsic apoptosis	33
4.3 Growth factors regulate metabolism	34
4.4 Growth factors maintain mitochondrial integrity	36
4.5 Regulation of the cell cycle is under control of growth factors	37
 Chapter 5 Scope of the thesis	39
 References	41

Part II:	Experimental data and results	62
Chapter 6	Growth factor starvation induces phosphorylation of glyoxalase I by a GSK-3β-dependent mechanism	61
	Introduction	62
	Materials and Methods	63
	Results	65
	Discussion	75
	References	78
Chapter 7	Glyoxalase I is a direct substrate for Ca²⁺/calmodulin Dependent Kinase II (CaMKII)	83
	Introduction	84
	Results and Discussion	85
	Conclusions	90
	Materials and Methods	91
	References	94
Chapter 8	Phosphorylation of glyoxalase I during the cell cycle	97
	Introduction	98
	Results	99
	Discussion	106
	Materials and Methods	108
	References	110
	Phosphorylation of glyoxalase I supplementary figures	114
Chapter 9	The Receptor for Advanced Glycation Endproducts (RAGE) is regulated at several levels during the cell cycle	117
	Introduction	118
	Results	119
	Discussion	130
	Materials and Methods	133
	References	136
Part III	General Conclusions	139
	General conclusions	140
	References	143
Part IV	Acknowledgements	145
	Dankuwel	146

Abbreviations

AG	aminoguanidine
AGE	advanced glycation endproduct
AGE-R	AGE-receptor
AIF	apoptosis-inducing factor
AMP	adenosine monophosphate
AMPK	AMP-activated protein kinase
Apaf-1	apoptotic protease activation factor 1
ASK1	apoptosis signal-regulating kinase 1
ATP	adenosine 5'-triphosphate
Bcl-2	B cell lymphoma 2
Bcl-x _L	B cell lymphoma-x _L
BH	Bcl-2 homology
CaMKII	Ca ²⁺ /calmodulin-dependent kinase II
CDK	cyclin dependent kinase
CDPK	calcium dependent protein kinase
C-t-Ab	C-terminus-directed anti-RAGE antibody
C-type	constant type
DHAP	dihydroxyacetonephosphate
Diablo	direct IAP-binding protein with low pI
DNA	deoxyribonucleic acid
DN RAGE	dominant negative RAGE
E.C.	enzyme commission
EGF	epidermal growth factor
ERK	extracellular signal-related kinase
G ₁	gap 1
G ₂	gap 2
G3P	glyceraldehyde-3-phosphate
GAPDH	glyceraldehyde-3-phosphate dehydrogenase
GF	growth factor
GLO1	glyoxalase I
GLO2	glyoxalase II
GLO3	glyoxalase III
Glut	glucose transporter

Abbreviations

GS	glycogen synthase
GSH	glutathione
GSK-3	glycogen synthase kinase-3
GSNO	S-nitrosoglutathione
GTP	guanosine-5'-triphosphate
HK	hexokinase
HLA-DR	human leukocyte antigen-DR
HMGB1	high mobility group box 1
HOG-MAP	high osmolarity glycerol-mitogen-activated protein
Hsp	heat-shock protein
HtrA2	high temperature requirement A2
HUVEC	human umbilical vein endothelial cell
IAP	inhibitory apoptosis protein
IL-3	interleukin-3
IL-7	interleukin-7
INK4	inhibitors of CDK4
IRS-1	insulin receptor substrate-1
JAK	Janus kinase
JNK	c-Jun N-terminal kinase
M	mitosis
MAPK	mitogen-activated protein kinase
MAPKK	mitogen-activated protein kinase kinase
MAPKKK	mitogen-activated protein kinase kinase kinase
MEK	mitogen extracellular kinase
MG	methylglyoxal
MG-AGEs	methylglyoxal-derived advanced glycation endproducts
mTOR	mammalian target of rapamycin
NAD	nicotinamide adenine dinucleotide
NADPH	nicotinamide adenine dinucleotide phosphate
NF- κ B	nuclear factor- κ B
NO	nitric oxide
O-GlcNAc	O-linked N-acetylglucosamine

Abbreviations

PDGF	platelet derived growth factor
PFK	phosphofructokinase
pI	isoelectric point
PI3K	phosphoinositol-3-kinase
PKA	protein kinase A
PKB	protein kinase B
pRb	retinoblastoma protein
PTK	protein-tyrosine kinase
R	restriction point
RAGE	receptor for advanced glycation endproducts
ROS	reactive oxygen species
rpS6	ribosomal protein S6
S	DNA synthesis
SAPK	stress-activated protein kinase
Smac	second mitochondria-derived activator of caspase
SnRK	SNF1-related protein kinase
sRAGE	soluble RAGE
STAT	signal transducer and activator of transcription
TNF	tumor necrosis factor
TP	triosephosphate
TPI	triosephosphate isomerase
VDAC	voltage-dependent anion channel
V-type	variable type

Summary

Whereas for unicellular organisms growth is controlled mainly by the availability of nutrients, this does not suffice for multicellular organisms. In multicellular organisms nutrients are abundant, so an additional regulatory mechanism is required. This regulatory mechanism is provided by the dependence on growth factors (GFs). These GFs are limited in the environment of the cells, and competition for GFs between cells determines the number, the size and the survival of these cells. Moreover, this competition defines only the fittest cells survive, and they survive only in the right place. GFs control several cellular processes, including the inhibition of the default apoptotic pathways, the rate of metabolism and cell division. Therefore, when GFs are depleted, metabolism collapses, cell division ceases and cell death is initiated.

Results from our group showed that phosphorylation of glyoxalase I (GLO1) is involved in tumor necrosis factor (TNF)-induced cell death. Phosphorylated GLO1 is directly or indirectly involved in modification of specific target proteins with methylglyoxal (MG), an unavoidable by-product of glycolysis. These events are a requisite for TNF-induced cell death and the data suggested that MG may function as a signaling molecule during regulation of cell death.

In this thesis we describe how GLO1 is phosphorylated upon withdrawal of GFs, which leads to a G_1 arrest and induction of apoptosis. GF depletion was also accompanied by MG-modification of specific target proteins. By pharmacological inhibition we could show that both events were GSK-3 β -dependent. Inhibition of GSK-3 β also significantly inhibited apoptosis induced by GF depletion. Next, we determined for which kinase GLO1 was a direct substrate. It turned out to be not GSK-3 β , but by an *in vitro* kinase assay, we found that GLO1 was a direct substrate of Ca^{2+} /calmodulin-dependent kinase II (CaMKII). This was later confirmed in cells by coexpression of CaMKII with GLO1.

In addition we found that phosphorylation of GLO1 also occurs during cell division by a PI3K-dependent mechanism.

Given the possibility that effects of MG-modified proteins can be mediated by the Receptor for Advanced Glycation End products (RAGE), we subsequently set off a study to investigate whether RAGE was regulated during the cell cycle and after GF withdrawal-induced G_1 arrest and apoptosis. We discovered a novel 17 kDa isoform of RAGE. This 17 kDa RAGE isoform was generated by proteolytic cleavage that could be inhibited by the general caspase inhibitor z-VAD-FMK. Processing of RAGE may thus represent an additional regulatory mechanism in RAGE signaling.

Samenvatting

Waar bij eencellige organismen de groei voornamelijk bepaald wordt door de beschikbaarheid van nutriënten, is dit niet afdoende bij meercellige organismen. Bij meercellige organismen zijn nutriënten doorgaans abundant aanwezig in het milieu van cellen. Bijgevolg is er een bijkomend regulerend mechanisme vereist. Dit regulerend mechanisme wordt voorzien in de afhankelijkheid van groeifactoren (GFs). Deze GFs zijn slechts in beperkte mate aanwezig in het milieu van de cel, en de concurrentie voor deze GFs bepaalt het aantal, de grootte en de overleving van cellen. Bovendien zorgt deze concurrentie ervoor dat enkel de gezondste cellen zullen prolifereren en dat cellen van een bepaald type enkel op de juiste plaats zullen overleven. GFs sturen verscheidene cellulaire processen, waaronder het verhinderen van de intrinsieke apoptotische signalisatie, het in stand houden van het cellulaire metabolisme en het reguleren van de celdeling. Bijgevolg, zal, wanneer GFs onttrokken worden, het cellulaire metabolisme vervallen, de celdeling stilvallen en celdood in gang worden gezet.

Resultaten verworven in onze onderzoeksgroep hebben aangetoond dat fosforylering van glyoxalase I (GLO1) betrokken is in tumor necrose factor (TNF)-geïnduceerde celdood. Gefosforyleerd GLO1 is vervolgens rechtstreeks of onrechtstreeks betrokken bij de modificatie van specifieke doelwitproteïnen met methylglyoxal (MG), een onvermijdbaar bijproduct van de glycolyse. Deze gebeurtenissen zijn vereist bij TNF-geïnduceerde celdood en de data suggereren dat MG zou functioneren als een signaalmolecule bij de regulatie van celdood.

In deze scriptie beschrijven we hoe GLO1 ook wordt gefosforyleerd tijdens een groeistop in de G_1 fase van de celcyclus en de hierop volgende apoptose, welke worden veroorzaakt door het onttrekken van GFs. Door farmacologische inhibitie konden we aantonen dat deze gebeurtenissen afhankelijk waren van GSK-3 β . Inhibitie van GSK-3 β inhibeerde eveneens significant de inductie van apoptose door GF depletie. Vervolgens bepaalden we voor welk kinase GLO1 een rechtstreeks substraat vormt. Dit bleek niet GSK-3 β te zijn, maar door een *in vitro* kinase assay vonden we dat GLO1 een rechtstreeks substraat is van Ca^{2+} /calmodulin-afhankelijk kinase II (CaMKII). Dit werd later in cellen bevestigd door co-expressie van CaMKII met GLO1. Daarenboven vonden we dat fosforylering van GLO1 ook voorkomt tijdens de celcyclus, en dit op een PI3K-afhankelijke wijze.

Gezien de mogelijkheid dat de effecten van MG-gemodificeerde proteïnen worden gemedieerd door de receptor voor gevorderde glycatie eindproducten (RAGE), werd vervolgens een aanvang genomen met een studie naar de regulatie van RAGE tijdens de celcyclus en bij GF-onttrekking geïnduceerde groeistop in de G_1 fase en de daaropvolgende apoptose. We ontdekten een nieuwe 17 kDa isovorm van RAGE. Deze 17 kDa RAGE isovorm werd opgewekt door proteolytische verknippen, wat verhinderd kon worden door de caspase inhibitor z-VAD-FMK. Verknippen van RAGE zou bijgevolg een bijkomend regulerend mechanisme kunnen zijn in RAGE signalisatie.

Part I: General Introduction

Chapter 1: The glyoxalase system

1.1 Introduction

For almost a hundred years the glyoxalase (GLO) system has been known to exist in animals. The discovery dates back to 1913, when two research groups independently found enzymes that converted α -oxoaldehydes into corresponding α -hydroxyacids (1,2). From then on, research focussed on its putative role in glycolysis. However, when the findings of Embden, Meyerhof and colleagues led to the elucidation of the glycolysis (3,4), it turned out that the GLO system did not lie on the main route to glucose degradation. Thus the hypothesis for a role of the GLO system in glycolysis was abandoned. Later it was found that the endproduct of the glyoxalase system was D-lactate and not L-lactate, which is the endproduct of glycolysis (5). This was the definite proof that the GLO system was not involved in glycolysis and interest in the GLO system severely diminished.

The ubiquitous nature of the GLO system, nevertheless, suggests a fundamental and conserved role (6). In the 1960s Szent-Györgyi took an interest in the GLO system. From his research on cancer cells he proposed a role for glyoxalase I (GLO1) in the regulation of cell division (7-9). However, until today his hypothesis could not be proven since a mechanistic link between GLO1 and the regulation of the cell cycle could not be found.

1.2 Detoxification of α -oxoaldehydes

An important function of the GLO system is the detoxification of reactive α -oxoaldehydes, mainly methylglyoxal (MG). This reaction is dependent on reduced glutathione (GSH) (10) and results in the formation of the corresponding α -hydroxyacid (5).

The GLO system comprises of two enzymes: glyoxalase I (GLO1; E.C. 4.4.1.5) and glyoxalase II (GLO2; E.C. 3.1.2.6). α -oxoaldehydes react nonenzymatically with reduced GSH, thereby forming a hemithioacetal. GLO1 catalyses the conversion of this hemithioacetal to a S-D-hydroxyacylglutathione (11-14), which, in turn, is hydrolyzed to the corresponding α -hydroxyacid, hereby regenerating reduced GSH (15) (Figure 1.1). The committed step in this pathway is the reaction catalysed by GLO1 since, under physiological conditions, it is considered irreversible (13).

The GLO system has a broad substrate specificity for α -oxoaldehydes (16-18), but the reaction rates are inversely correlated to the hydrophobicity of the side chains of the




Figure 1.1. The glyoxalase system.

The GLO system comprises of 2 enzymes, GLO1 and GLO2. α -oxoaldehydes spontaneously react with GSH to form a hemithioacetal, which is converted to a S-D-hydroxyacylglutathione by GLO1. This S-D-hydroxyacylglutathione is subsequently hydrolyzed to the corresponding α -hydroxyacid, hereby regenerating reduced GSH. (R = CH₃ for MG).

substrate (19). MG is generally considered to be the primary physiological substrate for the glyoxalase system. Under physiological conditions the GLO system reaction rates for the catalysis of MG are near diffusion-controlled limits (20-22), making the GLO system very efficient, and keeping the intracellular levels of MG low. The efficiency of the detoxification of MG is primarily determined by the spontaneous formation of the hemithioacetal. In aqueous solutions at pH 7, MG appears in α -oxoaldehyde, but also in a mono- and dehydrated form. When physiological concentrations of GSH are present, the equilibrium is shifted towards the formation of the hemithioacetal, and MG is efficiently removed by the GLO system (23,24) (see also chapter 2: Methylglyoxal).

1.3 Glyoxalase I

As previously mentioned the first step in the detoxification process by the glyoxalase system is the conversion of the hemithioacetal, which is spontaneously formed by MG and GSH, into S-D-lactoylglutathione. This reaction is catalysed by GLO1 (Figure 1.2). The crystal structure of GLO1 shows that the active site is situated in a barrel which is only formed at the dimer interface (25). This active site comprises a Zn²⁺ ion interacting with side chains from both subunits. The side chains involved are Glu₃₃ and Glu₉₉ from one subunit and His₁₂₆ and Glu₁₇₂ from the other. The structure of the active site suggests the mechanism is a proton transfer between the C-1 and C-2 of the hemithioacetal, whereby the ene-diol intermediate is stabilized in the *cis* configuration with both oxygens orientated towards the Zn²⁺ ion (25-28). The intermediate is rapidly ketonized to the hemithioacetal. It has been proposed that Glu₁₇₂ and Glu₉₉ are the catalytic bases for the S-substrate and R-substrate respectively (29-31).

GLO1 has been described to exist in various organisms, going from prokaryotes over yeast to plants, mammals and humans (e.g. (11,20,32-42)). In mammals and higher eukaryotes GLO1 functions as a dimeric metalloenzyme, using Zn²⁺ as a cofactor (see




Figure 1.2. Proposed catalytic mechanism for glyoxalase I.

The active site of GLO1 is situated at the dimer interface and contains a Zn^{2+} ion which interacts with side chains from both monomers (designated A and B). The substrate is the R-hemithioacetal and Glu₉₉ initiates the reaction as catalytic base. See text for more details (Adaptation from (27,30)).

above). GLO1 from *Escherichia coli*, however, needs Ni^{2+} for its activation (36). In yeast GLO1 functions as a monomer and contains two active sites (43). Monomeric human GLO1 is a protein of 184 amino acids with a molecular mass of approximately 21 kDa. It has 51% homology at the nucleotide level and 42% homology at the amino acid level with bacterial GLO1 (44). The GLO1 dimer is extremely stable and shows a high resistance to reducing conditions (45). GLO1 is present in a concentration of approximately 0,2 $\mu\text{g}/\text{mg}$ protein in most human tissue (46).

The human GLO1 gene is located at chromosome 6 (47) and its locus is linked to the HLA-DR (human leukocyte antigen-DR) (48,49). The gene for GLO1 was first described as one with five exons (45,50), but recent analysis indicates that it contains six exons (51). Human GLO1 is expressed from an autosomal diallelic gene and thus is expressed in three different phenotypes, GLO1 1-1, GLO1 2-2 and GLO1 1-2, representing the two homozygous and heterozygous expression patterns that are characteristic for co-dominant expression (52). GLO1 1-1 is electrophoretically slower than GLO1 2-2 (53). The difference in the two alleles is a single nucleotide polymorphism, resulting in an amino

acid substitution at position 111 from an alanine to a glutamic acid, thereby changing the electrophoretic properties (54). This variation also results in a decrease in glyoxalase activity (55). Several nucleotide polymorphisms have been described in the 5' and 3' untranslated regions and these polymorphisms may affect GLO1 expression, stability or posttranslational regulation. This may also affect the function of GLO1 (51). It has been claimed that some GLO1 phenotypes show correlation with diseases such as autism (55), multiple sclerosis (56), diabetes (57-59) and prostate cancer (60).

1.4 Posttranslational modification of glyoxalase I

There are several reports on posttranslational modification of GLO1. Studies in plants, yeast and mammals show that GLO1 is subject to phosphorylation and responds to nitric oxide (NO), indicating that posttranslational modification is conserved and is important for its biological function.

Phosphorylation: phosphorylation has been described to occur in mammalian cells, in yeast and in plants. Phosphorylation of GLO1 in mammalian L929 cells has been described as a necessity for the induction of necrosis by tumor necrosis factor (TNF) (61). This phosphorylation doesn't alter the detoxification capacity of the glyoxalase system for MG.

In yeast, treatment of the α -type budding strain of *Saccharomyces cerevisiae* with mating factor also induces phosphorylation of GLO1 (62). In contrast to the phosphorylation in mammalian cells, in yeast the activity of GLO1 was enhanced (63).

In *Arabidopsis*, GLO1 was phosphorylated upon the over expression of the kinase SnRK2.8, a kinase involved in resistance to stress (64). Concomitantly, a rise in GLO1 activity was induced. In rice leaves, gibberellin, a regulator of growth, induces phosphorylation of GLO1 (65), albeit this is a novel type of plant GLO1 (41). Possible changes in GLO1 activity were not reported.

Response to NO: research in yeast and mammalian cells indicates that GLO1 is responsive to NO. In mammalian endothelial cells the addition of NO leads to the inactivation of GLO1 (66). The addition of NO also leads to an increase of the isoelectric point (pI) of GLO1. The effects induced by NO on GLO1 are dependent on the formation of S-nitrosoglutathione (GSNO), suggesting that the substrate recognition site of GLO1 is most likely involved (67). Later it was shown that at least three cystein residues were

sensitive to NO-induced modification (68). Also in yeast and *in vitro* the addition of GSNO inhibits the activity of GLO1 (69).

Interplay between phosphorylation and NO has also been shown. In the TNF-induced necrosis in L929 cells NO-induced modification and phosphorylation occur independently. The NO-mediated modification is not required for the TNF-induced phosphorylation, although TNF-induced phosphorylation does primarily occur on the NO-responsive form of GLO1 (68).

1.5 Glyoxalase II

The second step in the detoxification of α -oxoaldehydes is the hydrolysis of the S-D-hydroxyacylglutathione formed in the first step (see above) to an α -hydroxyacid, hereby regenerating reduced GSH. Although GLO2 is a very efficient enzyme for the hydrolysis of S-D-lactoylglutathione (70), which is considered its physiological and preferred substrate, it is considered to be the rate limiting enzyme of the glyoxalase system (71). GLO2 is a monomer, consisting of two domains. The active site extends over the domain interface and contains a binuclear Zn-binding site. The active site also contains a water molecule, presumably in the form of a hydroxide ion. This hydroxide ion probably acts as the nucleophile catalysing the hydrolysis (72), whereby the Tyr¹⁷⁵ residue contributes to the binding of the GSH derivative (73).

Like GLO1, GLO2 has been described in various organisms, from prokaryotes over plants to mammals (74-82). GLO2 exists as multiple isozymes and has been located not only in the cytosol, but also in mitochondria. The human gene for GLO2 has been assigned to chromosome 16 (83) and some polymorphism has been described (84).

In mammals, the GLO2 gene gives rise to two distinct mRNA species transcribed from 9 and 10 exons respectively. The former of the two encodes for two protein species, mitochondrial targeted GLO2 and cytosolic GLO2, both initiated from different AUG codons. The latter mRNA species only encodes for cytosolic GLO2 (85). The mitochondrial GLO2 is located in the matrix and intermembrane space (86).

In plants and yeasts however, cytosolic and mitochondrial GLO2 are encoded by different genes (76,87). Multiple forms of mitochondrial GLO2 exist in plants and GLO2 can be located in the matrix and in the intermembrane space (86). Some mitochondrial isozymes have a Fe³⁺Zn²⁺ metal center instead of the two Zn²⁺ ions described for human

GLO2 (88). In *Arabidopsis thaliana* the different GLO2 isozymes are differently expressed in plants, with cytosolic GLO2 being more abundant in flower buds and mitochondrial GLO2 being more abundant in roots (76).

Quite interesting is the fact that GLO2 appears to be regulated by members of the p53 family of proteins. In the temporal cortex of p53^{-/-} mice GLO2 is decreased (89). Xu and Chen report that GLO2 is up-regulated by p63 and p73, two members of the p53 family that have fundamental roles in development (90). They postulate that GLO2 serves as a pro-survival factor and plays a critical role in the normal development and in the pathogenesis of various human diseases (e.g. cancer and diabetes).

1.6 Glyoxalase III

In *Escherichia coli*, the existence of a novel GLO enzyme, glyoxalase III (GLO3), has been described, which catalyses the conversion of MG to D-lactate (91). GLO3 works independently of the other two glyoxalases and in contrast to GLO1 and GLO2, does not require GSH as a co-factor. Specificity of this enzyme is limited to MG and phenylglyoxal. GLO3 is inactivated by oxidative stress (92). Besides the fact that GLO3 seems to play a role in the survival of non-growing *Escherichia coli* cultures (93), not much is known about this enzyme.

1.7 Possible functions of the glyoxalase system

It is quite obvious that the detoxification of α -oxoaldehydes, MG in particular, is a very important function for the glyoxalase system. However, the complex nature of the glyoxalase system (e.g. the fact that it comprises two enzymes, and not just one), the ubiquitous and conserved nature of the glyoxalase system, the existence of several other detoxification mechanisms for MG (see also the chapter 2: Methylglyoxal) and the existence of MG synthases in eukaryotes and prokaryotes (94-97) suggest that there are other, perhaps more fundamental functions for the glyoxalase system. Moreover, because of the different levels of regulation and modification of the different enzymes it is unlikely that the only role for the glyoxalase system is detoxification of MG. Two possible functions that will be emphasised here are an involvement in cell division and a role during stress.

A possible function in cell division: In plants it was observed that there was a strong correlation between GLO1 activity and cell proliferation (98,99). The observation

was made that in strongly dividing parts of the plant the concentration of MG was low and the GLO1 activity was high. In non-dividing parts, the level of MG was higher and GLO1 activity was undetectable. Also, when the cell cycle was inhibited, a decrease in GLO1 activity was induced and when GLO1 was inhibited, a significant inhibition of the increase in cell number and DNA content were observed. When soybean cell suspensions or carrot cell suspensions were treated with auxin, the activity of GLO1 was modulated together with proliferation (100,101). Gibberellin, another important growth regulator in plants, induces phosphorylation of GLO1 (65).

In yeast GLO1 activity correlates with the overall growth status, although it is not involved in the regulation of cell cycle (102). Furthermore GLO1 activity alters and is phosphorylated after treatment with mating factor (62,63). Mating factors are known to induce G₁ arrest in haploid yeast cells (103,104).

In addition, in animals a correlation between cell proliferation and activity of the glyoxalase system was found. Principato et al. found that GLO1 activity was high in embryonic chicken tissue, and GLO2 activity was low. This relation was inversed in mature chicken tissues, where a low GLO1 activity and a high GLO2 activity were observed (105). Also, in regenerating liver after partial hepatectomy, high GLO1 activity and low GLO2 activity were found. These activities returned to control levels after regeneration (106). During the maturation of human erythrocytes it was found that GLO1 activity increased, but a decreased activity of GLO1 was seen in mature, old erythrocytes (107). A very strong correlation was also found in the development of *Bufo bufo* embryos. Already at the earliest stages of development significant levels of GLO1 activity could be measured, and GLO1 activity rose during development. GLO1 to GLO2 activity ratios are very high at the beginning of development and gradually decrease as the development progresses (108). When differentiation is induced in HL-60 and K526, both leukaemia cell lines, a concomitant decrease in GLO1 to GLO2 activity ratios is seen (71). High expression levels of GLO1 are considered a risk factor for the development of certain types of cancer, and numerous tumours show a significant increase in levels and activity of GLO1 and a high GLO1 to GLO2 activity ratio in comparison to corresponding non-tumour tissues and surrounding tissues (109-117). Albert Szent-Györgyi was a strong believer in a very fundamental role for GLO1 and its substrate MG in the regulation of cell division and built his 'promine/retine' theory around it (e.g.(7-9,118,119) reviewed in (120)). However, in general the glyoxalase system is considered to be a 'marker for cell growth and division', but a causal relationship between the glyoxalase system and cell growth remains obscure.

A possible function in stress tolerance: A firm link between expression levels and activity of GLO1 and the resistance to abiotic stress has been established in plants. In *Brassica juncea* it was found that GLO1 was upregulated in response to salt, water and

heavy metal stresses (40). Moreover, in tomato plants an induction of GLO1 and GLO2 was observed in response to several forms of stress (121). Transgenic plants over-expressing GLO1 also showed an increase in tolerance towards salt and MG. Research in several plant species showed that in response to salinity, cold and drought, MG concentrations in the plant increase, which results in the inhibition of seed germination. This effect was enhanced by the down-regulation of GLO1 (122). Over-expressing components of the glyoxalase system led to enhanced tolerance towards salinity or heavy metal stress in several plant species (123-126). Furthermore it was observed that overexpression of SnRK2.8, a kinase involved in resistance to several types of stress in plants, induces phosphorylation of GLO1 (64).

In yeast, osmotic stress leads to alteration in GLO1 levels or activity. Inoue and co-workers reported the induction of GLO1 gene expression in *Saccharomyces cerevisiae* in response to osmotic stress (127). GLO1 activity is regulated in *Schizosaccharomyces pombe* in response to osmotic stress, as well. No change in mRNA levels or protein levels was observed in this case, so the effect was induced by posttranslational modification of GLO1 (128). In *Candida albicans* the regulation of GLO1 by the transcription factor Cap1p is part of the stress response induced by oxidative stress (129).

In animals, as well, the GLO1 is involved in resistance to different kinds of stress. Expression of GLO1 is up-regulated in the parasitic nematode *Onchocerca volvulus* under conditions of oxidative stress (37). Miller et al. showed that GLO1 provides protection of human retinal pericytes against hyperglycemia-induced cell death (130). Overexpression of GLO1 in bovine endothelial cells protected against the effects of hyperglycemia (131). It has been reported that tumors with high levels of GLO1 are more resistant to antitumor induced-apoptosis (132). Therefore a lot of effort has been put into finding inhibitors for GLO1 to be used as antitumor agents (e.g. (133-136)).

Regulation of the glyoxalase system may be a requirement for organisms to meet altered metabolic demands imposed by stress. In a sense, cell division can be regarded as a form of stress and the regulation of the glyoxalase system may be a consequence of adaptation to that stress. Indications of this, are, for instance, the involvement of similar signaling pathways that are involved in both stress response and cell division (e.g. Ca^{2+} (136,137) and MAPK pathways (127)).

Chapter 2: Methylglyoxal

2.1 Introduction

Methylglyoxal (2-oxopropanal, MG) has long been thought of simply as a by-product of glycolysis, which is the most fundamental metabolic pathway. Recent research, however indicates that there might be more to this molecule than just a role as cytotoxic by-product.

MG is a reactive dicarbonyl compound known to form covalent adducts with proteins and nucleic acids (138). In aqueous solutions it is present in three forms, the unhydrated, monohydrate and dihydrate forms, which are in equilibrium. The presence of physiological concentrations of glutathione (GSH) shifts this equilibrium towards the formation of the hemithioacetal, and MG is efficiently removed by the glyoxalase system (Figure 2.1) (23,24,139).




Figure 2.1. Methylglyoxal in aqueous solutions.

The main sources of MG are triose-phosphate (TP) glycolytic intermediates. Non-enzymatic fragmentation and elimination of phosphate from glyceraldehyde-3-phosphate (G3P) and dihydroxyacetonephosphate (DHAP) (140,141) or leakage of phosphor-enediolate from the active site of triosephosphate isomerase (TPI) (142,143) contribute most to the formation of MG (Figure 2.2). Metabolism of acetone and threonine also adds to the formation of MG (144-146), but under normal metabolic conditions both the latter pathways can be considered to be of minor importance. MG can also be formed enzymatically by methylglyoxal synthases (94-97,147-149), although the importance of these enzymes has not been further investigated in higher eukaryotes and therefore remains obscure.

An important regulator of intracellular MG levels is the glyceraldehyde-3-phosphate dehydrogenase (GAPDH). Reduced activity of GAPDH leads to the accumulation of TPs, leading to a subsequent increase in MG formation (150). GAPDH activity is influenced by e.g. oxidative stress (151) and by MG modification (152). This suggests that MG is able to propagate its own formation.

Cells contain considerable amounts of MG. The majority of the MG present *in vivo* however, is bound to proteins (see also below). Chaplen and colleagues detected up to 310 μ M of MG in chinese hamster ovary cells and estimate that 5-10% of cellular proteins may be modified to physiologically significant levels (153).




Figure 2.2. Main metabolic origins of methylglyoxal in mammalian cells. MG is mainly formed by non-enzymatic or enzymatic elimination of phosphate from triose-phosphate intermediates from glycolysis. Also acetone and threonine metabolism contributes to MG formation in mammalian cells, albeit considerably less.


2.2 Modification of proteins

MG is able to modify proteins under physiological conditions (154). The reactions proceed even at physiological concentrations of MG and lead to the reversible and irreversible modification of arginine and lysine residues and to the reversible modification of cysteine residues (Figure 2.3). However, when compared to cysteine residues, there is a clear preference for arginine and lysine residues to be modified (155). Although from model studies with $N\alpha$ -acetylamino acids the reaction of MG with protein is expected to be the most rapid at cysteine residues, *in vivo* MG-modified cysteines have not yet been detected, most likely because of their unstable and reversible nature (154). The modification of cysteine

with MG yields a hemithioacetal (156) (Figure 2.3.A). An example of such a reaction would be the reversible and non-enzymatic reaction between MG and GSH.


Under physiological conditions the primary targets are the arginine residues (157). A major physiological glycation adduct formed from MG with arginine is the imidazolone N_δ -(5-hydro-5-methyl-4-imidazolone-2-yl)-ornithine (Figure 2.3.B) (158,159).

A.



cysteine residue hemithioacetal

B.



arginine residue

N_δ -(5-hydro-5-methyl-4-imidazolone-2-yl)

Figure 2.3. The reaction of methylglyoxal with cysteine and arginine residues.

A. Reaction of MG with a cysteine residue (e.g. glutathione). **B.** Reaction of MG with an arginine residue. R = peptide chain.

Further irreversible reactions of the MG-modified arginine and lysine residues lead to the formation of Advanced Glycation Endproducts (AGEs). This reaction is slow and is called the maillard reaction. Formation of these AGEs is often accompanied by oxidative stress and leads to the dysfunctioning of protein and cells. AGEs contribute to the development of numerous pathophysiological conditions, such as diabetes, Alzheimer's disease and atherosclerosis, and aging (for reviews see (158,160-164)).

Glycation of a protein by MG generally induces partial unfolding; this renders proteins instable and induces degradation. MG modification, therefore, might function as a signal for the degradation of protein (154,157).

Interestingly, MG modification displays a high specificity. MG modification is restricted to preferential proteins (165-167), as well as to specific residues of these target proteins (168-170).

2.3 Detoxification of methylglyoxal

Besides the glyoxalase system (see chapter 1: The glyoxalase system) several other pathways contribute to the detoxification of MG.

1. Aldose reductase is dependent on NADPH and catalyses reduction of a wide range of aldehydes; MG is reduced to acetol, or when MG forms the hemithioacetal with GSH, to lactaldehyde. Both can then be reduced to propanediol, also catalyzed by aldose reductase (171-173).
2. Betaine aldehyde dehydrogenase is NAD-dependent and oxidizes numerous aldehydes, among which, MG. The efficiency of MG oxidation is considerably low in comparison with betaine aldehyde, its physiological substrate (174).
3. 2-oxoaldehyde dehydrogenase catalyses the NADP-dependent oxidation of MG to pyruvate. This enzyme seems primarily of importance in the liver, but has been purified from erythrocytes, as well (175-178).

However, under normal physiological conditions the glyoxalase system is the most important pathway for MG detoxification (24,127,171,179). Expression and activity of the glyoxalase system decrease with age, leading to a rise in MG and, consequently, AGE formation (165-167,180,181).

2.4 Methylglyoxal as a cytostatic and cytotoxic molecule

Ever since Szent-Györgyi hypothesized a function of MG as a regulator of cell division ((7-9,118), reviewed in (120)), a lot of attention has been given to this proposed role of MG.

MG exerts its activity on proliferation and survival in different ways:

1. MG signals are mediated by receptors (see chapter on RAGE)
2. MG and S-D-lactoylglutathione, an intermediate in the detoxification of MG, are capable of inhibiting DNA-synthesis and inducing DNA-damage, leading to G₁ arrest and subsequently to apoptosis (182-189).
3. Reactive oxygen species (ROS) are generated during the reaction of MG with amino acid residues (190-193). Cell lines that are more sensitive to oxidative stress are therefore

also more sensitive to the effects of MG (194,195). A first effect of high concentrations of ROS and MG is a depletion of cellular GSH, hereby impairing protection against oxidative stress and detoxification in general (194,196-199). Superoxide formation also leads to the activation of p38 MAPK and JNK, signaling cascades leading to apoptosis (195,200-202) (see also paragraph 'MG as a signaling molecule').

4. In human embryonic kidney cells (HEK293) and a mouse fibroblast cell line (NIH3T3) MG has been shown to render these cells resistant to the mitogenic action of insulin-like growth factor-1 (IGF-1). This effect was associated with a stronger and prolonged activation of ERK (extracellular signal-related kinase) and an up-regulation of p21 (Waf1/Cip1) (203).

5. The impairment of insulin signaling by MG is well documented. MG reacts with insulin, hereby changing the biological functions of insulin (204). In addition, downstream effector proteins of the insulin receptor are modified, impinging on insulin signaling (205). Other receptors, such as the platelet derived growth factor (PDGF) receptor and epidermal growth factor (EGF) receptor, are also inhibited in their signaling by MG (206-208).

6. Glycolysis and mitochondrial respiration are sensitive to MG. Tumor cells seem particularly sensitive to these inhibitory effects (209-215). This inhibition is mainly the consequence of inactivation of GAPDH, but also of hexokinase (HK). At the mitochondrial level, the inhibition occurs at complex I of the respiratory electron transport chain.

7. The covalent modification of proteins by MG plays a role in cell survival, as well. Two examples are the modification of heat-shock protein 27 (Hsp27) (169,216-219) and of the mitochondrial permeability transition pore (220,221). In both cases MG reversibly modifies arginine residues, hereby influencing the structure and function of the involved protein. These MG modifications protect cells against apoptotic stimuli, somewhat in contrast to the other examples, where MG induces cell cycle arrest in the G₁ phase or apoptosis.

2.5 Methylglyoxal in other signaling pathways

Despite its evident cytostatic and cytotoxic properties, MG is inevitably present in every cell under physiological conditions, from prokaryotes to higher eukaryotes. Not only is it a by-product of glycolysis, the most fundamental metabolic pathway, several other metabolic pathways lead to the formation of MG in microorganisms and higher eukaryotes, as well. These facts, together with the well-documented existence of prokaryotic as well as eukaryotic MG synthases, make the explanation of MG as a mere flaw of metabolism unsatisfying. Recent reports suggest a possible role for MG as a signaling molecule.

In plants: Under stress conditions a rise in intracellular MG concentration has been observed (122). This seems to be a general mechanism, as a rise of intracellular MG concentrations has been measured in several different plant systems. This suggests that MG, therefore, could act as a signal for plants to respond to stress.

In yeast: In *Saccharomyces cerevisiae* it has been shown that the activity of Yap1, a transcription factor that mediates the oxidative stress response, is modulated by methylglyoxal (222). The localisation of Yap1 in the cytoplasm or the nucleus is influenced by intracellular concentrations of MG. Yap1 is most likely directly modified by MG, presumably on a cysteine residue. This modification is reversible and independent of ROS formation in the cell. It has also been reported that extracellular MG activates several signal transduction pathways, like the high osmolarity glycerol (HOG)-mitogen-activated protein (MAP) kinase pathway, hereby eliciting the induction of several genes (223,224). In addition, MG transiently activates the uptake of Ca^{2+} from the medium (224). Both responses are independent of ROS formation, neither are they caused by changes in osmolarity due to the higher concentration of extracellular MG. Maeta and colleagues hypothesize there might exist a sensory system for external MG on the cell surface conveying signals to the nucleus. A likely candidate for this is the osmosensor Sln1.

In *Schizosaccharomyces pombe* an alternative mechanism has been shown to exist. Here extracellular MG activated a yeast homologue of the stress-activated protein kinase (SAPK) pathway, leading to activation of the transcription factor Spc1 and independent of this Spc1-SAPK pathway Pap1, a Yap1 homologue in *S. pombe*, is activated, as well (170). The activation of Spc1 through the SAPK pathway takes place irrespective of intracellular MG content. Activation of Pap1, on the other hand, is reversible and dependent on intracellular MG. ROS are not involved in either of the pathways. MG could, therefore, be a metabolic signaling molecule. Intracellular levels of MG vary depending on growth conditions and stress situations. These alterations in MG concentrations could then elicit a response through the modification of different signaling molecules (225,226).

In mammals: In mammalian cells the pathways involved in the mediation of MG-induced signals and the outcome are dependent on the cell type. MG selectively induces apoptosis in Jurkat cells (195,201). The mechanism of MG-induced apoptosis is cell type-dependent since several other cell lines were shown to be insensitive to MG. MG-induced cell death was mediated through JNK (c-Jun N-terminal kinase), showed typical apoptotic morphology (e.g. DNA fragmentation) and was dependent on caspase-3. Administration of MG to rat mesangial cells induces apoptosis. This apoptosis is dependent on the activation of apoptosis signal-regulating kinase 1 (ASK1) through the

formation of ROS and is mediated by the p38 α MAPK pathway (202). Up-regulation of p38 MAPK pathway is also involved in the down regulation of glucose transporter-4 (GLUT4) in adipocytes (227). In human umbilical vein endothelial cells (HUVECs) MG triggers several pathways (200,228). A first pathway involves protein-tyrosine kinase (PTK) activation for tyrosine phosphorylation of several cellular proteins, leading to the activation of ERK. Another pathway triggered by MG is the induction of apoptosis. This signal transduction cascade is independent of PTK activity, but involves ROS induced activation of JNK, p38 kinase and c-Jun. Interestingly enough, glyoxal, which differs from MG only by a methyl group, is only capable of inducing the first signal, but unable to induce the second. This strongly accentuates the specificity of MG as a signaling molecule.

In retinal Müller cells and in microvascular endothelial cells from mouse kidney increased glycolysis leads to increased MG modification of the corepressor mSin3A. MG modification of mSin3A results in the recruitment of O-linked N-acetylglucosamine (O-GlcNAc) transferase to an mSin3A-SP3 complex, leading to the O-GlcNAc modification of Sp3. This modification results in a decreased DNA-binding capacity of the repressor complex, thus changing the gene expression pattern of the respective cells (229,230). The extent of this modification is a direct reflection of intracellular MG concentrations which is determined by a variety of cellular processes. This suggests a possible function for MG as an integrator to co-ordinately regulate distinct patterns of gene expression. In the mouse fibrosarcoma cell line L929 tumor necrosis factor (TNF)-induced necrosis is dependent on the accumulation of MG. The formation of specific MG-derived modified protein is observed and these steps are important in these signaling processes (61).

Chapter 3: RAGE, the receptor for advanced glycation endproducts

3.1 Introduction

One mechanism by which methylglyoxal-derived advanced glycation endproducts (MG-AGEs) exert their cellular function is through interaction with specific receptors. Various receptors and proteins that can bind to AGEs have been described, e.g. macrophage scavenger receptors type I and II (231-234), oligosaccharyl transferase-48 (AGE receptor 1; AGE-R1) (235,236), 80K-H phosphoprotein (AGE-R2) (235,236), galectin-3 (AGE-R3) (237), and lactoferrin and lysozyme (238). The one that has been studied the best, however, is the receptor for advanced glycation endproducts (RAGE) (239,240).

3.2 The receptor for AGEs

RAGE was first isolated from bovine lung extract by affinity purification using AGE-modified albumin (240). RAGE is a member of the immunoglobulin superfamily of cell surface receptors. It is composed of one V-type and two C-type immunoglobulin domains which are extracellular, a single-pass transmembrane spanning domain and a short highly charged intracellular domain (239,241). The principal ligand binding domain is the V domain (242), however, one ligand, namely S100A6 has been reported to interact specifically with the C2 domain (243), hereby emphasizing the complex nature of RAGE signaling. Furthermore, recent reports indicate that the V domain and the C1 domain do not function as independent domains, but as a single structural unit (244). The highly charged cytoplasmic tail is important for signal transduction (241).

At least three different major isoforms of RAGE exist, the full length RAGE (FLRAGE), an isoform described as a dominant negative RAGE (DNRAGE), lacking the intracellular domain, and a soluble RAGE (sRAGE), lacking both the intracellular domain and the transmembrane domain (Figure 3.1). These isoforms are the result of alternative splicing from a single RAGE gene (245), although some isoforms may be generated by protein cleavage (246). Other splice variants and polymorphism have also been described (reviewed in (247)).

RAGE expression has been detected in all tissues and at a cellular level RAGE is found at the surface of most cell lines that have been examined (248). The correlation between mRNA levels, detected by Northern Blot, and protein levels seems small, indicating multiple levels of transcriptional, posttranscriptional and translational regulation. Of note is that RAGE expression is very low under physiological conditions, but under pathological




Figure 3.1. Major isoforms of RAGE.

Multiple isoforms of RAGE have been described. The three major isoforms are depicted and all contain an N-terminal V-type immunoglobulin domain and two C-type domains. sRAGE is not attached to the cell and has been hypothesized to attenuate RAGE signaling by capturing ligands prior to membrane interaction. DNRAGE does not possess a signaling function since it lacks the intracellular domain and therefore acts as a decoy receptor. sRAGE and DNRAGE could however exert a signaling function when involved in receptor complex formation. RAGE signaling is in part determined by the orchestrated effects of the different RAGE receptors. Adapted from (247).

conditions RAGE expression is strongly upregulated (249-252).

3.3 Signaling through RAGE

A striking feature of RAGE signaling is its multi-ligand character. These ligands include AGEs (242,253,254), amyloid- β peptide ($A\beta$) and β -sheet fibrils (255,256), several members of the S100/calgranulin family (257), amphoterin (often referred to as high mobility group box I (HMGB1) DNA-binding protein) (258,259), and Mac-1, a member of the β 2-integrins (260). Multiple cellular signaling cascades have been shown to be activated by the binding of ligands to RAGE (Figure 3.2). The first signaling pathway to be described was the activation of p21^{ras} through the induction of reactive oxygen species (ROS), leading to a cascade of mitogen-activated protein kinases (MAPKs) (261,262). This pathway has been reported in several cell types (263-266). The cytoplasmic tail of RAGE could function as a docking site for the extracellular signal-related kinase 1/2 (ERK1/2) MAPK (267). Other signaling pathways induced by the ligand binding of RAGE are other MAPK pathways, such as the p38 pathway and the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) pathway (266,268-270), several JAK/STAT (Janus kinase/

signal transducer and activator of transcription) pathways (271-273), the cdc42 and rac pathway of rho-GTPases (241), and the phosphoinositol-3-kinase (PI3K) pathway (274). Most of these pathways lead to activation of NF- κ B (nuclear factor- κ B) (e.g. (241,261,263-266,270,275,276)). Of note is that RAGE signaling hereby provides a positive feedback loop since the RAGE gene contains two NF- κ B responsive elements in its promoter (277,278).

The outcome of RAGE signaling is dependent on the cell type, the available ligands and their relative concentrations, the presence of additional splice variants and isoforms, the presence of RAGE polymorphisms and their different affinities for the available substrates, and the interplay between all of these different factors. An additional factor making RAGE signaling even more complex is the observation that some RAGE-mediated effects are independent of the cytoplasmic tail and only require the extracellular domain (284). This suggests engagement of RAGE in receptor complexes.




Figure 3.2. Schematic overview of signaling through RAGE.

Numerous signaling pathways can be initiated upon RAGE activation. Of note is the central role of NF- κ B. Other transcription factors such as c-Jun are also involved in RAGE-mediated signaling. RAGE activation can also result in apoptosis, independent of gene-induction by activated transcription factors. See text for more details (recently reviewed in (279-283)).

3.4 Involvement of RAGE in proliferation and survival

Engagement of RAGE by one of its ligands can give rise to the activation of numerous key cell signaling pathways (see also above). The outcome depends on several parameters (e.g. cell type, ligand, ligand concentration, ...). Several of these signaling cascades are important in the regulation of proliferation and cell survival, two processes of which the molecular pathways are closely intertwined.

Although several reports implicate the engagement of RAGE in the induction of cell division, RAGE signaling is also involved in differentiation. Proliferation and differentiation are separate and mutually exclusive events that must occur sequentially and in that order. Sajital and co-workers provided evidence for a role of amphoterin/RAGE signaling in growth retardation and survival of neuroblastoma cells when differentiation was induced with retinoic acid. RAGE was shown to play a significant role in cell survival by inducing an increase of the anti-apoptotic protein Bcl-2 (285). In rat L6 myoblasts amphoterin/RAGE interaction also stimulates differentiation and inhibits proliferation through the activation of the p38 pathway (286,287). Amphoterin activates RAGE partly in an autocrine manner in order to exert its activity on myoblasts (288).

When compared to normal tissue, a strong reduction in RAGE mRNA and even more in RAGE protein has been observed in lung cancer cells (289,290). This suggests that RAGE exerts an inhibitory activity on lung cell proliferation and promotes differentiation. Indeed, when full-length RAGE is over-expressed, proliferation is impaired and cells are rendered less responsive to proliferative stimuli. Activity of the p44/p42 MAPK pathway is decreased upon RAGE overexpression and the cells start to differentiate. A more direct link between RAGE and cell cycle control was seen in mesangial cells. Binding of AGE to RAGE leads to the activation of STAT5, which induces the p21^{waf} gene, a potent inhibitor of cyclin dependent kinases (CDKs), key regulators of cell cycle (271).

RAGE, however, has also been described to promote proliferation. Taguchi and colleagues noted amphoterin/RAGE interaction induces lung tumor proliferation through activation of p44/p42, p38 and SAP/JNK MAP kinase pathways (268). Furthermore, RAGE stimulation with S100B or AGEs could function as a mitogenic stimulus in neuronal cells. Ligand binding of RAGE leads to the activation of the p44/p42 MAPK pathway and results in the upregulation and activation of cyclin D₁/cdk4 and the transition into the S-phase (291).

Of special interest is the complex nature of the involvement of amphoterin/RAGE interaction in T cell activation driven by dendritic cells. In response to inflammatory stimuli, dendritic cells actively secrete amphoterin. This amphoterin is extensively acetylated and serves as an autocrine ligand for RAGE activation, leading to a well-characterized differentiation program, referred to as maturation, through activation of p38, ERK1/2 (p44/p42) and NF- κ B. However, the secreted amphoterin also drives proliferation, survival and

polarization of naïve T cells through interaction with RAGE (292,293).

RAGE signaling has been shown to be involved in cell survival. Also here, RAGE exerts a very dualistic function, as it has been implicated in both the survival of cells as in the induction of cell death, sometimes even in the same cell type. Activation of RAGE by amphotericin can promote cell survival by inducing expression of Bcl-2 (B cell lymphoma-2) through activation of NF- κ B (294). These effects, however, are seen when nanomolar concentrations of amphotericin are present. When, on the other hand, micromolar concentrations of amphotericin are present, apoptosis was induced. The latter outcome was dependent on the induction of ROS, which led to the activation of mitogen extracellular kinase (MEK) and consequent cytochrome c release from mitochondria and caspase-3 activation. Furthermore, when S100P activated RAGE in an autocrine manner, ERK mediated NF- κ B activation leads to proliferation and survival (295,296). This dual role for RAGE is also observed in neuronal cells, where low concentrations of S100B protect cells against hypoxia or ischemia-induced apoptosis, whereas high concentrations of S100B exert a RAGE-mediated cytotoxic effect (249). On the other hand, AGE-modified collagen induces caspase-3, -8, and, to a lesser extent, -9 dependent apoptosis upon binding to RAGE (297,298).

Chapter 4: Growth factors

4.1 Introduction

Growth, proliferation and survival of cells are highly regulated processes and in multicellular organisms depend on the presence of extracellular stimuli (299). The availability of growth factors (GFs) seems decisive herein (300). These GFs control cell survival, either through the inhibition of the intrinsic apoptotic pathways or by actively promoting survival. Both mechanisms are strongly intertwined and rely partly on the same signaling pathways. Also, it is clear that neither of the mechanisms is sufficient on its own, both are needed to maintain cell size and viability (301-307). Literature on this subject is vast. A large number of cytokines exist, the web of downstream signaling pathways is complex, and the outcome is largely dependent on the cell type, making knowledge on this matter fragmentary. Valuable tools in the study of GF signaling are hematopoietic cells. For their survival, these cell lines depend on well defined GFs, such as interleukin-3 (IL-3), -2 or -7. A central mediator of interleukin signaling is the phosphoinositide-3-kinase (PI3K) and Akt/protein kinase B (PKB) pathway (308-313). A simplified schematic overview of PI3K/Akt signaling is given in figure 4.1. Akt is a versatile kinase with numerous substrates and Akt signaling contributes to diverse physiological processes, including the regulation of apoptosis, the regulation of the cell cycle and the regulation of glucose metabolism.




Figure 4.1. The PI3K/Akt pathway has a central role in growth factor signaling.

Binding of GFs to their receptors leads to activation of PI3K, either directly or indirectly through scaffolding adaptors such as insulin receptor substrate-1 (IRS-1). These events ultimately lead to the activation of Akt. Subsequently numerous substrates can be phosphorylated by Akt. This phosphorylation can lead to either activation (arrows) or inactivation (blocking arrows). Two major downstream branches in Akt signaling are the mTOR pathway, which gets

activated by Akt, and the GSK-3 pathway, which gets inactivated by Akt phosphorylation. Several downstream targets are shown, which are regulated directly by Akt or indirectly through the mTOR or GSK-3 branch. These targets illustrate how Akt signaling is involved in the regulation of different physiological processes, such as apoptosis, the cell cycle and glucose metabolism (see text for further discussion of these processes). Of note, both GSK-3 and mTOR signaling lead to inhibition of IRS-1, hereby negatively regulating Akt signaling. Recent reviews about PI3K/Akt signaling are (314,315), see text for more references.

4.2 Growth factors inhibit intrinsic apoptosis

In the absence of appropriate GFs cells, by default, undergo apoptosis (316) (Figure 4.2). Members of the BH3-only subfamily of the Bcl-2 family¹ are activated upon GF-withdrawal. Once activated, these BH3-only members promote apoptosis, most likely by antagonizing the prosurvival anti-apoptotic Bcl-2 family members such as Bcl-2, Bcl-x_L and Mcl-1 or by facilitating the activity of the pro-apoptotic Bcl-2 family members Bax and Bak (321,323-328). Bax and Bak subsequently participate in the permeabilization of the outer mitochondrial membrane, hereby releasing prodeath factors like cytochrome c, Smac, apoptosis-inducing factor (AIF), and endonuclease G. Release of cytochrome c then leads to the activation of caspase-9, which in turn activates caspase-3 (329-331). Numerous studies indicate that Bad is the BH3-only member involved in GF withdrawal-induced apoptosis (332-335). Recent reports, on the other hand, claim that, not Bad, but Puma or Bim are involved (336-339).

GF signaling can intersect with this default apoptosis at several levels. A critical step in the induction of apoptosis is the permeabilization of the outer mitochondrial membrane and the subsequent release of cytochrome c. An important manner in which GFs inhibit apoptosis is, therefore, upstream of this event, by directly hampering the action of the BH3-only proteins. Proteins like Bad, Bim and Puma are under tight transcriptional control and are regulated at the posttranslational level. The best studied example herein is Bad. GFs are able to induce multiple phosphorylations of Bad (309,342-346), hereby marking it for binding to 14-3-3 proteins (342,347-351) or blocking its interaction with Bcl-2 and Bcl-x_L (352-354) and inactivating its pro-apoptotic function. The same mechanism has been shown to be involved in the regulation of the activity of Bim (338). GFs also regulate the BH3-only proteins at the transcriptional level in part by inactivating the forkhead transcription factors (336-339,355-357). In addition, GFs counter apoptosis by inducing the expression of anti-apoptotic members of the Bcl-2 family, partly through the activation of NF-κB (358-366). It is interesting to note that GFs can also regulate this intrinsic apoptotic pathway downstream of cytochrome c release. This inhibition can be obtained in part by the direct phosphorylation of caspase-9 (367-369).

¹ The control and regulation of mitochondria-related apoptotic events occurs through members of the Bcl-2 (B cell lymphoma 2) family of proteins. Based on the presence of conserved Bcl-2 homology (BH) domains, and their functions, three subfamilies can be described (310,311).

The anti-apoptotic Bcl-2 proteins, such as Bcl-2, Bcl-x_L and Mcl-1, contain all four BH-domains. The pro-apoptotic proteins can be separated into two sub-families. First there is the multidomain Bcl-2 family, including Bax and Bak. This family shares three BH-domains, but lacks the N-terminal BH4 domain (312,313). And finally there is the pro-apoptotic BH3-only subfamily, containing only the BH3 domain. This family includes e.g. Bid, Bad, Bim, Puma, Noxa (314-318).




Figure 4.2. Growth Factor withdrawal leads to mitochondria-dependent apoptosis.

Upon the withdrawal of GFs Bax and Bak are activated by the BH3-only proteins (e.g. Bad, Bim, ...), leading to the permeabilization of the outer mitochondrial membrane and the subsequent release of cytochrome c and other prodeath factors from the intermembrane space. The precise mechanism of Bax/Bak activation is still unclear. Either BH-3 only proteins bind to and neutralize anti-apoptotic Bcl-2 proteins, leading to the release of Bax and Bak, or BH3-only proteins can interact directly with Bax and Bak. The released cytochrome c, together with Apaf-1 and caspase-9, forms the apoptosome. Smac/Diablo and Omi/HtrA2 relieve the inhibition of caspases induced by IAP. Subsequently the executioner caspase-3 is activated, which results in apoptosis. AIF, endonuclease G, Omi/HtrA2 and the dysfunctioning of the mitochondria further contribute to cell death in a caspase-independent manner (for reviews see (328,340,341)).

4.3 Growth factors regulate metabolism

Although the most obvious way for GFs to regulate survival is by inhibiting apoptosis (see above), GFs are dependent on the presence of nutrients to exert their function (306,370,371). A key indication for this is the observation that even the presence of Bcl-2 or the deficiency of pro-apoptotic Bcl-2 family members fails to rescue cells from cell death upon GF withdrawal (301,372-376). Instead, in the absence of GFs, cells undergo progressive atrophy, ultimately leading to cell death (304,371,377). It is becoming clear

that metabolism is not merely driven by increasing metabolic demands, due to GF-driven proliferation. GFs directly regulate metabolism, in parallel to growth, proliferation and survival (378,379). A decrease or inhibition of glucose metabolism can lead to the initiation of cell death and, inversely, GF withdrawal-induced cell death is associated with a reduced glucose metabolism (380).

A first key step in the regulation of glucose metabolism is the uptake of glucose through facilitative glucose transporters, such as Glut1, the predominant glucose transporter in hematopoietic cells (Figure 4.3). A strong correlation is seen between the ability of GFs to maintain viability and the expression of glucose transporters (380,381). It has also been shown that overexpression of Glut1 can delay the onset of GF starvation-induced apoptosis in IL-3 dependent cells, although this Glut1 overexpression does not maintain long-term survival (306). An important mechanism by which GFs regulate glucose uptake is by promoting localization and retention of Glut1 on the cell surface (379,382). GF withdrawal leads to a rapid decline in membrane expression of Glut1, which can be prevented by activation of PKB/Akt signalling (303,305,306,371,372,383). In addition to the regulation of glucose uptake, GFs have been shown to regulate the membrane expression of several other nutrient transporters, like 4F2hc, an amino acid transporter, transferrin receptor, and low-density lipoprotein (LDL) receptor in a PKB/Akt and mTOR-dependent fashion (384-386).




Figure 4.3. Schematic overview of the regulation of the metabolism by Growth Factors.

GLycolytic metabolism is regulated at several steps by GFs. Also the membrane localisation of numerous nutrient transporters is under the control of GFs. See text for more details (adapted from (378,387)).

Besides determining the uptake of glucose and metabolites, GFs strongly regulate glycolysis itself (Figure 4.3). Upon GF withdrawal, the flux of glucose through the glycolytic pathway declines and the production of L-lactate drops (303,306,388). Rate-determining steps in glycolysis are the steps catalyzed by hexokinase (HK) and phosphofructokinase 1 (PFK-1) and a decrease in reaction rate of either one of these enzymes results in a decrease in the overall rate of glycolysis. HK mediates the first committed step of glycolysis, the phosphorylation of glucose to glucose-6-phosphate. GF stimulation leads to an increase of HK expression levels and HK activity (306,389-392). Several reports have confirmed a general association between high levels of HK activity and cell survival (370,393,394). Furthermore, it has been described that overexpression of HK is able to attenuate apoptosis (305,370,395,396). The most important rate-limiting committed step in glycolysis is the conversion of fructose-6-phosphate to fructose-1,6-bisphosphate, catalyzed by PFK-1. Besides regulating the expression level (306,397), GFs also control the activity of PFK-2 (phosphofructokinase 2) (397-401). PFK-2 is activated by phosphorylation and this leads to an increase of fructose-2,6-bisphosphate, an important allosteric activator of PFK-1 (402). In addition, other enzymes involved in metabolism, such as lactate dehydrogenase (LDH) and triosephosphate isomerase, are regulated by GFs (389).

4.4 Growth factors maintain mitochondrial physiology

For a long time the maintenance of metabolism and the inhibition of apoptosis were considered to be independent pathways and were studied as such (306,309,331,352,353,370). A central point in the induction of apoptosis is the permeabilization of the outer membrane of the mitochondria, which leads to the release of cytochrome c and other pro-apoptotic proteins. A decrease in metabolism would lead to a perturbation of mitochondrial physiology, resulting in a subsequent rupture of the outer membrane (304,403,404). On the other hand, Bcl-x_L is capable of preventing apoptosis, even in the absence of glucose or GF stimulation (370,388), suggesting that the regulation of metabolism and induction of apoptosis can be dissociated. Several reports, however, point towards a role for HK in the regulation of mitochondria-dependent apoptosis. HKs are known to reside in mitochondria upon GF stimulation and to bind to the voltage-dependent anion channel (VDAC) (370,405). The VDAC is a β -barrel protein, spanning the outer mitochondrial membrane and has been implicated in the induction of apoptosis (406,407). Mitochondrial bound HK is able to attenuate the induction of apoptosis by antagonizing the activity of Bax and Bak (396,408,409). In liver mitochondria HK IV and Bad have been shown to reside in a holoenzyme complex (307). The phosphorylation status of Bad influences HK activity and HK activity is required in order to keep Bad phosphorylated. This mechanism provides strong evidence for the coordination of apoptosis and glycolysis and indicates an interdependence between pathways preventing apoptosis and pathways promoting survival.

4.5 Regulation of the cell cycle is under control of growth factors

Besides preventing apoptosis and stimulating metabolism (see above), GFs are known to exert another function, namely the induction of proliferation. Overexpression of Bcl-2 or Bcl-x_L can protect cells from GF withdrawal-induced apoptosis, but cells undergo growth arrest (404,410,411). Mammalian cells depend on GFs for the regulation of the transgression through the cell cycle (Figure 4.4). It has been known for a long time, however, that this dependency does not last throughout the entire cell cycle. Already during G₁ phase, several hours before the entry into S phase, the restriction point (R) is defined as the moment when a transition occurs from GF dependence to GF independence. After R, cells can complete their division cycle irrespective of the presence of mitogenic signals (412-414). Cell cycle progression relies on a highly regulated chain of signaling and transcriptional events, involving the sequential transcription and activation of cyclin dependent kinases (CDKs, reviewed in (415)). Important herein is the phosphorylation state of the retinoblastoma protein (pRb) (416). pRb binds to the E2F family of transcription factors (417,418) and to chromatin remodeling enzymes, such as histone deacetylase 1 and 2 (419,420), hereby inhibiting their activity and suppressing cell cycle progression. In early G₁ pRb is hypophosphorylated and exerts its inhibitory activity. In order to pass from G₁ to S phase the cyclin D-CDK4/6 complex partially phosphorylates pRb, causing the release of the E2F family of proteins. Many promoters of genes which are important for cell cycle progression, such as cyclins E (421) and A (422) are activated by E2F proteins. Subsequently pRb is further phosphorylated by the cyclin E-CDK2 complex, hereby intensifying the transcription of E2F-regulated genes, leading to G₁-S transition. Phosphorylation of pRb and the subsequent induction of cyclin E are often considered to be the molecular mechanism behind R, although this is contested by several reports (423-426).

GFs regulate cell cycle entry and G₁-S transition mainly by lowering the concentration of CDK inhibitors, consisting of the inhibitors of CDK4 (INK4) and the more broadly acting Cip/Kip families, and by regulating the levels and activity of cyclins and CDKs (415,427,428).

The withdrawal of GFs results in two distinguishable and sequential responses. The first response is a cell cycle arrest, which is followed by apoptosis. Moreover, not only do these processes differ kinetically and mechanistically, it has been shown that lower concentrations of GFs suffice to protect cells from apoptosis, but are not able to induce cell division (429). Of note is the observation that Bcl-2 family proteins have regulatory properties in cell cycle. Bcl-2 and Bcl-x_L are known to enhance G₀ arrest and significantly delay cell cycle progression, in part, by inducing p27^{Kip1} expression (430-434). These effects on cell cycle can not be separated from the prosurvival functions and can be

countered by Bad and Bax (435-437). Recent reports that pro-apoptotic proteins like Bad and caspase-9 are regulated in a cell cycle-dependent manner underline how much cell cycle and apoptotic signaling are interlaced (367,368,438). By combining the regulation of these processes, GFs might provide protection against inappropriate proliferation.




Figure 4.4. Schematic overview of the cell cycle.

Several phases can be distinguished in the cell cycle; G₁ (gap 1), S (DNA synthesis), G₂ (gap 2) and M (mitosis). Late in G₁, cells become independent of GFs to complete their division cycle. This point is referred to as the restriction point (R). Formation and activation of cyclin-cdk complexes determines the progression of the cell cycle and is strictly regulated throughout the different phases. The presence and activity of the respective complexes is indicated with arrows. A central role of the cyclin-cdk complexes is the phosphorylation of retinoblastoma protein (pRB), a general cell cycle inhibitor that prevents G₁-S transition by repressing E2F dependent transcription. The role of GFs is the regulation the levels and activity of G₁ cyclins and CDKs and lowering the levels of inhibitory proteins. Withdrawal of GFs renders cells incapable of passing R and results in G₁ arrest. See also text for more details.

Scope of the thesis

The requirement of growth factors (GFs) is of fundamental importance in physiological homeostasis. For unicellular organism growth regulation is mainly controlled by the availability of nutrients. However, for multicellular organism an additional regulatory mechanism is indispensable. Here nutrient availability is normally not limiting and thus does not suffice to control cellular growth and division in multicellular organisms. This additional regulatory mechanism is provided by the dependence on GFs. GFs exert their activity by regulating several fundamental processes, such as metabolism, proliferation and survival. Although these processes are regulated as independent branches of GF-induced signaling, extensive crosstalk is needed between them.

In this context we set out to study the involvement of the ubiquitous and highly conserved glyoxalase system and its physiological substrate methylglyoxal, an unavoidable by-product of glucose metabolism, in the regulation of cell death and cell cycle. Earlier findings in our group demonstrated that phosphorylated glyoxalase I and its substrate methylglyoxal are important in TNF-induced cell death in the fibrosarcoma cell line L929. However, the ubiquitous nature of the glyoxalase system and the inevitable presence of methylglyoxal under physiological conditions, suggest a more fundamental role. The **scope** of this study was to investigate **the potential involvement of phosphorylated glyoxalase I in the GF-withdrawal induced G₁ arrest and apoptosis and in the regulation of the cell cycle.**

Research in mammalian cells, as well as in prokaryotic and plant cells indicate a strong correlation between the rate of growth on the one hand and the expression level and/or activity of glyoxalase I on the other. However, a mechanistic link has never been found. We therefore investigated whether the phosphorylation of glyoxalase I might be involved in the regulation of the cell cycle. Two cellular model systems were used to study the posttranslational modifications and activity of glyoxalase I in detail. First Ba/F3 cells were depleted of IL-3 to induce G₁ arrest and subsequently apoptosis. GF starvation has been shown to result in a significant decline in glucose metabolism. This would result in a change in intracellular levels of methylglyoxal, making this an attractive model system to study the glyoxalase system and possible modulation of the detoxification of MG. Concurrently attention was given to the **formation of specific methylglyoxal-derived advanced glycation endproducts (MG-AGEs)** and we attempted to situate these events in progression of the IL-3 depletion induced signaling. Time course experiments were also performed on IL-3 dependent Ba/F3 cells going through the cell cycle synchronously. Finally, in vitro studies were commenced to further explore the pathway upstream of the **phosphorylation of glyoxalase I.**

A possible mechanism used by MG-AGEs to exert their influence on the cell cycle and/or apoptosis is through activation of the Receptor for Advanced Glycation Endproducts (RAGE). This multiligand receptor has been reported to be involved in numerous cellular processes, including proliferation and survival. A dual role has been imputed on RAGE in both processes. RAGE has been involved in proliferation as well as in differentiation, and in survival as well as in cell death. The **aim** of the second part of this thesis was thus dedicated to the study of **RAGE during GF-depletion induced G₁ arrest apoptosis, and during the cell cycle.**

Reference List

1. **Dakin, H. D. and H. W. Dudley.** 1913. AN ENZYME CONCERNED WITH THE FORMATION OF HYDROXY ACIDS FROM KETONIC ALDEHYDES. *Journal of Biological Chemistry* **14**:155-157.
2. **Neuberg, C.** The destruction of lactic aldehyde and methylglyoxal by animal organs. *Biochemische Zeitschrift* **49**, 502-506. 1913.
3. **Embden, D. G., H. J. Deitricke, and G. Kraft.** The intermediate processes in glycolysis in muscle. *Klinische Wochenschrift* **12**, 213-215. 1933.
4. **Meyerhof, O.** The appearance and transformation of α -glycerophosphoric acid in the enzymic carbohydrate hydrolysis. *Biochemische Zeitschrift* **264**, 40-71. 1933.
5. **RACKER, E.** 1951. The mechanism of action of glyoxalase. *J.Biol.Chem.* **190**:685-696.
6. **Hopkins, F. G. and E. J. Morgan.** 1945. On the distribution of glyoxalase and glutathione. *Biochem.J.* **39**:320-324.
7. **Egyud, L. G. and A. Szent-Gyorgyi.** 1966. On the regulation of cell division. *Proc.Natl.Acad.Sci.U.S.A* **56**:203-207.
8. **Szent-Gyorgyi, A. and L. Egyud.** 1966. On Regulators of Cell Division. *Science* **152**:676-677.
9. **Szent-Gyorgyi, A., L. G. Egyud, and J. A. McLaughlin.** 1967. Keto-aldehydes and cell division. *Science* **155**: 539-541.
10. **Jowett, M. and J. H. Quastel.** 1933. The glyoxalase activity of the red blood cell: The function of glutathione. *Biochem.J.* **27**:486-498.
11. **Aronsson, A. C., E. Marmstal, and B. Mannervik.** 1978. Glyoxalase I, a zinc metalloenzyme of mammals and yeast. *Biochem.Biophys.Res.Comm.* **81**:1235-1240.
12. **Mannervik, B. Glyoxalase I.** Enzymatic basis of detoxification (Jacoby,W., ed.) **2**, 263-293. 1980. Academic Press, New York.
13. **Sellin, S., L. E. Eriksson, A. C. Aronsson, and B. Mannervik.** 1983. Octahedral metal coordination in the active site of glyoxalase I as evidenced by the properties of Co(II)-glyoxalase I. *J.Biol.Chem.* **258**:2091-2093.
14. **Thornalley, P. J.** 1990. The Glyoxalase System - New Developments Towards Functional-Characterization of A Metabolic Pathway Fundamental to Biological Life. *Biochemical Journal* **269**:1-11.
15. **Uotila, L.** Glutathione thiol esterases. *Glutathione: Chemical, Biochemical and Medical Aspects, Coenzymes and Cofactors III part A*, 767-804. 1989.
16. **Reeves, H. C. and S. J. Aji.** 1965. Enzymatic Synthesis and Metabolism of Hydroxypyruvic Aldehyde. *Journal of Biological Chemistry* **240**:569-573.
17. **Vander Jagt, D. L., E. Daub, J. A. Krohn, and L. P. Han.** 1975. Effects of pH and thiols on the kinetics of yeast glyoxalase I. An evaluation of the random pathway mechanism. *Biochemistry* **14**:3669-3675.
18. **Weaver, R. H. and H. A. Lardy.** 1961. Synthesis and Some Biochemical Properties of Phosphohydroxypyruvic Aldehyde and of 3-Phosphoglyceryl Glutathione Thiol Ester. *Journal of Biological Chemistry* **236**:313-317.
19. **Douglas, K. T.** 1987. Mechanism of action of glutathione-dependent enzymes. *Adv.Enzymol.Relat Areas Mol.Biol.* **59**:103-167.
20. **Marmstal, E., A. C. Aronsson, and B. Mannervik.** 1979. Comparison of Glyoxalase-I Purified from Yeast (*Saccharomyces-Cerevisiae*) with the Enzyme from Mammalian Sources. *Biochemical Journal* **183**:23-30.
21. **Marmstal, E. and B. Mannervik.** 1979. Purification, Characterization and Kinetic Studies of Glyoxalase-I from Rat-Liver. *Biochimica et Biophysica Acta* **566**:362-370.
22. **Shih, M. J., J. W. Edinger, and D. J. Creighton.** 1997. Diffusion-dependent kinetic properties of glyoxalase I and estimates of the steady-state concentrations of glyoxalase-pathway intermediates in glycolyzing erythrocytes. *European Journal of Biochemistry* **244**:852-857.
23. **Creighton, D. J., M. Migliorini, T. Pourmotabbed, and M. K. Guha.** 1988. Optimization of Efficiency in the Glyoxalase Pathway. *Biochemistry* **27**:7376-7384.
24. **Vander Jagt, D. L. and L. A. Hunsaker.** 2003. Methylglyoxal metabolism and diabetic complications: roles of aldose reductase, glyoxalase-1, betaine aldehyde dehydrogenase and 2-oxoaldehyde dehydrogenase. *Chemico-Biological Interactions* **143**:341-351.
25. **Cameron, A. D., B. Olin, M. Ridderstrom, B. Mannervik, and T. A. Jones.** 1997. Crystal structure of human glyoxalase .1. Evidence for gene duplication and 3D domain swapping. *Embo Journal* **16**:3386-3395.
26. **Cameron, A. D., M. Ridderstrom, B. Olin, M. J. Kavarana, D. J. Creighton, and B. Mannervik.** 1999. Reaction mechanism of glyoxalase I explored by an X-ray crystallographic analysis of the human enzyme in complex with a transition state analogue. *Biochemistry* **38**:13480-13490.
27. **Ridderstrom, M., A. D. Cameron, T. A. Jones, and B. Mannervik.** 1998. Involvement of an active-site Zn²⁺ ligand mechanism of human glyoxalase I. *Journal of Biological Chemistry* **273**:21623-21628.
28. **Garciaigniguez, L., L. Powers, B. Chance, S. Sellin, B. Mannervik, and A. S. Mildvan.** 1984. X-Ray Absorption Studies of the Zn²⁺ Site of Glyoxalase-I. *Biochemistry* **23**:685-689.
29. **Feierberg, I., V. Luzhkov, and J. Aqvist.** 2000. Computer simulation of primary kinetic isotope effects in the proposed rate-limiting step of the glyoxalase I catalyzed reaction. *Journal of Biological Chemistry* **275**:22657-22662.
30. **Himo, F. and P. E. M. Siegbahn.** 2001. Catalytic mechanism of glyoxalase I: A theoretical study. *Journal of the American Chemical Society* **123**:10280-10289.
31. **Richter, U. and M. Krauss.** 2001. Active site structure and mechanism of human glyoxalase I - An ab initio theoretical study. *Journal of the American Chemical Society* **123**:6973-6982.
32. **Akoachere, M., R. Iozef, S. Rahlfs, M. Deponte, B. Mannervik, D. J. Creighton, H. Schirmer, and K. Becker.**

2005. Characterization of the glyoxalases of the malarial parasite *Plasmodium falciparum* and comparison with their human counterparts. *Biological Chemistry* **386**:41-52.
33. **Antognelli, C., R. Romani, F. Baldracchini, A. De Santis, G. Andreani, and V. Talesa.** 2003. Different activity of glyoxalase system enzymes in specimens of *Sparus auratus* exposed to sublethal copper concentrations. *Chemo-Biological Interactions* **142**:297-305.
34. **Clugston, S. L., E. Daub, R. Kinach, D. Miedema, J. F. J. Barnard, and J. F. Honek.** 1997. Isolation and sequencing of a gene coding for glyoxalase I activity from *Salmonella typhimurium* and comparison with other glyoxalase I sequences. *Gene* **186**:103-111.
35. **Clugston, S. L., E. Daub, and J. F. Honek.** 1998. Identification of glyoxalase I sequences in *Brassica oleracea* and *Sporobolus stapfianus*: Evidence for gene duplication events. *Journal of Molecular Evolution* **47**:230-234.
36. **Clugston, S. L., J. F. J. Barnard, R. Kinach, D. Miedema, R. Ruman, E. Daub, and J. F. Honek.** 1998. Overproduction and characterization of a dimeric non-zinc glyoxalase I from *Escherichia coli*: Evidence for optimal activation by nickel ions. *Biochemistry* **37**:8754-8763.
37. **Sommer, A., P. Fischer, K. Krause, K. Boettcher, P. M. Brophy, R. D. Walter, and E. Liebau.** 2001. A stress-responsive glyoxalase I from the parasitic nematode *Onchocerca volvulus*. *Biochemical Journal* **353**:445-452.
38. **Sukdeo, N., S. L. Clugston, E. Daub, and J. F. Honek.** 2004. Distinct classes of glyoxalase I: metal specificity of the *Yersinia pestis*, *Pseudomonas aeruginosa* and *Neisseria meningitidis* enzymes. *Biochemical Journal* **384**:111-117.
39. **Veena and S. K. Sopory.** 1997. Molecular cloning of glyoxalase I from *Brassica juncea*. *Faseb Journal* **11**:A976.
40. **Veena, V. S. Reddy, and S. K. Sopory.** 1999. Glyoxalase I from *Brassica juncea*: Molecular cloning, regulation and its over-expression confer tolerance in transgenic tobacco under stress. *Plant Journal* **17**:385-395.
41. **Usui, Y., M. Nakase, H. Hotta, A. Urisu, N. Aoki, K. Kitajima, and T. Matsuda.** 2001. A 33-kDa allergen from rice (*Oryza sativa* L. Japonica) - cDNA cloning, expression, and identification as a novel glyoxalase I. *Journal of Biological Chemistry* **276**:11376-11381.
42. **Murata, K., Y. Inoue, H. Rhee, and A. Kimura.** 1989. 2-Oxoaldehyde metabolism in microorganisms. *Can.J.Microbiol.* **35**:423-431.
43. **Frickel, E. M., P. Jemth, M. Widersten, and B. Mannervik.** 2001. Yeast glyoxalase I is a monomeric enzyme with two active sites. *Journal of Biological Chemistry* **276**:1845-1849.
44. **Ranganathan, S., E. S. Walsh, A. K. Godwin, and K. D. Tew.** 1993. Cloning and Characterization of Human Colon Glyoxalase-I. *Journal of Biological Chemistry* **268**:5661-5667.
45. **Clapper, M. L., S. J. Hoffman, and K. D. Tew.** 1991. Glutathione S-Transferases in Normal and Malignant Human Colon Tissue. *Biochimica et Biophysica Acta* **1096**:209-216.
46. **Larsen, K., A. C. Aronsson, E. Marmstal, and B. Mannervik.** 1985. Immunological Comparison of Glyoxalase-I from Yeast and Mammals and Quantitative-Determination of the Enzyme in Human-Tissues by Radioimmunoassay. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* **82**:625-638.
47. **Bender, K. and K. H. Grzeschik.** 1976. Assignment of Genes for Human Glyoxalase-I to Chromosome-6 and for Human Esterase-D to Chromosome-13. *Cytogenetics and Cell Genetics* **16**:93-96.
48. **Kompf, J., S. Bissbort, and F. Schunter.** 1976. Confirmation of Linkage Between Loci for HI-A and Glyoxalase-1. *Human Genetics* **32**:197-198.
49. **Mayr, W. R., D. Mayr, J. Kompf, S. Bissbort, and H. Ritter.** 1976. Possible Linkage of HI-A and Glo. *Human Genetics* **31**:241-242.
50. **Ranganathan, S., P. J. Ciaccio, E. S. Walsh, and K. D. Tew.** 1999. Genomic sequence of human glyoxalase-I: analysis of promoter activity and its regulation. *Gene* **240**:149-155.
51. **Gale, C. P. and P. J. Grant.** 2004. The characterisation and functional analysis of the human glyoxalase-1 gene using methods of bioinformatics. *Gene* **340**:251-260.
52. **Kompf, J., S. Bissbort, S. Gussmann, and H. Ritter.** 1975. Polymorphism of Red-Cell Glyoxalase I (Ec-4.4.1.5) - New Genetic Marker in Man - Investigation of 169 Mother-Child Combinations. *Humangenetik* **27**:141-143.
53. **Kompf, J., S. Bissbort, and H. Ritter.** 1975. Red-Cell Glyoxalase I (Ec-4.4.1.5) - Formal Genetics and Linkage Relations. *Humangenetik* **28**:249-251.
54. **Kim, N. S., S. Sekine, N. Kiuchi, and S. Kato.** 1995. Cdna Cloning and Characterization of Human Glyoxalase-I Isoforms from Ht-1080 Cells. *Journal of Biochemistry* **117**:359-361.
55. **Junaid, M. A., D. Kowal, M. Barua, P. S. Pullarkat, S. S. Brooks, and R. K. Pullarkat.** 2004. Proteomic studies identified a single nucleotide polymorphism in glyoxalase I as autism susceptibility factor. *American Journal of Medical Genetics Part A* **131A**:11-17.
56. **Sidoti, A., C. Antognelli, C. Rinoldi, R. D'Angelo, V. Dattola, P. Girlanda, V. Talesa, and A. Amato.** 2007. Glyoxalase IA111E, paraoxonase 1 q192R and L55M polymorphisms: susceptibility factors of multiple sclerosis? *Multiple Sclerosis* **13**:446-453.
57. **Mccann, V. J., R. E. Davis, T. A. Welborn, I. J. Constable, and D. G. Beale.** 1981. Glyoxalase Phenotypes in Patients with Diabetes-Mellitus. *Australian and New Zealand Journal of Medicine* **11**:380-382.
58. **Mccann, V. J., R. E. Davis, and T. A. Welborn.** 1981. Distribution of Glyoxalase Phenotypes in Diabetic-Patients in Western Australia. *Diabetologia* **21**:76.
59. **Thornalley, P. J.** 1991. Population-Genetics of Human Glyoxalases. *Heredity* **67**:139-142.
60. **Samadi, A. A., S. A. Fullerton, D. G. Tortorellis, G. B. Johnson, S. D. Davidson, M. S. Choudhury, C. Mallouh, H. Tazaki, and S. Konno.** 2001. Glyoxalase I phenotype as a potential risk factor for prostate

- carcinoma. *Urology* **57**:183-187.
61. **Van Herreweghe, F., J. Q. Mao, F. W. R. Chaplen, J. Grooten, K. Gevaert, J. Vandekerckhove, and K. Vancompernelle.** 2002. Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methyl glyoxal-derived AGE. *Proceedings of the National Academy of Sciences of the United States of America* **99**:949-954.
 62. **Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura.** 1990. Sexual-Response of *Saccharomyces-Cerevisiae* - Phosphorylation of Yeast Glyoxalase-I by A Cell Extract of Mating Factor Treated-Cells. *Journal of Biochemistry* **108**:4-6.
 63. **Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura.** 1989. Sexual-Response in *Saccharomyces-Cerevisiae* - Alteration of Enzyme-Activity in the Glyoxalase System by Mating Factor. *Biochemical and Biophysical Research Communications* **165**:1091-1095.
 64. **Shin, R., S. Alvarez, A. Y. Burch, J. M. Jez, and D. P. Schachtman.** 2007. Phosphoproteomic identification of targets of the Arabidopsis sucrose nonfermenting-like kinase SnRK2.8 reveals a connection to metabolic processes. *Proceedings of the National Academy of Sciences of the United States of America* **104**:6460-6465.
 65. **Khan, M. M. K., A. Jan, H. Karibe, and S. Komatsu.** 2005. Identification of phosphoproteins regulated by gibberellin in rice leaf sheath. *Plant Molecular Biology* **58**:27-40.
 66. **Mitsumoto, A., K. R. Kim, G. Oshima, M. Kunimoto, K. Okawa, A. Iwamatsu, and Y. Nakagawa.** 1999. Glyoxalase I is a novel nitric-oxide-responsive protein. *Biochemical Journal* **344**:837-844.
 67. **Mitsumoto, A., K. R. Kim, G. Oshima, M. Kunimoto, K. Okawa, A. Iwamatsu, and Y. Nakagawa.** 2000. Nitric oxide inactivates glyoxalase I in cooperation with glutathione. *Journal of Biochemistry* **128**:647-654.
 68. **de, H., V. D. Rondas, J. Vandekerckhove, and K. Vancompernelle.** 2007. Tumour necrosis factor induces phosphorylation primarily of the nitric-oxide-responsive form of glyoxalase I. *Biochem.J.* **407**:121-128.
 69. **Sahoo, R., R. Sengupta, and S. Ghosh.** 2003. Nitrosative stress on yeast: inhibition of glyoxalase-I and glyceraldehyde-3-phosphate dehydrogenase in the presence of GSNO. *Biochemical and Biophysical Research Communications* **302**:665-670.
 70. **Guha, M. K., D. L. Vanderjagt, and D. J. Creighton.** 1988. Diffusion-Dependent Rates for the Hydrolysis Reaction Catalyzed by Glyoxalase-Ii from Rat Erythrocytes. *Biochemistry* **27**:8818-8822.
 71. **Hooper, N. I., M. J. Tisdale, and P. J. Thornalley.** 1987. Glyoxalase Activity During Differentiation of Human-Leukemia Cells-Invitro. *Leukemia Research* **11**:1141-1148.
 72. **Cameron, A. D., M. Ridderstrom, B. Olin, and B. Mannervik.** 1999. Crystal structure of human glyoxalase II and its complex with a glutathione thiolester substrate analogue. *Structure* **7**:1067-1078.
 73. **Ridderstrom, M., P. Jemth, A. D. Cameron, and B. Mannervik.** 2000. The active-site residue Tyr-175 in human glyoxalase II contributes to binding of glutathione derivatives. *Biochimica et Biophysica Acta-Protein Structure and Molecular Enzymology* **1481**:344-348.
 74. **Campos-Bermudez, V. A., N. R. Leite, R. Krog, A. J. Costa-Filho, F. C. Soncini, G. Oliva, and A. J. Vila.** 2007. Biochemical and structural characterization of *Salmonella typhimurium* glyoxalase II: new insights into metal ion selectivity. *Biochemistry* **46**:11069-11079.
 75. **Cho, M. Y., C. D. Bae, J. B. Park, and T. H. Lee.** 1998. Purification and cloning of glyoxalase II from rat liver. *Exp.Mol.Med.* **30**:53-57.
 76. **Maiti, M. K., S. Krishnasamy, H. A. Owen, and C. A. Makaroff.** 1997. Molecular characterization of glyoxalase II from *Arabidopsis thaliana*. *Plant Mol.Biol.* **35**:471-481.
 77. **Principato, G. B., G. Rosi, V. Talesa, E. Giovannini, and S. J. Norton.** 1987. A comparative study on glyoxalase II from vertebrata. *Enzyme* **37**:164-168.
 78. **Ridderstrom, M., F. Saccucci, U. Hellman, T. Bergman, G. Principato, and B. Mannervik.** 1996. Molecular cloning, heterologous expression, and characterization of human glyoxalase II. *J.Biol.Chem.* **271**:319-323.
 79. **Ridderstrom, M. and B. Mannervik.** 1997. Molecular cloning and characterization of the thiolesterase glyoxalase II from *Arabidopsis thaliana*. *Biochem.J.* **322 (Pt 2)**:449-454.
 80. **Talesa, V., G. B. Principato, S. J. Norton, S. Contenti, C. Mangiabene, and G. Rosi.** 1990. Isolation of glyoxalase II from bovine liver mitochondria. *Biochem.Int.* **20**:53-58.
 81. **Trincao, J., S. M. Sousa, L. Barata, C. Bonifacio, S. Carvalho, A. M. Tomas, A. E. Ferreira, C. Cordeiro, F. A. Ponces, and M. J. Romao.** 2006. Purification, crystallization and preliminary X-ray diffraction analysis of the glyoxalase II from *Leishmania infantum*. *Acta Crystallogr.Sect.F.Struct.Biol.Cryst.Comm.* **62**:805-807.
 82. **Yadav, S. K., S. L. Singla-Pareek, M. Kumar, A. Pareek, M. Saxena, N. B. Sarin, and S. K. Sopory.** 2007. Characterization and functional validation of glyoxalase II from rice. *Protein Expr.Purif.* **51**:126-132.
 83. **Honey, N. K. and T. B. Shows.** 1981. Assignment of the glyoxalase II gene (HAGH) to human chromosome 16. *Hum.Genet.* **58**:358-361.
 84. **Board, P. G.** 1980. Genetic polymorphism of human erythrocyte glyoxalase II. *Am.J.Hum.Genet.* **32**:690-694.
 85. **Cordell, P. A., T. S. Futers, P. J. Grant, and R. J. Pease.** 2004. The human hydroxyacylglutathione hydrolase (HAGH) gene encodes both cytosolic and mitochondrial forms of glyoxalase II. *Journal of Biological Chemistry* **279**:28653-28661.
 86. **Talesa, V., L. Uotila, M. Koivusalo, G. Principato, E. Giovannini, and G. Rosi.** 1989. Isolation of glyoxalase II from two different compartments of rat liver mitochondria. Kinetic and immunochemical characterization of the enzymes. *Biochim.Biophys.Acta* **993**:7-11.
 87. **Bito, A., M. Haider, I. Hadler, and M. Breitenbach.** 1997. Identification and phenotypic analysis of two glyoxalase II encoding genes from *Saccharomyces cerevisiae*, GLO2 and GLO4, and intracellular localization of the corresponding proteins. *J.Biol.Chem.* **272**:21509-21519.
 88. **Marasinghe, G. P. K., I. M. Sander, B. Bennett, G. Periyannan, K. W. Yang, C. A. Makaroff, and M. W.**

- Crowder.** 2005. Structural studies on a mitochondrial glyoxalase II. *Journal of Biological Chemistry* **280**:40668-40675.
89. **Araki, N., T. Morimasa, T. Sakai, H. Tokuoh, S. Yunoue, M. Kamo, K. Miyazaki, K. Abe, H. Saya, and A. Tsugita.** 2000. Comparative analysis of brain proteins from p53-deficient mice by two-dimensional electrophoresis. *Electrophoresis* **21**:1880-1889.
90. **Xu, Y. and X. B. Chen.** 2006. Glyoxalase II, a detoxifying enzyme of glycolysis byproduct methylglyoxal and a target of p63 and p73, is a pro-survival factor of the p53 family. *Journal of Biological Chemistry* **281**:26702-26713.
91. **Misra, K., A. B. Banerjee, S. Ray, and M. Ray.** 1995. Glyoxalase-iii from *Escherichia-Coli* - A Single Novel Enzyme for the Conversion of Methylglyoxal Into D-Lactate Without Reduced Glutathione. *Biochemical Journal* **305**:999-1003.
92. **Okado-Matsumoto, A. and I. Fridovich.** 2000. The role of alpha,beta-dicarbonyl compounds in the toxicity of short chain sugars. *Journal of Biological Chemistry* **275**:34853-34857.
93. **Benov, L., F. Sequeira, and A. F. Beema.** 2004. Role of rpoS in the regulation of glyoxalase III in *Escherichia coli*. *Acta Biochimica Polonica* **51**:857-860.
94. **Cooper, R. A.** 1975. Methylglyoxal synthase. *Methods Enzymol.* **41**:502-508.
95. **Hopper, D. J. and R. A. Cooper.** 1972. The purification and properties of *Escherichia coli* methylglyoxal synthase. *Biochem.J.* **128**:321-329.
96. **Murata, K., Y. Fukuda, K. Watanabe, T. Saikusa, M. Shimosaka, and A. Kimura.** 1985. Characterization of Methylglyoxal Synthase in *Saccharomyces-Cerevisiae*. *Biochemical and Biophysical Research Communications* **131**:190-198.
97. **Ray, S. and M. Ray.** 1981. Isolation of Methylglyoxal Synthase from Goat Liver. *Journal of Biological Chemistry* **256**:6230-6233.
98. **Deswal, R., T. N. Chakaravarty, and S. K. Sopory.** 1993. The Glyoxalase System in Higher-Plants - Regulation in Growth and Differentiation. *Biochemical Society Transactions* **21**:527-530.
99. **Sethi, U., A. Basu, and S. Guhamukherjee.** 1988. Control of Cell-Proliferation and Differentiation by Regulating Polyamine Biosynthesis in Cultures of Brassica and Its Correlation with Glyoxalase-I Activity. *Plant Science* **56**: 167-175.
100. **Ghosh, A. K.** 1977. Polymorphism of Red-Cell Glyoxalase-1 - with Special Reference to South and Southeast-Asia and Oceania. *Human Genetics* **39**:91-95.
101. **Paulus, C., B. Kollner, and H. J. Jacobsen.** 1993. Physiological and Biochemical-Characterization of Glyoxalase-I, A General Marker for Cell-Proliferation, from A Soybean Cell-Suspension. *Planta* **189**:561-566.
102. **Dudani, A. K., L. K. Srivastava, and R. Prasad.** 1984. Glyoxalase-I Activity and Cell-Cycle Regulation in Yeast. *Biochemical and Biophysical Research Communications* **119**:962-967.
103. **Bucking-Throm, E., W. Duntze, L. H. Hartwell, and T. R. Manney.** 1973. Reversible arrest of haploid yeast cells in the initiation of DNA synthesis by a diffusible sex factor. *Exp.Cell Res.* **76**:99-110.
104. **Duntze, W., D. Stotzler, E. Bucking-Throm, and S. Kalbitzer.** 1973. Purification and partial characterization of -factor, a mating-type specific inhibitor of cell reproduction from *Saccharomyces cerevisiae*. *Eur.J.Biochem.* **35**:357-365.
105. **Principato, G. B., M. Bodo, M. G. Biagioni, G. Rosi, and F. S. Liotti.** 1982. Glyoxalases and glutathione reductase activity changes in chicken liver during embryo development and after hatching. *Acta Embryol.Morphol.Exp.* **3**:173-179.
106. **Principato, G. B., P. Locci, G. Rosi, V. Tlesa, and E. Giovannini.** 1983. Activity Changes of Glyoxalase-I Glyoxalase-II and Glutathione-Reductase in Regenerating Rat-Liver. *Biochemistry International* **6**:249-255.
107. **McLellan, A. C. and P. J. Thornalley.** 1989. Glyoxalase Activity in Human Red Blood-Cells Fractioned by Age. *Mechanisms of Ageing and Development* **48**:63-71.
108. **Amicarelli, F., P. Sacchetta, S. Colafarina, S. Angelucci, M. Miranda, and C. Di Ilio.** 1998. Glyoxalases activity during *Bufo bufo* embryo development. *Mechanisms of Ageing and Development* **100**:261-267.
109. **Antognelli, C., F. Baldracchini, V. N. Tlesa, E. Costantini, A. Zucchi, and E. Mearini.** 2006. Overexpression of glyoxalase system enzymes in human kidney tumor. *Cancer Journal* **12**:222-228.
110. **Ayoub, F., M. Zaman, P. Thornalley, and J. Masters.** 1993. Glyoxalase activities in human tumour cell lines in vitro. *Anticancer Res.* **13**:151-155.
111. **Davidson, S. D., D. M. Milanese, C. Mallouh, M. S. Choudhury, H. Tazaki, and S. Konno.** 2002. A possible regulatory role of glyoxalase I in cell viability of human prostate cancer. *Urol.Res.* **30**:116-121.
112. **Di Ilio, C., S. Angelucci, A. Pennelli, A. Zezza, R. Tenaglia, and P. Sacchetta.** 1995. Glyoxalase activities in tumor and non-tumor human urogenital tissues. *Cancer Lett.* **96**:189-193.
113. **Jerzykowski, T., R. Winter, W. Matuszewski, and D. Piskorska.** 1978. A re-evaluation of studies on the distribution of glyoxalases in animal and tumour tissues. *Int.J.Biochem.* **9**:853-860.
114. **Mearini, E., R. Romani, L. Mearini, C. Antognelli, A. Zucchi, T. Baroni, M. Porena, and V. N. Tlesa.** 2002. Differing expression of enzymes of the glyoxalase system in superficial and invasive bladder carcinomas. *Eur.J.Cancer* **38**:1946-1950.
115. **Ranganathan, S. and K. D. Tew.** 1993. Analysis of glyoxalase-I from normal and tumor tissue from human colon. *Biochim.Biophys.Acta* **1182**:311-316.
116. **Rulli, A., L. Carli, R. Romani, T. Baroni, E. Giovannini, G. Rosi, and V. Tlesa.** 2001. Expression of glyoxalase I and II in normal and breast cancer tissues. *Breast Cancer Res.Treat.* **66**:67-72.
117. **Selicharova, I., K. Smutna, M. Sanda, K. Ubik, E. Matouskova, E. Bursikova, M. Brozova, J. Vydra, and J. Jiracek.** 2007. 2-DE analysis of a new human cell line EM-G3 derived from breast cancer progenitor cells and

- comparison with normal mammary epithelial cells. *Proteomics* **7**:1549-1559.
118. **Egyud, L. G. and A. Szent-Gyorgyi.** 1966. Cell division, SH, ketoaldehydes, and cancer. *Proc. Natl. Acad. Sci. U.S.A* **55**:388-393.
 119. **Szent-Gyorgyi, A., A. HEGYELI, and J. A. McLaughlin.** 1963. Growth and cellular constituents. *Proc. Natl. Acad. Sci. U.S.A* **49**:878-879.
 120. **Kalapos, M. P.** 1999. On the promine/retine theory of cell division: now and then. *Biochimica et Biophysica Acta-General Subjects* **1426**:1-16.
 121. **Espartero, J., I. Sanchez-Aguayo, and J. M. Pardo.** 1995. Molecular characterization of glyoxalase-I from a higher plant; upregulation by stress. *Plant Mol. Biol.* **29**:1223-1233.
 122. **Yadav, S. K., S. L. Singla-Pareek, M. Ray, M. K. Reddy, and S. K. Sopory.** 2005. Methylglyoxal levels in plants under salinity stress are dependent on glyoxalase I and glutathione. *Biochemical and Biophysical Research Communications* **337**:61-67.
 123. **Singla-Pareek, S. L., M. K. Reddy, and S. K. Sopory.** 2003. Genetic engineering of the glyoxalase pathway in tobacco leads to enhanced salinity tolerance. *Proc. Natl. Acad. Sci. U.S.A* **100**:14672-14677.
 124. **Singla-Pareek, S. L., S. K. Yadav, A. Pareek, M. K. Reddy, and S. K. Sopory.** 2006. Transgenic tobacco overexpressing glyoxalase pathway enzymes grow and set viable seeds in zinc-spiked soils. *Plant Physiology* **140**:613-623.
 125. **Singla-Pareek, S. L., S. K. Yadav, A. Pareek, M. K. Reddy, and S. K. Sopory.** 2007. Enhancing salt tolerance in a crop plant by overexpression of glyoxalase II. *Transgenic Res.*
 126. **Yadav, S. K., S. L. Singla-Pareek, M. K. Reddy, and S. K. Sopory.** 2005. Transgenic tobacco plants overexpressing glyoxalase enzymes resist an increase in methylglyoxal and maintain higher reduced glutathione levels under salinity stress. *FEBS Lett.* **579**:6265-6271.
 127. **Inoue, Y., Y. Tsujimoto, and A. Kimura.** 1998. Expression of the glyoxalase I gene of *Saccharomyces cerevisiae* is regulated by high osmolarity glycerol mitogen-activated protein kinase pathway in osmotic stress response. *Journal of Biological Chemistry* **273**:2977-2983.
 128. **Takatsume, Y., S. Izawa, and Y. Inoue.** 2005. Unique regulation of glyoxalase I activity during osmotic stress response in the fission yeast *Schizosaccharomyces pombe*: neither the mRNA nor the protein level of glyoxalase I increase under conditions that enhance its activity. *Archives of Microbiology* **183**:224-227.
 129. **Wang, Y., Y. Y. Cao, X. M. Jia, Y. B. Cao, P. H. Gao, X. P. Fu, K. Ying, W. S. Chen, and Y. Y. Jiang.** 2006. Cap1p is involved in multiple pathways of oxidative stress response in *Candida albicans*. *Free Radic. Biol. Med.* **40**:1201-1209.
 130. **Miller, A. G., D. G. Smith, M. Bhat, and R. H. Nagaraj.** 2006. Glyoxalase I is critical for human retinal capillary pericyte survival under hyperglycemic conditions. *Journal of Biological Chemistry* **281**:11864-11871.
 131. **Shinohara, M., P. J. Thornalley, I. Giardino, P. Beisswenger, S. R. Thorpe, J. Onorato, and M. Brownlee.** 1998. Overexpression of glyoxalase-I in bovine endothelial cells inhibits intracellular advanced glycation endproduct formation and prevents hyperglycemia-induced increases in macromolecular endocytosis. *Journal of Clinical Investigation* **101**:1142-1147.
 132. **Sakamoto, H., T. Mashima, A. Kizaki, S. Dan, Y. Hashimoto, M. Naito, and T. Tsuruo.** 2000. Glyoxalase I is involved in resistance of human leukemia cells to antitumor agent-induced apoptosis. *Blood* **95**:3214-3218.
 133. **Hamilton, D. S., M. J. Kavarana, E. M. Sharkey, J. L. Eiseman, and D. J. Creighton.** 1999. A new method for rapidly generating inhibitors of glyoxalase I inside tumor cells using S-(N-aryl-N-hydroxycarbamoyl)ethylsulfoxides. *J. Med. Chem.* **42**:1823-1827.
 134. **Sakamoto, H., T. Mashima, S. Sato, Y. Hashimoto, T. Yamori, and T. Tsuruo.** 2001. Selective activation of apoptosis program by S-p-bromobenzylglutathione cyclopentyl diester in glyoxalase I-overexpressing human lung cancer cells. *Clinical Cancer Research* **7**:2513-2518.
 135. **Sharkey, E. M., H. B. O'Neill, M. J. Kavarana, H. Wang, D. J. Creighton, D. L. Sentz, and J. L. Eiseman.** 2000. Pharmacokinetics and antitumor properties in tumor-bearing mice of an enediol analogue inhibitor of glyoxalase I. *Cancer Chemother. Pharmacol.* **46**:156-166.
 136. **Thornalley, P. J., M. J. Ladan, S. J. S. Ridgway, and Y. Kang.** 1996. Antitumor activity of S-(p-bromobenzyl)glutathione diesters in vitro: A structure-activity study. *Journal of Medicinal Chemistry* **39**:3409-3411.
 137. **Deswal, R. and S. K. Sopory.** 1999. Glyoxalase I from *Brassica juncea* is a calmodulin stimulated protein. *Biochimica et Biophysica Acta-Molecular Cell Research* **1450**:460-467.
 138. **White, J. S. and K. R. Rees.** 1982. Inhibitory Effects of Methyl Glyoxal on Dna, Rna and Protein-Synthesis in Cultured Guinea-Pig Keratocytes. *Chemico-Biological Interactions* **38**:339-347.
 139. **Rae, C., S. J. Bernersprice, B. T. Bulliman, and P. W. Kuchel.** 1990. Kinetic-Analysis of the Human Erythrocyte Glyoxalase System Using H-1-Nmr and A Computer-Model. *European Journal of Biochemistry* **193**:83-90.
 140. **Phillips, S. A. and P. J. Thornalley.** 1993. The Formation of Methylglyoxal from Triose Phosphates - Investigation Using A Specific Assay for Methylglyoxal. *European Journal of Biochemistry* **212**:101-105.
 141. **Richard, J. P.** 1993. Mechanism for the Formation of Methylglyoxal from Triosephosphates. *Biochemical Society Transactions* **21**:549-553.
 142. **Ahmed, N., S. Battah, N. Karachalias, R. Babaei-Jadidi, M. Horanyi, K. Baroti, S. Hollan, and P. J. Thornalley.** 2003. Increased formation of methylglyoxal and protein glycation, oxidation and nitrosation in triosephosphate isomerase deficiency. *Biochimica et Biophysica Acta-Molecular Basis of Disease* **1639**:121-132.
 143. **Pompliano, D. L., A. Peyman, and J. R. Knowles.** 1990. Stabilization of A Reaction Intermediate As A Catalytic Device - Definition of the Functional-Role of the Flexible Loop in Triosephosphate Isomerase. *Biochemistry* **29**:

- 3186-3194.
144. **Koop, D. R. and J. P. Casazza.** 1985. Identification of Ethanol-Inducible P-450 Isozyme 3AAs the Acetone and Acetol Monooxygenase of Rabbit Microsomes. *Journal of Biological Chemistry* **260**:3607-3612.
145. **Lyles, G. A. and J. Chalmers.** 1992. The Metabolism of Aminoacetone to Methylglyoxal by Semicarbazide-Sensitive Amine Oxidase in Human Umbilical Artery. *Biochemical Pharmacology* **43**:1409-1414.
146. **Reichard, G. A., C. L. Skutches, R. D. Hoeldtke, and O. E. Owen.** 1986. Acetone Metabolism in Humans During Diabetic-Ketoacidosis. *Diabetes* **35**:668-674.
147. **Gracy, R. W., P. K. Tsai, and P. M. Yuan.** 1975. Isolation and Characterization of A Methylglyoxal Synthetase from *Proteus Vulgaris*. *Federation Proceedings* **34**:575.
148. **Hopper, D. J. and R. A. Cooper.** 1971. The regulation of *Escherichia coli* methylglyoxal synthase; a new control site in glycolysis? *FEBS Lett.* **13**:213-216.
149. **Rose, I. A. and J. S. Nowick.** 2002. Methylglyoxal synthetase, enol-pyruvate dehydrogenase, glutathione and the glyoxalase system. *Journal of the American Chemical Society* **124**:13047-13052.
150. **Beisswenger, P. J., S. K. Howell, K. Smith, and B. S. Szwegold.** 2003. Glyceraldehyde-3-phosphate dehydrogenase activity as an independent modifier of methylglyoxal levels in diabetes. *Biochimica et Biophysica Acta-Molecular Basis of Disease* **1637**:98-106.
151. **Knight, R. J., K. F. Kofoed, H. R. Schelbert, and D. B. Buxton.** 1996. Inhibition of glyceraldehyde-3-phosphate dehydrogenase in post-ischaemic myocardium. *Cardiovascular Research* **32**:1016-1023.
152. **Lee, H. J., S. K. Howell, R. J. Sanford, and P. J. Beisswenger.** 2005. Methylglyoxal can modify GAPDH activity and structure. *Maillard Reaction: Chemistry at the Interface of Nutrition, Aging, and Disease* **1043**:135-145.
153. **Chaplen, F. W. R., W. E. Fahl, and D. C. Cameron.** 1998. Evidence of high levels of methylglyoxal in cultured Chinese hamster ovary cells. *Proceedings of the National Academy of Sciences of the United States of America* **95**:5533-5538.
154. **Lo, T. W. C., M. E. Westwood, A. C. Mclellan, T. Selwood, and P. J. Thornalley.** 1994. Binding and Modification of Proteins by Methylglyoxal Under Physiological Conditions - A Kinetic and Mechanistic Study with N-Alpha-Acetylarginine, N-Alpha-Acetylcysteine, and N-Alpha-Acetyllysine, and Bovine Serum-Albumin. *Journal of Biological Chemistry* **269**:32299-32305.
155. **Oya, T., N. Hattori, Y. Mizuno, S. Miyata, S. Maeda, T. Osawa, and K. Uchida.** 1999. Methylglyoxal modification of protein - Chemical and immunochemical characterization of methylglyoxal-arginine adducts. *Journal of Biological Chemistry* **274**:18492-18502.
156. **Weber, A. L.** 1984. Prebiotic Formation of Energy-Rich Thioesters from Glyceraldehyde and N-Acetylcysteine. *Origins of Life and Evolution of the Biosphere* **15**:17-27.
157. **Westwood, M. E. and P. J. Thornalley.** 1995. Molecular Characteristics of Methylglyoxal-Modified Bovine and Human Serum Albumins - Comparison with Glucose-Derived Advanced Glycation Endproduct-Modified Serum Albumins. *Journal of Protein Chemistry* **14**:359-372.
158. **Thornalley, P. J.** 1996. Pharmacology of methylglyoxal: Formation, modification of proteins and nucleic acids, and enzymatic detoxification - A role in pathogenesis and antiproliferative chemotherapy. *General Pharmacology* **27**:565-573.
159. **Thornalley, P. J.** 2005. Dicarbonyl intermediates in the Maillard reaction. *Maillard Reaction: Chemistry at the Interface of Nutrition, Aging, and Disease* **1043**:111-117.
160. **Brownlee, M.** 2001. Biochemistry and molecular cell biology of diabetic complications. *Nature* **414**:813-820.
161. **Singh, R., A. Barden, T. Mori, and L. Beilin.** 2001. Advanced glycation end-products: a review. *Diabetologia* **44**:129-146.
162. **Singh, R., A. Barden, T. Mori, and L. Beilin.** 2002. Advanced glycation end-products: a review (vol 44, pg 129, 2001). *Diabetologia* **45**:293.
163. **Vlassara, H. and M. R. Palace.** 2002. Diabetes and advanced glycation endproducts. *Journal of Internal Medicine* **251**:87-101.
164. **Wautier, J. L. and A. M. Schmidt.** 2004. Protein glycation - A firm link to endothelial cell dysfunction. *Circulation Research* **95**:233-238.
165. **Jana, C. K., N. Das, and R. S. Sohal.** 2002. Specificity of age-related carbonylation of plasma proteins in the mouse and rat. *Archives of Biochemistry and Biophysics* **397**:433-439.
166. **Poggioli, S., H. Bakala, and B. Friguet.** 2002. Age-related increase of protein glycation in peripheral blood lymphocytes is restricted to preferential target proteins. *Experimental Gerontology* **37**:1207-1215.
167. **Poggioli, S., J. Mary, H. Bakala, and B. Friguet.** 2004. Evidence of preferential protein targets for age-related modifications in peripheral blood lymphocytes. *Strategies for Engineered Negligible Senescence: Why Genuine Control of Aging May be Foreseeable* **1019**:211-214.
168. **Gao, Y. and Y. S. Wang.** 2006. Site-selective modifications of arginine residues in human hemoglobin induced by methylglyoxal. *Biochemistry* **45**:15654-15660.
169. **Sakamoto, H., T. Mashima, K. Yamamoto, and T. Tsuruo.** 2002. Modulation of heat-shock protein 27 (Hsp27) anti-apoptotic activity by methylglyoxal modification. *Journal of Biological Chemistry* **277**:45770-45775.
170. **Takatsume, Y., S. Izawa, and Y. Inoue.** 2006. Methylglyoxal as a signal initiator for activation of the stress-activated protein kinase cascade in the fission yeast *Schizosaccharomyces pombe*. *Journal of Biological Chemistry* **281**:9086-9092.
171. **Vander Jagt, D. L., R. K. Hassebrook, L. A. Hunsaker, W. T. Brown, and R. E. Royer.** 2001. Metabolism of the 2-oxoaldehyde methylglyoxal by aldose reductase and by glyoxalase-I: roles for glutathione in both enzymes and implications for diabetic complications. *Chemico-Biological Interactions* **130**:549-562.

172. Jagt, D. L. V., N. S. Kolb, T. J. V. Jagt, J. Chino, F. J. Martinez, L. A. Hunsaker, and R. E. Royer. 1995. Substrate-Specificity of Human Aldose Reductase - Identification of 4-Hydroxynonenal As An Endogenous Substrate. *Biochimica et Biophysica Acta-Protein Structure and Molecular Enzymology* **1249**:117-126.
173. Vanderjagt, D. L., B. Robinson, K. K. Taylor, and L. A. Hunsaker. 1992. Reduction of Trioses by NADPH-Dependent Aldo-Keto Reductases - Aldose Reductase, Methylglyoxal, and Diabetic Complications. *Journal of Biological Chemistry* **267**:4364-4369.
174. Izaguirre, G., A. Kikonyogo, and R. Pietruszko. 1998. Methylglyoxal as substrate and inhibitor of human aldehyde dehydrogenase: Comparison of kinetic properties among the three isozymes. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* **119**:747-754.
175. Fujii, E., H. Iwase, I. Ishiikarakasa, Y. Yajima, and K. Hotta. 1995. The Presence of 2-Keto-3-Deoxygluconic Acid and Oxaldehyde Dehydrogenase-Activity in Human Erythrocytes. *Biochemical and Biophysical Research Communications* **210**:852-857.
176. Monder, C. 1965. Oxidation of methylglyoxal to pyruvic acid by a sheep-liver enzyme. *Biochim.Biophys.Acta* **99**: 573-575.
177. Monder, C. 1967. Alpha-keto aldehyde dehydrogenase, an enzyme that catalyzes the enzymic oxidation of methylglyoxal to pyruvate. *J.Biol.Chem.* **242**:4603-4609.
178. Oimomi, M., F. Hata, N. Igaki, T. Nakamichi, S. Baba, and H. Kato. 1989. Purification of Alpha-Ketoaldehyde Dehydrogenase from the Human-Liver and Its Possible Significance in the Control of Glycation. *Experientia* **45**: 463-466.
179. Takatsume, Y., S. Izawa, and Y. Inoue. 2004. Identification of thermostable glyoxalase I in the fission yeast *Schizosaccharomyces pombe*. *Archives of Microbiology* **181**:371-377.
180. Kuhla, B., K. Boeck, H. J. Luth, A. Schmidt, B. Weigle, M. Schnitz, V. Ogunlade, G. Munch, and T. Arendt. 2006. Age-dependent changes of glyoxalase I expression in human brain. *Neurobiology of Aging* **27**:815-822.
181. Kuhla, B., K. Boeck, A. Schmidt, V. Ogunlade, T. Arendt, G. Munch, and H. J. Luth. 2007. Age- and stage-dependent glyoxalase I expression and its activity in normal and Alzheimer's disease brains. *Neurobiology of Aging* **28**:29-41.
182. Ayoub, F. M., R. E. Allen, and P. J. Thornalley. 1993. Inhibition of Proliferation of Human Leukemia 60-Cells by Methylglyoxal *In vitro*. *Leukemia Research* **17**:397-401.
183. Clelland, J. D., R. E. Allen, and P. J. Thornalley. 1992. Inhibition of Growth of Human Leukemia-60 Cells by S-2-Hydroxyacylglutathiones and Monoethyl Ester Derivatives. *Biochemical Pharmacology* **44**:1953-1959.
184. Edwards, L. G., A. Adesida, and P. J. Thornalley. 1996. Inhibition of human leukaemia 60 cell growth by s-D-lactoylglutathione *in vitro*. Mediation by metabolism to n-D-lactoylcysteine and induction of apoptosis. *Leukemia Research* **20**:17-26.
185. Kang, Y. B., L. G. Edwards, and P. J. Thornalley. 1996. Effect of methylglyoxal on human leukaemia 60 cell growth: Modification of DNA, G(1) growth arrest and induction of apoptosis. *Leukemia Research* **20**:397-405.
186. Thornalley, P. J. and M. J. Tisdale. 1988. Inhibition of Proliferation of Human Promyelocytic Leukemia HL60 Cells by S-D-Lactoylglutathione *In vitro*. *Leukemia Research* **12**:897-904.
187. Milanesa, D. M., M. S. Choudhury, C. Mallouh, H. Tazaki, and S. Konno. 2000. Methylglyoxal-induced apoptosis in human prostate carcinoma: Potential modality for prostate cancer treatment. *European Urology* **37**: 728-734.
188. Amicarelli, F., T. Bucciarelli, A. Poma, P. Aimola, C. Di Ilio, A. M. Ragnelli, and M. Miranda. 1998. Adaptive response of human melanoma cells to methylglyoxal injury. *Carcinogenesis* **19**:519-523.
189. Maeta, K., K. Mori, Y. Takatsume, S. Izawa, and Y. Inoue. 2005. Diagnosis of cell death induced by methylglyoxal, a metabolite derived from glycolysis, in *Saccharomyces cerevisiae*. *Fems Microbiology Letters* **243**:87-92.
190. Di Loreto, S., V. Caracciolo, S. Colafarina, P. Sebastiani, A. Gasbarri, and F. Amicarelli. 2004. Methylglyoxal induces oxidative stress-dependent cell injury and up-regulation of interleukin-10 and nerve growth factor in cultured hippocampal neuronal cells. *Brain Research* **1006**:157-167.
191. Du, X. L., D. Edelstein, L. Rossetti, I. G. Fantus, H. Goldberg, F. Ziyadeh, J. Wu, and M. Brownlee. 2000. Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proceedings of the National Academy of Sciences of the United States of America* **97**:12222-12226.
192. Okado, A., Y. Kawasaki, Y. Hasuike, M. Takahashi, T. Teshima, J. Fujii, and N. Taniguchi. 1996. Induction of apoptotic cell death by methylglyoxal and 3-deoxyglucosone in macrophage-derived cell lines. *Biochemical and Biophysical Research Communications* **225**:219-224.
193. Yim, M. B., S. O. Kang, and P. B. Chock. 2000. Enzyme-like activity of glycated cross-linked proteins in free radical generation. *Reactive Oxygen Species: from Radiation to Molecular Biology* **899**:168-181.
194. Amicarelli, F., S. Colafarina, F. Cattani, A. Cimini, C. Di Ilio, M. P. Ceru, and M. Miranda. 2003. Scavenging system efficiency is crucial for cell resistance to ROS-mediated methylglyoxal injury. *Free Radical Biology and Medicine* **35**:856-871.
195. Du, J., H. Suzuki, F. Nagase, A. A. Akhand, T. Yokoyama, T. Miyata, K. Kurokawa, and I. Nakashima. 2000. Methylglyoxal induces apoptosis in Jurkat leukemia T cells by activating c-jun N-terminal kinase. *Journal of Cellular Biochemistry* **77**:333-344.
196. Beard, K. M., N. Shangari, B. Wu, and P. J. O'Brien. 2003. Metabolism, not autoxidation, plays a role in alpha-oxoaldehyde- and reducing sugar-induced erythrocyte GSH depletion: Relevance for diabetes mellitus. *Molecular and Cellular Biochemistry* **252**:331-338.
197. Choudhary, D., D. Chandra, and R. K. Kale. 1997. Influence of methylglyoxal on antioxidant enzymes and

- oxidative damage. *Toxicology Letters* **93**:141-152.
198. **Di Loreto, S., V. Zimmitti, P. Sebastiani, C. Cervelli, S. Falone, and F. Amicarelli.** 2007. Methylglyoxal causes strong weakening of detoxifying capacity and apoptotic cell death in rat hippocampal neurons. *Int.J.Biochem.Cell Biol.*
199. **Thornalley, P. J.** 1998. Glutathione-dependent detoxification of alpha-oxoaldehydes by the glyoxalase system: involvement in disease mechanisms and antiproliferative activity of glyoxalase I inhibitors. *Chemico-Biological Interactions* **112**:137-151.
200. **Akhand, A. A., K. Hossain, H. Mitsui, M. Kato, T. Miyata, R. Inagi, J. Du, K. Takeda, Y. Kawamoto, H. Suzuki, K. Kurokawa, and I. Nakashima.** 2001. Glyoxal and methylglyoxal trigger distinct signals for map family kinases and caspase activation in human endothelial cells. *Free Radical Biology and Medicine* **31**:20-30.
201. **Du, J., H. Suzuki, F. Nagase, A. A. Akhand, X. Y. Ma, T. Yokoyama, T. Miyata, and I. M. Nakashima.** 2001. Superoxide-mediated early oxidation and activation of ASK1 are important for initiating methylglyoxal-induced apoptosis process. *Free Radical Biology and Medicine* **31**:469-478.
202. **Liu, B. F., S. Miyata, Y. Hirota, S. Higo, H. Miyazaki, M. Fukunaga, Y. Hamada, S. Ueyama, O. Muramoto, A. Uriuhara, and M. Kasuga.** 2003. Methylglyoxal induces apoptosis through activation of p38 mitogen-activated protein kinase in rat mesangial cells. *Kidney International* **63**:947-957.
203. **Du, J., S. Cai, H. Suzuki, A. A. Akhand, X. Ma, Y. Takagi, T. Miyata, I. Nakashima, and F. Nagase.** 2003. Involvement of MEKK1/ERK/P21Waf1/Cip1 signal transduction pathway in inhibition of IGF-I-mediated cell growth response by methylglyoxal. *J.Cell Biochem.* **88**:1235-1246.
204. **Jia, X. M., D. J. H. Olson, A. R. S. Ross, and L. Y. Wu.** 2006. Structural and functional changes in human insulin induced by methylglyoxal. *Faseb Journal* **20**:1555-+.
205. **Riboulet-Chavey, A., A. Pierron, I. Durand, J. Murdaca, J. Giudicelli, and E. Van Obberghen.** 2006. Methylglyoxal impairs the insulin signaling pathways independently of the. formation of intracellular reactive oxygen species. *Diabetes* **55**:1289-1299.
206. **Cantero, A. V., M. Portero-Otin, V. Ayala, N. Auge, M. Sanson, M. Elbaz, J. C. Thiers, R. Pamplona, R. Salvayre, and A. Negre-Salvayre.** 2007. Methylglyoxal induces advanced glycation end product (AGEs) formation and dysfunction of PDGF receptor-beta: implications for diabetic atherosclerosis. *Faseb Journal* **21**: 3096-3106.
207. **Portero-Otin, M., R. Pamplona, M. J. Bellmunt, C. Ruiz, A. Lafarga, R. Salvayre, and A. Negre-Salvayre.** 2000. Maillard-reaction derived oxoaldehydes inhibit epidermal-growth factor receptor activity in human cultured endothelial cells. *Diabetes* **49**:A333-A334.
208. **Portero-Otin, M., R. Pamplona, M. J. Bellmunt, M. C. Ruiz, J. Prat, R. Salvayre, and A. Negre-Salvayre.** 2002. Advanced glycation end product precursors impair epidermal growth factor receptor signaling. *Diabetes* **51**:1535-1542.
209. **Biswas, S., M. Ray, S. Misra, D. P. Dutta, and S. Ray.** 1997. Selective inhibition of mitochondrial respiration and glycolysis in human leukaemic leucocytes by methylglyoxal. *Biochemical Journal* **323**:343-348.
210. **de Arriba, S. G., G. Stuchbury, J. Yarin, J. Burnell, C. Loske, and G. Munch.** 2007. Methylglyoxal impairs glucose metabolism and leads to energy depletion in neuronal cells - protection by carbonyl scavengers. *Neurobiology of Aging* **28**:1044-1050.
211. **Halder, J., M. Ray, and S. Ray.** 1993. Inhibition of Glycolysis and Mitochondrial Respiration of Ehrlich Ascites-Carcinoma Cells by Methylglyoxal. *International Journal of Cancer* **54**:443-449.
212. **Ray, M., J. Halder, S. K. Dutta, and S. Ray.** 1991. Inhibition of Respiration of Tumor-Cells by Methylglyoxal and Protection of Inhibition by Lactaldehyde. *International Journal of Cancer* **47**:603-609.
213. **Ray, S., S. Dutta, J. Halder, and M. Ray.** 1994. Inhibition of Electron Flow-Through Complex I of the Mitochondrial Respiratory-Chain of Ehrlich Ascites-Carcinoma Cells by Methylglyoxal. *Biochemical Journal* **303**:69-72.
214. **Ray, S., S. Biswas, and M. Ray.** 1997. Similar nature of inhibition of mitochondrial respiration of heart tissue and malignant cells by methylglyoxal. A vital clue to understand the biochemical basis of malignancy. *Molecular and Cellular Biochemistry* **171**:95-103.
215. **Ray, M., N. Basu, and S. Ray.** 1997. Inactivation of glyceraldehyde-3-phosphate dehydrogenase of human malignant cells by methylglyoxal. *Molecular and Cellular Biochemistry* **177**:21-26.
216. **Oya-Ito, T., B. F. Liu, and R. H. Nagaraj.** 2006. Effect of methylglyoxal modification and phosphorylation on the chaperone and anti-apoptotic properties of heat shock protein 27. *Journal of Cellular Biochemistry* **99**:279-291.
217. **Schalkwijk, C. G., J. van Bezu, R. C. van der Schors, K. Uchida, C. D. A. Stehouwer, and V. W. M. van Hinsbergh.** 2006. Heat-shock protein 27 is a major methylglyoxal-modified protein in endothelial cells (vol 580, pg 1565, 2006). *Febs Letters* **580**:2400.
218. **Schalkwijk, C. G., J. van Bezu, R. C. van der Schors, K. Uchida, C. D. A. Stehouwer, and V. W. M. van Hinsbergh.** 2006. Heat-shock protein 27 is a major methylglyoxal-modified protein in endothelial cells. *Febs Letters* **580**:1565-1570.
219. **van Heijst, J. W. J., H. W. M. Niessen, R. J. Musters, V. W. M. van Hinsbergh, K. Hoekman, and C. G. Schalkwijk.** 2006. Argpyrimidine-modified Heat Shock Protein 27 in human non-small cell lung cancer: A possible mechanism for evasion of apoptosis. *Cancer Letters* **241**:309-319.
220. **Johans, M., E. Milanese, M. Franck, C. Johans, J. Liobikas, M. Panagiotaki, L. Greci, G. Principato, P. K. J. Kinnunen, P. Bernardi, P. Costantini, and O. Eriksson.** 2005. Modification of permeability transition pore arginine(s) by phenylglyoxal derivatives in isolated mitochondria and mammalian cells - Structure-function

- relationship of arginine ligands. *Journal of Biological Chemistry* **280**:12130-12136.
221. **Speer, O., S. Morkunaite-Haimi, J. Liobikas, M. Franck, L. Hensbo, M. D. Linder, P. K. J. Kinnunen, T. Wallimann, and O. Eriksson.** 2003. Rapid suppression of mitochondrial permeability transition by methylglyoxal - Role of reversible arginine modification. *Journal of Biological Chemistry* **278**:34757-34763.
 222. **Maeta, K., S. Izawa, S. Okazaki, S. Kuge, and Y. Inoue.** 2004. Activity of the Yap1 transcription factor in *Saccharomyces cerevisiae* is modulated by methylglyoxal, a metabolite derived from glycolysis. *Molecular and Cellular Biology* **24**:8753-8764.
 223. **Aguilera, J., S. Rodriguez-Vargas, and J. A. Prieto.** 2005. The HOG MAP kinase pathway is required for the induction of methylglyoxal-responsive genes and determines methylglyoxal resistance in *Saccharomyces cerevisiae*. *Molecular Microbiology* **56**:228-239.
 224. **Maeta, K., S. Izawa, and Y. Inoue.** 2005. Methylglyoxal, a metabolite derived from glycolysis, functions as a signal initiator of the high osmolarity glycerol-mitogen-activated protein kinase cascade and Calcineurin/Crz1-mediated pathway in *Saccharomyces cerevisiae*. *Journal of Biological Chemistry* **280**:253-260.
 225. **Aguilera, J. and J. A. Prieto.** 2001. The *Saccharomyces cerevisiae* aldose, reductase is implied in the metabolism of methylglyoxal in response to stress conditions. *Current Genetics* **39**:273-283.
 226. **Aguilera, J. and J. A. Prieto.** 2004. Yeast cells display a regulatory mechanism in response to methylglyoxal. *Fems Yeast Research* **4**:633-641.
 227. **Carlson, C. J., S. Koterski, R. J. Sciotti, G. B. Poccard, and C. M. Rondinone.** 2003. Enhanced basal activation of mitogen-activated protein kinases in adipocytes from type 2 diabetes - Potential role of p38 in the downregulation of GLUT4 expression. *Diabetes* **52**:634-641.
 228. **Akhand, A. A., K. Hossain, M. Kato, T. Miyata, J. Du, H. Suzuki, K. Kurokawa, and I. Nakashima.** 2001. Glyoxal and methylglyoxal induce aggregation and inactivation of ERK in human endothelial cells. *Free Radical Biology and Medicine* **31**:1228-1235.
 229. **Yao, D., T. Taguchi, T. Matsumura, R. Pestell, D. Edelstein, I. Giardino, G. Suske, N. Rabbani, P. J. Thornalley, V. P. Sarthy, H. P. Hammes, and M. Brownlee.** 2007. High glucose increases angiopoietin-2 transcription in microvascular endothelial cells through methylglyoxal modification of mSin3A. *Journal of Biological Chemistry* **282**:31038-31045.
 230. **Yao, D. C., T. Taguchi, T. Matsumura, R. Pestell, D. Edelstein, I. Giardino, G. Suske, N. Ahmed, P. J. Thornalley, V. P. Sarthy, H. P. Hammes, and M. Brownlee.** 2006. Methylglyoxal modification of mSin3A links glycolysis to angiopoietin-2 transcription. *Cell* **124**:275-286.
 231. **Horiuchi, S., T. Higashi, K. Ikeda, T. Saishoji, Y. Jinnouchi, H. Sano, R. Shibayama, T. Sakamoto, and N. Araki.** 1996. Advanced glycation end products and their recognition by macrophage and macrophage-derived cells. *Diabetes* **45**:S73-S76.
 232. **Vlassara, H., M. Brownlee, and A. Cerami.** 1986. Novel Macrophage Receptor for Glucose-Modified Proteins Is Distinct from Previously Described Scavenger Receptors. *Journal of Experimental Medicine* **164**:1301-1309.
 233. **Yang, Z., Z. Makita, Y. Horii, S. Brunelle, A. Cerami, P. Sehajpal, M. Suthanthiran, and H. Vlassara.** 1991. 2 Novel Rat-Liver Membrane-Proteins That Bind Advanced Glycosylation Endproducts - Relationship to Macrophage Receptor for Glucose-Modified Proteins. *Journal of Experimental Medicine* **174**:515-524.
 234. **Araki, N., T. Higashi, T. Mori, R. Shibayama, Y. Kawabe, T. Kodama, K. Takahashi, M. Shichiri, and S. Horiuchi.** 1995. Macrophage Scavenger Receptor Mediates the Endocytic Uptake and Degradation of Advanced Glycation End-Products of the Maillard Reaction. *European Journal of Biochemistry* **230**:408-415.
 235. **Li, Y. M., T. Mitsuhashi, D. Wojciechowicz, R. Gilmore, and H. Vlassara.** 1995. Elucidation of Advanced Glycation-Receptor Components P60/Ost-48 and P90/80-K-H. *Diabetes* **44**:A113.
 236. **Li, Y. M., T. Mitsuhashi, D. Wojciechowicz, N. Shimizu, J. Li, A. Stitt, C. J. He, D. Banerjee, and H. Vlassara.** 1996. Molecular identity and cellular distribution of advanced glycation endproduct receptors: Relationship of p60 to OST-48 and p90 to 80K-H membrane proteins. *Proceedings of the National Academy of Sciences of the United States of America* **93**:11047-11052.
 237. **Vlassara, H., Y. M. Li, F. Imani, D. Wojciechowicz, Z. Yang, F. T. Liu, and A. Cerami.** 1995. Identification of Galectin-3 As A High-Affinity Binding-Protein for Advanced Glycation End-Products (Age) - A New Member of the Age-Receptor Complex. *Molecular Medicine* **1**:634-646.
 238. **Li, Y. M., A. X. Tan, and H. Vlassara.** 1995. Antibacterial Activity of Lysozyme and Lactoferrin Is Inhibited by Binding of Advanced Glycation-Modified Proteins to A Conserved Motif. *Nature Medicine* **1**:1057-1061.
 239. **Neeper, M., A. M. Schmidt, J. Brett, S. D. Yan, F. Wang, Y. C. E. Pan, K. Elliston, D. Stern, and A. Shaw.** 1992. Cloning and Expression of A Cell-Surface Receptor for Advanced Glycosylation End-Products of Proteins. *Journal of Biological Chemistry* **267**:14998-15004.
 240. **Schmidt, A. M., M. Vianna, M. Gerlach, J. Brett, J. Ryan, J. Kao, C. Esposito, H. Hegarty, W. Hurley, M. Clauss, F. Wang, Y. C. E. Pan, T. C. Tsang, and D. Stern.** 1992. Isolation and Characterization of 2 Binding-Proteins for Advanced Glycosylation End-Products from Bovine Lung Which Are Present on the Endothelial-Cell Surface. *Journal of Biological Chemistry* **267**:14987-14997.
 241. **Huttunen, H. J., C. Fages, and H. Rauvala.** 1999. Receptor for advanced glycation end products (RAGE)-mediated neurite outgrowth and activation of NF-kappa B require the cytoplasmic domain of the receptor but different downstream signaling pathways. *Journal of Biological Chemistry* **274**:19919-19924.
 242. **Kislinger, T., C. F. Fu, B. Huber, W. Qu, A. Taguchi, S. D. Yan, M. Hofmann, S. F. Yan, M. Pischetsrieder, D. Stern, and A. M. Schmidt.** 1999. N-epsilon-(carboxymethyl)lysine adducts of proteins are ligands for receptor for advanced glycation end products that activate cell signaling pathways and modulate gene expression. *Journal of Biological Chemistry* **274**:31740-31749.
 243. **Leclerc, E., G. Fritz, M. Weibel, C. W. Heizmann, and A. Galichet.** 2007. S100B and S100A6 differentially

- modulate cell survival by interacting with distinct RAGE (receptor for advanced glycation end products) immunoglobulin domains. *Journal of Biological Chemistry* **282**:31317-31331.
244. **Dattilo, B. M., G. Fritz, E. Leclerc, C. W. Vander Kooi, C. W. Heizmann, and W. J. Chazin.** 2007. The extracellular region of the receptor for advanced glycation end products is composed of two independent structural units. *Biochemistry* **46**:6957-6970.
245. **Yonekura, H., Y. Yamamoto, S. Sakurai, R. G. Petrova, M. J. Abedin, H. Li, K. Yasui, M. Takeuchi, Z. Makita, S. Takasawa, H. Okamoto, T. Watanabe, and H. Yamamoto.** 2003. Novel splice variants of the receptor for advanced glycation end-products expressed in human vascular endothelial cells and pericytes, and their putative roles in diabetes-induced vascular injury. *Biochemical Journal* **370**:1097-1109.
246. **Hanford, L. E., J. J. Enghild, Z. Valnickova, S. V. Petersen, L. M. Schaefer, T. M. Schaefer, T. A. Reinhart, and T. D. Oury.** 2004. Purification and characterization of mouse soluble receptor for advanced glycation end products (sRAGE). *Journal of Biological Chemistry* **279**:50019-50024.
247. **Ding, Q. X. and J. N. Keller.** 2005. Evaluation of rage isoforms, ligands, and signaling in the brain. *Biochimica et Biophysica Acta-Molecular Cell Research* **1746**:18-27.
248. **Brett, J., A. M. Schmidt, S. D. Yan, Y. S. Zou, E. Weidman, D. Pinsky, R. Nowygrod, M. Neeper, C. Przysiecki, A. Shaw, A. Migheli, and D. Stern.** 1993. Survey of the Distribution of A Newly Characterized Receptor for Advanced Glycation End-Products in Tissues. *American Journal of Pathology* **143**:1699-1712.
249. **Pichiule, P., J. C. Chavez, A. M. Schmidt, and S. J. Vannucci.** 2007. Hypoxia inducible factor-1 mediates neuronal expression of the receptor for advanced glycation endproducts following hypoxia/Ischemia. *Journal of Biological Chemistry* **M706407200**.
250. **Ritthaler, U., Y. Deng, Y. Zhang, J. Greten, M. Abel, B. Sido, J. Allenberg, G. Otto, H. Roth, A. Bierhaus, R. Ziegler, A. M. Schmidt, R. Waldherr, P. Wahl, D. M. Stern, and P. P. Nawroth.** 1995. Expression of Receptors for Advanced Glycation End-Products in Peripheral Occlusive Vascular-Disease. *American Journal of Pathology* **146**:688-694.
251. **Tanji, N., G. S. Markowitz, C. F. Fu, T. Kislinger, A. Taguchi, M. Pischetsrieder, D. Stern, A. M. Schmidt, and V. D. D'Agati.** 2000. Expression of advanced glycation end products and their cellular receptor RAGE in diabetic nephropathy and nondiabetic renal disease. *Journal of the American Society of Nephrology* **11**:1656-1666.
252. **Pachydaki, S. I., S. R. Tari, S. E. Lee, W. C. Ma, J. J. Tseng, A. A. Sosunov, G. Cataldergirmen, N. Scarmeas, C. Caspersen, S. Chang, W. M. Schiff, A. M. Schmidt, and G. R. Barile.** 2006. Upregulation of RAGE and its ligands in proliferative retinal disease. *Experimental Eye Research* **82**:807-815.
253. **Ikeda, K., T. Higashi, H. Sano, Y. Jinnouchi, M. Yoshida, T. Araki, S. Ueda, and S. Horiuchi.** 1996. N-epsilon-(carboxymethyl)lysine protein adduct is a major immunological epitope in proteins modified with advanced glycation end products of the Maillard reaction. *Biochemistry* **35**:8075-8083.
254. **Reddy, S., J. Bichler, K. J. WellsKnecht, S. R. Thorpe, and J. W. Baynes.** 1995. N-Epsilon-(Carboxymethyl)Lysine Is A Dominant Advanced Glycation End-Product (Age) Antigen in Tissue Proteins. *Biochemistry* **34**:10872-10878.
255. **Yan, S. D., X. Chen, J. Fu, M. Chen, H. J. Zhu, A. Roher, T. Slattery, L. Zhao, M. Nagashima, J. Morser, A. Migheli, P. Nawroth, D. Stern, and A. M. Schmidt.** 1996. RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* **382**:685-691.
256. **Yan, S. D., H. J. Zhu, A. P. Zhu, A. Golabek, H. Du, A. Roher, J. Yu, C. Soto, A. M. Schmidt, D. Stern, and M. Kindy.** 2000. Receptor-dependent cell stress and amyloid accumulation in systemic amyloidosis. *Nature Medicine* **6**:643-651.
257. **Hofmann, M. A., S. Drury, C. F. Fu, W. Qu, A. Taguchi, Y. Lu, C. Avila, N. Kambham, A. Bierhaus, P. Nawroth, M. F. Neurath, T. Slattery, D. Beach, J. McClary, M. Nagashima, J. Morser, D. Stern, and A. M. Schmidt.** 1999. RAGE mediates a novel proinflammatory axis: A central cell surface receptor for S100/calgranulin polypeptides. *Cell* **97**:889-901.
258. **Hori, O., J. Brett, T. Slattery, R. Cao, J. H. Zhang, J. X. Chen, M. Nagashima, E. R. Lundh, S. Vijay, D. Nitecki, J. Morser, D. Stern, and A. M. Schmidt.** 1995. The Receptor for Advanced Glycation End-Products (Rage) Is A Cellular-Binding Site for Amphoterin - Mediation of Neurite Outgrowth and Coexpression of Rage and Amphoterin in the Developing Nervous-System. *Journal of Biological Chemistry* **270**:25752-25761.
259. **Hori, O., R. Cao, J. Brett, T. Slattery, M. Nagashima, D. Nitecki, J. Morser, D. Stern, and A. M. Schmidt.** 1995. The Receptor for Advanced Glycation Endproducts (Rage) Is A Cell-Surface Receptor for Amphoterin in the Developing Central-Nervous-System (Cns) to Promote Neurite Outgrowth. *Faseb Journal* **9**:A382.
260. **Chavakis, T., A. Bierhaus, N. Al Fakhri, D. Schneider, S. Witte, T. Linn, M. Nagashima, J. Morser, B. Arnold, K. T. Preissner, and P. P. Nawroth.** 2003. The Pattern Recognition Receptor (RAGE) Is a Counterreceptor for Leukocyte Integrins: A Novel Pathway for Inflammatory Cell Recruitment. *J.Exp.Med.* **198**:1507-1515.
261. **Lander, H. M., J. M. Tauras, J. S. Ogiste, O. Hori, R. A. Moss, and A. M. Schmidt.** 1997. Activation of the receptor for advanced glycation end products triggers a p21(ras)-dependent mitogen-activated protein kinase pathway regulated by oxidant stress. *Journal of Biological Chemistry* **272**:17810-17814.
262. **Wautier, M. P., O. Chappay, S. Corda, D. M. Stern, A. M. Schimidt, and J. L. Wautier.** 2001. Activation of NADPH oxidase by AGE links oxidant stress to altered gene expression via RAGE. *American Journal of Physiology-Endocrinology and Metabolism* **280**:E685-E694.
263. **Cortizo, A. M., M. G. Lettieri, D. A. Barrio, N. Mercer, S. B. Etcheverry, and A. D. McCarthy.** 2003. Advanced glycation end-products (AGEs) induce concerted changes in the osteoblastic expression of their receptor RAGE and in the activation of extracellular signal-regulated kinases (ERK). *Molecular and Cellular Biochemistry* **250**: 1-10.

264. Li, J. H., W. S. Wang, X. R. Huang, M. Oldfield, A. M. Schmidt, M. E. Cooper, and H. Y. Lan. 2004. Advanced glycation end products induce tubular epithelial-myofibroblast transition through the RAGE-ERK1/2 MAP kinase signaling pathway. *American Journal of Pathology* **164**:1389-1397.
265. Shanmugam, N., Y. S. Kim, L. Lanting, and R. Natarajan. 2003. Regulation of cyclooxygenase-2 expression in monocytes by ligation of the receptor for advanced glycation end products. *Journal of Biological Chemistry* **278**:34834-34844.
266. Sorci, G., F. Riuzzi, A. L. Agneletti, C. Marchetti, and R. Donato. 2004. S100B causes apoptosis in a myoblast cell line in a RAGE-independent manner. *Journal of Cellular Physiology* **199**:274-283.
267. Ishihara, K., K. Tsutsumi, S. Kawane, M. Nakajima, and T. Kasaoka. 2003. The receptor for advanced glycation end-products (RAGE) directly binds to ERK by a D-domain-like docking site. *Febs Letters* **550**:107-113.
268. Taguchi, A., D. C. Blood, G. del Toro, A. Canet, D. C. Lee, W. Qu, N. Tanji, Y. Lu, E. Lalla, C. F. Fu, M. A. Hofmann, T. Kislinger, M. Ingram, A. Lu, H. Tanaka, O. Hori, S. Ogawa, D. M. Stern, and A. M. Schmidt. 2000. Blockade of RAGE-amphoterin signalling suppresses tumour growth and metastases. *Nature* **405**:354-360.
269. Yeh, C. H., L. T. Sturgis, S. J. Haidacher, X. N. Zhang, S. J. Sherwood, M. T. Crow, O. Juhasz, R. G. Tilton, and L. Denner. 2000. Requirement for p38 mitogen-activated protein Kinase in RAGE-Mediated NF-kappa B transcriptional activation and cytokine secretion. *Diabetes* **49**:A337-A338.
270. Yeh, C. H., L. Sturgis, J. Haidacher, X. N. Zhang, S. J. Sherwood, R. J. Bjercke, O. Juhasz, M. T. Crow, R. G. Tilton, and L. Denner. 2001. Requirement for p38 and p44/p42 mitogen-activated protein kinases in RAGE-mediated nuclear factor-kappa B transcriptional activation and cytokine secretion. *Diabetes* **50**:1495-1504.
271. Brizzi, M. F., P. Dentelli, A. Rosso, C. Calvi, R. Gambino, M. Cassader, G. Salvidio, G. Deferrari, G. Camussi, L. Pegoraro, G. Pagano, and P. Cavallo-Perin. 2004. RAGE- and TGF-beta receptor-mediated signals converge on STAT5 and P21(waf) to control cell-cycle progression of mesangial cells: a possible role in the development and progression of diabetic nephropathy. *Faseb Journal* **18**:1249-+.
272. Huang, J. S., J. Y. Guh, H. C. Chen, W. C. Hung, Y. H. Lai, and L. Y. Chuang. 2001. Role of receptor for advanced glycation end-product (RAGE) and the JAK/STAT-signaling pathway in AGE-induced collagen production in NRK-49F cells. *Journal of Cellular Biochemistry* **81**:102-113.
273. Sakaguchi, T., S. F. Yan, S. Du Yan, D. Belov, L. L. Rong, M. Sousa, M. Andrassy, S. P. Marso, S. Duda, B. Arnold, B. Liliensiek, P. P. Nawroth, D. M. Stern, A. M. Schmidt, and Y. Naka. 2003. Central role of RAGE-dependent neointimal expansion in arterial restenosis. *Journal of Clinical Investigation* **111**:959-972.
274. Dukic-Stefanovic, S., J. Gasic-Milenkovic, W. Deuther-Conrad, and G. Munch. 2003. Signal transduction pathways in mouse microglia N-11 cells activated by advanced glycation endproducts (AGEs). *Journal of Neurochemistry* **87**:44-55.
275. Rohleder, N., A. Bierhaus, J. M. Wolf, P. P. Nawroth, and C. Kirschbaum. 2001. Psychosocial stress rapidly induces activation of the transcription factor NF-kappa B in healthy adults. *Psychosomatic Medicine* **63**:187.
276. Yeh, C. H., L. T. Sturgis, S. J. Haidacher, X. N. Zhang, S. J. Sherwood, M. T. Crow, O. Juhasz, R. G. Tilton, and L. Denner. 2000. Requirement for p38 mitogen-activated protein Kinase in RAGE-Mediated NF-kappa B transcriptional activation and cytokine secretion. *Diabetes* **49**:A337-A338.
277. Li, J. F. and A. M. Schmidt. 1997. Characterization and functional analysis of the promoter of RAGE, the receptor for advanced glycation end products. *Journal of Biological Chemistry* **272**:16498-16506.
278. Tanaka, N., H. Yonekura, S. Yamagishi, H. Fujimori, Y. Yamamoto, and H. Yamamoto. 2000. The receptor for advanced glycation end products is induced by the glycation products themselves and tumor necrosis factor-alpha through nuclear Factor-kappa B, and by 17 beta-estradiol through Sp-1 in human vascular endothelial cells. *Journal of Biological Chemistry* **275**:25781-25790.
279. Bierhaus, A., P. M. Humpert, M. Morcos, T. Wendt, T. Chavakis, B. Arnold, D. M. Stern, and P. P. Nawroth. 2005. Understanding RAGE, the receptor for advanced glycation end products. *Journal of Molecular Medicine-Jmm* **83**:876-886.
280. Bierhaus, A., P. M. Humpert, D. M. Stern, B. Arnold, and P. P. Nawroth. 2005. Advanced glycation end product receptor - Mediated cellular dysfunction. *Maillard Reaction: Chemistry at the Interface of Nutrition, Aging, and Disease* **1043**:676-680.
281. Goldin, A., J. A. Beckman, A. M. Schmidt, and M. A. Creager. 2006. Advanced glycation end products - Sparking the development of diabetic vascular injury. *Circulation* **114**:597-605.
282. Hudson, B. I., T. Wendt, L. G. Bucciarelli, L. L. Rong, Y. Naka, S. F. Yan, and A. M. Schmidt. 2005. Diabetic vascular disease: It's all the RAGE. *Antioxidants & Redox Signaling* **7**:1588-1600.
283. Stern, D. M., S. Du Yan, S. F. Yan, and A. M. Schmidt. 2002. Receptor for advanced glycation endproducts (RAGE) and the complications of diabetes. *Ageing Research Reviews* **1**:1-15.
284. Adami, C., R. Bianchi, G. Pula, and R. Donato. 2004. S100B-stimulated NO production by BV-2 microglia is independent of RAGE transducing activity but dependent on RAGE extracellular domain. *Biochimica et Biophysica Acta-Molecular Cell Research* **1742**:169-177.
285. Sajithlal, G., H. Huttunen, H. Rauvala, and G. Munch. 2002. Receptor for advanced glycation end products plays a more important role in cellular survival than in neurite outgrowth during retinoic acid-induced differentiation of neuroblastoma cells. *Journal of Biological Chemistry* **277**:6888-6897.
286. Riuzzi, F., G. Sorci, and R. Donato. 2006. The amphoterin (HMGB1)/receptor for advanced glycation end products (RAGE) pair modulates myoblast proliferation, apoptosis, adhesiveness, migration, and invasiveness - Functional inactivation of rage in L6 myoblasts results in tumor formation in vivo. *Journal of Biological Chemistry* **281**:8242-8253.

287. **Sorci, G., F. Riuizi, C. Arcuri, I. Giambanco, and R. Donato.** 2004. Amphoterin stimulates myogenesis and counteracts the antimyogenic factors basic fibroblast growth factor and S100B via RAGE binding. *Molecular and Cellular Biology* **24**:4880-4894.
288. **Riuizi, F., G. Sorci, and R. Donato.** 2007. RAGE expression in rhabdomyosarcoma cells results in myogenic differentiation and reduced proliferation, migration, invasiveness, and tumor growth. *American Journal of Pathology* **171**:947-961.
289. **Bartling, B., H. S. Hofmann, B. Weigle, R. E. Silber, and A. Simm.** 2005. Down-regulation of the receptor for advanced glycation end-products (RAGE) supports non-small cell lung carcinoma. *Carcinogenesis* **26**:293-301.
290. **Bartling, B., N. Demling, R. E. Silber, and A. Simm.** 2006. Proliferative stimulus of lung fibroblasts on lung cancer cells is impaired by the receptor for advanced glycation end-products. *American Journal of Respiratory Cell and Molecular Biology* **34**:83-91.
291. **Schmidt, A., B. Kuhla, K. Bigl, G. Munch, and T. Arendt.** 2007. Cell cycle related signaling in neuro2a cells proceeds via the receptor for advanced glycation end products. *Journal of Neural Transmission* **114**:1413-1424.
292. **Dumitriu, I. E., P. Baruah, B. Valentinis, R. E. Voll, M. Herrmann, P. P. Nawroth, B. Arnold, M. E. Bianchi, A. A. Manfredi, and P. Rovere-Querini.** 2005. Release of high mobility group box 1 by dendritic cells controls T cell activation via the receptor for advanced glycation end products. *Journal of Immunology* **174**:7506-7515.
293. **Dumitriu, I. E., M. E. Bianchi, M. Bacci, A. A. Manfredi, and P. Rovere-Querini.** 2007. The secretion of HMGB1 is required for the migration of maturing dendritic cells. *Journal of Leukocyte Biology* **81**:84-91.
294. **Huttunen, H. J., J. Kuja-Panula, G. Sorci, A. L. Agneletti, R. Donato, and H. Rauvala.** 2000. Coregulation of neurite outgrowth and cell survival by amphoterin and S100 proteins through receptor for advanced glycation end products (RAGE) activation. *Journal of Biological Chemistry* **275**:40096-40105.
295. **Arumugam, T., D. M. Simeone, A. M. Schmidt, and C. D. Logsdon.** 2004. S100P stimulates cell proliferation and survival via receptor for activated glycation end products (RAGE). *Journal of Biological Chemistry* **279**:5059-5065.
296. **Fuentes, M. K., S. S. Nigavekar, T. Arumugam, C. D. Logsdon, A. M. Schmidt, J. C. Park, and E. H. Huang.** 2007. RAGE activation by S100P in colon cancer stimulates growth, migration, and cell signaling pathways. *Diseases of the Colon & Rectum* **50**:1230-1240.
297. **Alikhani, M., C. M. MacLellan, M. Raptis, S. Vora, P. C. Trackman, and D. T. Graves.** 2007. Advanced glycation end products induce apoptosis in fibroblasts through activation of ROS, MAP kinases, and the FOXO1 transcription factor. *American Journal of Physiology-Cell Physiology* **292**:C850-C856.
298. **Alikhani, Z. B., M. Alikhani, C. M. Boyd, K. Nagao, P. C. Trackman, and D. T. Graves.** 2005. Advanced glycation end products enhance expression of pro-apoptotic genes and stimulate fibroblast apoptosis through cytoplasmic and mitochondrial pathways. *Journal of Biological Chemistry* **280**:12087-12095.
299. **Conlon, I. and M. Raff.** 1999. Size control in animal development. *Cell* **96**:235-244.
300. **Raff, M. C.** 1992. Social Controls on Cell-Survival and Cell-Death. *Nature* **356**:397-400.
301. **Akashi, K., M. Kondo, U. vonFreedenJeffry, R. Murray, and I. L. Weissman.** 1997. Bcl-2 rescues T lymphopoiesis in interleukin-7 receptor-deficient mice. *Cell* **89**:1033-1041.
302. **Chi, M. M. Y., J. Pingsterhaus, M. Carayannopoulos, and K. H. Moley.** 2000. Decreased glucose transporter expression triggers BAX-dependent apoptosis in the murine blastocyst. *Journal of Biological Chemistry* **275**:40252-40257.
303. **Plas, D. R., S. Talapatra, A. L. Edinger, J. C. Rathmell, and C. B. Thompson.** 2001. Akt and Bcl-x(L) promote growth factor-independent survival through distinct effects on mitochondrial physiology. *Journal of Biological Chemistry* **276**:12041-12048.
304. **Rathmell, J. C., M. G. Vander Heiden, M. H. Harris, K. A. Frauwirth, and C. B. Thompson.** 2000. In the absence of extrinsic signals, nutrient utilization by lymphocytes is insufficient to maintain either cell size or viability. *Molecular Cell* **6**:683-692.
305. **Rathmell, J. C., C. J. Fox, D. R. Plas, P. S. Hammerman, R. M. Cinalli, and C. B. Thompson.** 2003. Akt-directed glucose metabolism can prevent Bax conformation change and promote growth factor-independent survival. *Molecular and Cellular Biology* **23**:7315-7328.
306. **Vander Heiden, M. G., D. R. Plas, J. C. Rathmell, C. J. Fox, M. H. Harris, and C. B. Thompson.** 2001. Growth factors can influence cell growth and survival through effects on glucose metabolism. *Molecular and Cellular Biology* **21**:5899-5912.
307. **Danial, N. N., C. F. Gramm, L. Scorrano, C. Y. Zhang, S. Krauss, A. M. Ranger, S. R. Datta, M. E. Greenberg, L. J. Licklider, B. B. Lowell, S. P. Gygi, and S. J. Korsmeyer.** 2003. BAD and glucokinase reside in a mitochondrial complex that integrates glycolysis and apoptosis. *Nature* **424**:952-956.
308. **Barata, J. T., A. Silva, J. G. Brandao, L. M. Nadler, A. A. Cardoso, and V. A. Boussiotis.** 2004. Activation of P13K is indispensable for interleukin 7-mediated viability, proliferation, glucose use, and growth of T cell acute lymphoblastic leukemia cells. *Journal of Experimental Medicine* **200**:659-669.
309. **Craddock, B. L., E. A. Orchiston, H. J. Hinton, and M. J. Welham.** 1999. Dissociation of apoptosis from proliferation, protein kinase B activation, and BAD phosphorylation in interleukin-3-mediated phosphoinositide 3-kinase signaling. *Journal of Biological Chemistry* **274**:10633-10640.
310. **Minshall, C., S. Arkins, G. G. Freund, and K. W. Kelley.** 1996. Requirement for phosphatidylinositol 3'-kinase to protect hemopoietic progenitors against apoptosis depends upon the extracellular survival factor. *J.Immunol.* **156**:939-947.
311. **Plas, D. R. and C. B. Thompson.** 2005. Akt-dependent transformation: there is more to growth than just

- surviving. *Oncogene* **24**:7435-7442.
312. **Scheid, M. P., R. W. Lauener, and V. Duronio.** 1995. Role of Phosphatidylinositol 8-Oh-Kinase Activity in the Inhibition of Apoptosis in Hematopoietic-Cells - Phosphatidylinositol 3-Oh-Kinase Inhibitors Reveal A Difference in Signaling Between Interleukin-3 and Granulocyte-Macrophage Colony-Stimulating Factor. *Biochemical Journal* **312**:159-162.
 313. **Songyang, Z., D. Baltimore, L. C. Cantley, D. R. Kaplan, and T. F. Franke.** 1997. Interleukin 3-dependent survival by the Akt protein kinase. *Proc.Natl.Acad.Sci.U.S.A* **94**:11345-11350.
 314. **Chang, F., J. T. Lee, P. M. Navolanic, L. S. Steelman, J. G. Shelton, W. L. Blalock, R. A. Franklin, and J. A. McCubrey.** 2003. Involvement of PI3K/Akt pathway in cell cycle progression, apoptosis, and neoplastic transformation: a target for cancer chemotherapy. *Leukemia* **17**:590-603.
 315. **Manning, B. D. and L. C. Cantley.** 2007. AKT/PKB signaling: Navigating downstream. *Cell* **129**:1261-1274.
 316. **Raff, M. C., B. A. Barres, J. F. Burne, H. S. R. Coles, Y. Ishizaki, and M. D. Jacobson.** 1994. Programmed Cell-Death and the Control of Cell-Survival. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **345**:265-268.
 317. **Adams, J. M. and S. Cory.** 1998. The Bcl-2 protein family: Arbiters of cell survival. *Science* **281**:1322-1326.
 318. **Reed, J. C., K. S. Doctor, and A. Godzik.** 2004. The domains of apoptosis: a genomics perspective. *Sci.STKE*. **2004**:re9.
 319. **Suzuki, M., R. J. Youle, and N. Tjandra.** 2000. Structure of Bax: Coregulation of dimer formation and intracellular localization. *Cell* **103**:645-654.
 320. **Wei, M. C., W. X. Zong, E. H. Y. Cheng, T. Lindsten, V. Panoutsakopoulou, A. J. Ross, K. A. Roth, G. R. MacCregor, C. B. Thompson, and S. J. Korsmeyer.** 2001. Proapoptotic BAX and BAK: A requisite gateway to mitochondrial dysfunction and death. *Science* **292**:727-730.
 321. **Chen, L., S. N. Willis, A. Wei, B. J. Smith, J. I. Fletcher, M. G. Hinds, P. M. Colman, C. L. Day, J. M. Adams, and D. C. S. Huang.** 2005. Differential targeting of prosurvival Bcl-2 proteins by their BH3-only ligands allows complementary apoptotic function. *Molecular Cell* **17**:393-403.
 322. **Huang, D. C. S. and A. Strasser.** 2000. BH3-Only proteins - Essential initiators of apoptotic cell death. *Cell* **103**: 839-842.
 323. **Kuwana, T., L. Bouchier-Hayes, J. E. Chipuk, C. Bonzon, B. A. Sullivan, D. R. Green, and D. D. Newmeyer.** 2005. BH3 domains of BH3-only proteins differentially regulate bax-mediated mitochondrial membrane permeabilization both directly and indirectly. *Molecular Cell* **17**:525-535.
 324. **Letai, A., M. C. Bassik, L. D. Walensky, M. D. Sorcinelli, S. Weiler, and S. J. Korsmeyer.** 2002. Distinct BH3 domains either sensitize or activate mitochondrial apoptosis, serving as prototype cancer therapeutics. *Cancer Cell* **2**:183-192.
 325. **Willis, S. N., L. Chen, G. Dewson, A. Wei, E. Naik, J. I. Fletcher, J. M. Adams, and D. C. S. Huang.** 2005. Proapoptotic Bak is sequestered by Mcl-1 and Bcl-x(L), but not Bcl-2, until displaced by BH3-only proteins. *Genes & Development* **19**:1294-1305.
 326. **Cheng, E. H. Y. A., M. C. Wei, S. Weiler, R. A. Flavell, T. W. Mak, T. Lindsten, and S. J. Korsmeyer.** 2001. BCL-2, BCL-X-L sequester BH3 domain-only molecules preventing BAX- and BAK-mediated mitochondrial apoptosis. *Molecular Cell* **8**:705-711.
 327. **Maurer, U., C. Charvet, A. S. Wagman, E. Dejardin, and D. R. Green.** 2006. Glycogen synthase kinase-3 regulates mitochondrial outer membrane permeabilization and apoptosis by destabilization of MCL-1. *Molecular Cell* **21**:749-760.
 328. **Gustafsson, A. B. and R. A. Gottlieb.** 2007. Bcl-2 family members and apoptosis, taken to heart. *American Journal of Physiology-Cell Physiology* **292**:C45-C51.
 329. **Green, D. R.** 2005. Apoptotic pathways: Ten minutes to dead. *Cell* **121**:671-674.
 330. **Li, P., D. Nijhawan, I. Budihardjo, S. M. Srinivasula, M. Ahmad, E. S. Alnemri, and X. D. Wang.** 1997. Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. *Cell* **91**:479-489.
 331. **Wang, X. D.** 2001. The expanding role of mitochondria in apoptosis. *Genes & Development* **15**:2922-2933.
 332. **Ranger, A. M., J. P. Zha, H. Harada, S. R. Datta, N. N. Danial, A. P. Gilmore, J. L. Kutok, M. M. Le Beau, M. E. Greenberg, and S. J. Korsmeyer.** 2003. Bad-deficient mice develop diffuse large B cell lymphoma. *Proceedings of the National Academy of Sciences of the United States of America* **100**:9324-9329.
 333. **Datta, S. R., A. M. Ranger, M. Z. Lin, J. F. Sturgill, Y. C. Ma, C. W. Cowan, P. Dikkes, S. J. Korsmeyer, and M. E. Greenberg.** 2002. Survival factor-mediated BAD phosphorylation raises the mitochondrial threshold for apoptosis. *Developmental Cell* **3**:631-643.
 334. **Zha, J. P., H. Harada, K. Osipov, J. Jockel, G. Waksman, and S. J. Korsmeyer.** 1997. BH3 domain of BAD is required for heterodimerization with BCL-X-L and pro-apoptotic activity. *Journal of Biological Chemistry* **272**: 24101-24104.
 335. **Zha, J. P., H. Harada, E. Yang, J. Jockel, and S. J. Korsmeyer.** 1996. Serine phosphorylation of death agonist BAD in response to survival factor results in binding to 14-3-3 not BGL-X(L). *Cell* **87**:619-628.
 336. **Ekert, P. G., A. M. Jabbour, A. Manoharan, J. E. Heraud, J. Yu, M. Pakusch, E. M. Michalak, P. N. Kelly, B. Callus, T. Kiefer, A. Verhagen, J. Silke, A. Strasser, C. Borner, and D. L. Vaux.** 2006. Cell death provoked by loss of interleukin-3 signaling is independent of Bad, Bim, and PI3 kinase, but depends in part on Puma. *Blood* **108**:1461-1468.
 337. **Erlacher, M., V. Labi, C. Manzi, G. Bock, A. Tzankov, G. Hacker, E. Michalak, A. Strasser, and A. Villunger.** 2006. Puma cooperates with Bim, the rate-limiting BH3-only protein in cell death during lymphocyte development, in apoptosis induction. *Journal of Experimental Medicine* **203**:2939-2951.

338. **Harada, H., B. Quearry, A. Ruiz-Vela, and S. J. Korsmeyer.** 2004. Survival factor-induced extracellular signal-regulated kinase phosphorylates BIM, inhibiting its association with BAX and proapoptotic activity. *Proceedings of the National Academy of Sciences of the United States of America* **101**:15313-15317.
339. **You, H., M. Pellegrini, K. Tsuchihara, K. Yamamoto, G. Hacker, M. Erlacher, A. Villunger, and T. W. Mak.** 2006. FOXO3a-dependent regulation of Puma in response to cytokine/growth factor withdrawal. *Journal of Experimental Medicine* **203**:1657-1663.
340. **Donovan, M. and T. G. Cotter.** 2004. Control of mitochondrial integrity by Bcl-2 family members and caspase-independent cell death. *Biochimica et Biophysica Acta-Molecular Cell Research* **1644**:133-147.
341. **Elmore, S.** 2007. Apoptosis: A review of programmed cell death. *Toxicologic Pathology* **35**:495-516.
342. **Datta, S. R., H. Dudek, X. Tao, S. Masters, H. A. Fu, Y. Gotoh, and M. E. Greenberg.** 1997. Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell* **91**:231-241.
343. **Datta, S. R., A. M. Ranger, M. Z. Lin, J. F. Sturgill, Y. C. Ma, C. W. Cowan, P. Dikkes, S. J. Korsmeyer, and M. E. Greenberg.** 2002. Survival factor-mediated BAD phosphorylation raises the mitochondrial threshold for apoptosis. *Developmental Cell* **3**:631-643.
344. **delPeso, L., M. GonzalezGarcia, C. Page, R. Herrera, and G. Nunez.** 1997. Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt. *Science* **278**:687-689.
345. **Scheid, M. P. and V. Duronio.** 1998. Dissociation of cytokine-induced phosphorylation of bad and activation of PKB/akt: Involvement of MEK upstream of bad phosphorylation. *Proceedings of the National Academy of Sciences of the United States of America* **95**:7439-7444.
346. **Bonni, A., A. Brunet, A. E. West, S. R. Datta, M. A. Takasu, and M. E. Greenberg.** 1999. Cell survival promoted by the Ras-MAPK signaling pathway by transcription-dependent and -independent mechanisms. *Science* **286**:1358-1362.
347. **Chiang, C. W., G. Harris, C. Ellig, S. C. Masters, R. Subramanian, S. Shenolikar, B. E. Wadzinski, and E. Yang.** 2001. Protein phosphatase 2A activates the proapoptotic function of BAD in interleukin-3-dependent lymphoid cells by a mechanism requiring 14-3-3 dissociation. *Blood* **97**:1289-1297.
348. **Chiang, C. W., C. Kanies, K. W. Kim, W. B. Fang, C. Parkhurst, M. H. Xie, T. Henry, and E. Yang.** 2003. Protein phosphatase 2A dephosphorylation of phosphoserine 112 plays the gatekeeper role for BAD-mediated apoptosis. *Molecular and Cellular Biology* **23**:6350-6362.
349. **Harada, H., J. S. Andersen, M. Mann, N. Terada, and S. J. Korsmeyer.** 2001. p70S6 kinase signals cell survival as well as growth, inactivating the pro-apoptotic molecule BAD. *Proceedings of the National Academy of Sciences of the United States of America* **98**:9666-9670.
350. **Masters, S. C., H. Z. Yang, S. R. Datta, M. E. Greenberg, and H. Fu.** 2001. 14-3-3 inhibits bad-induced cell death through interaction with serine-136. *Molecular Pharmacology* **60**:1325-1331.
351. **Schurmann, A., A. F. Mooney, L. C. Sanders, M. A. Sells, H. G. Wang, J. C. Reed, and G. M. Bokoch.** 2000. p21-activated kinase 1 phosphorylates the death agonist Bad and protects cells from apoptosis. *Molecular and Cellular Biology* **20**:453-461.
352. **Tan, Y., M. R. Demeter, H. Ruan, and M. J. Comb.** 2000. BAD Ser-155 phosphorylation regulates BAD/Bcl-XL interaction and cell survival. *Journal of Biological Chemistry* **275**:25865-25869.
353. **Zhou, X. M., Y. M. Liu, G. Payne, R. J. Lutz, and T. Chittenden.** 2000. Growth factors inactivate the cell death promoter BAD by phosphorylation of its BH3 domain on Ser(155). *Journal of Biological Chemistry* **275**:25046-25051.
354. **Scheid, M. P., K. M. Schubert, and V. Duronio.** 1999. Regulation of bad phosphorylation and association with Bcl-x(L) by the MAPK/Erk kinase. *Journal of Biological Chemistry* **274**:31108-31113.
355. **Brunet, A., A. Bonni, M. J. Zigmond, M. Z. Lin, P. Juo, L. S. Hu, M. J. Anderson, K. C. Arden, J. Blenis, and M. E. Greenberg.** 1999. Akt promotes cell survival by phosphorylating and inhibiting a forkhead transcription factor. *Cell* **96**:857-868.
356. **Brunet, A., J. Park, H. Tran, L. S. Hu, B. A. Hemmings, and M. E. Greenberg.** 2001. Protein kinase SGK mediates survival signals by phosphorylating the forkhead transcription factor FKHL1 (FOXO3a). *Molecular and Cellular Biology* **21**:952-965.
357. **Tang, E. D., G. Nunez, F. G. Barr, and K. L. Guan.** 1999. Negative regulation of the forkhead transcription factor FKHR by Akt. *Journal of Biological Chemistry* **274**:16741-16746.
358. **Dumon, S., S. C. R. Santos, F. Debierre-Grockiego, V. Gouilleux-Gruart, L. Cocault, C. Boucheron, P. Mollat, S. Gisselbrecht, and F. Gouilleux.** 1999. IL-3 dependent regulation of Bcl-x(L) gene expression by STAT5 in a bone marrow derived cell line. *Oncogene* **18**:4191-4199.
359. **Jones, R. G., M. Parsons, M. Bonnard, V. S. F. Chan, W. C. Yeh, J. R. Woodgett, and P. S. Ohashi.** 2000. Protein kinase B regulates T lymphocyte survival, nuclear factor kappa B activation, and Bcl-X-L levels in vivo. *Journal of Experimental Medicine* **191**:1721-1733.
360. **Kuribara, R., T. Kinoshita, A. Miyajima, T. Shinjyo, T. Yoshihara, T. Inukai, K. Ozawa, A. T. Look, and T. Inaba.** 1999. Two distinct interleukin-3-mediated signal pathways, Ras-NFIL3 (E4BP4) and Bcl-x(L), regulate the survival of murine pro-B lymphocytes. *Molecular and Cellular Biology* **19**:2754-2762.
361. **Leverrier, Y., J. Thomas, G. R. Perkins, M. Mangeney, M. K. L. Collins, and J. Marvel.** 1997. In bone marrow derived Baf-3 cells, inhibition of apoptosis by IL-3 is mediated by two independent pathways. *Oncogene* **14**:425-430.
362. **Leverrier, Y., J. Thomas, A. L. Mathieu, W. Low, B. Blanquie, and J. Marvel.** 1999. Role of PI3-kinase in Bcl-X induction and apoptosis inhibition mediated by IL-3 or IGF-1 in Baf-3 cells. *Cell Death and Differentiation* **6**:290-296.
363. **Wang, J. M., M. Z. Lai, and H. F. Yang-Yen.** 2003. Interleukin-3 stimulation of mcl-1 gene transcription involves

- activation of the PU.1 transcription factor through a p38 mitogen-activated protein kinase-dependent pathway. *Molecular and Cellular Biology* **23**:1896-1909.
364. **Zong, W. X., L. C. Edelstein, C. L. Chen, J. Bash, and C. Gelinas.** 1999. The prosurvival Bcl-2 homolog Bfl-1/A1 is a direct transcriptional target of NF-kappa B that blocks TNF alpha-induced apoptosis. *Genes & Development* **13**:382-387.
 365. **Nakata, S., I. Matsumura, H. Tanaka, S. Ezoe, Y. Satoh, J. Ishikawa, T. Era, and Y. Kanakura.** 2004. NF-kappa B family proteins participate in multiple steps of hematopoiesis through elimination of reactive oxygen species. *Journal of Biological Chemistry* **279**:55578-55586.
 366. **Romashkova, J. A. and S. S. Makarov.** 1999. NF-kappaB a target of AKT in anti-apoptotic PDGF signalling NF-kappaB a target of AKT in anti-apoptotic PDGF signalling. *Nature* **401**:86.
 367. **Allan, L. A., N. Morrice, S. Brady, G. Magee, S. Pathak, and P. R. Clarke.** 2003. Inhibition of caspase-9 through phosphorylation at Thr 125 by ERK MAPK. *Nature Cell Biology* **5**:647-U45.
 368. **Allan, L. A. and P. R. Clarke.** 2007. Phosphorylation of caspase-9 by CDK1/cyclin B1 protects mitotic cells against apoptosis. *Molecular Cell* **26**:301-310.
 369. **Martin, M. C., L. A. Allan, M. Lickrish, C. Sampson, N. Morrice, and P. R. Clarke.** 2005. Protein kinase A regulates caspase-9 activation by Apaf-1 downstream of cytochrome c. *Journal of Biological Chemistry* **280**:15449-15455.
 370. **Gottlob, K., N. Majewski, S. Kennedy, E. Kandel, R. B. Robey, and N. Hay.** 2001. Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. *Genes & Development* **15**:1406-1418.
 371. **Lum, J. J., D. E. Bauer, M. Kong, M. H. Harris, C. Li, T. Lindsten, and C. B. Thompson.** 2005. Growth factor regulation of autophagy and cell survival in the absence of apoptosis. *Cell* **120**:237-248.
 372. **Ciofani, M. and J. C. Zuniga-Pflucker.** 2005. Notch promotes survival of pre-T cells at the beta-selection checkpoint by regulating cellular metabolism. *Nature Immunology* **6**:881-888.
 373. **Khaled, A. R., W. Q. Li, J. Q. Huang, T. J. Fry, A. S. Khaled, C. L. Mackall, K. Muegge, H. A. Young, and S. K. Durum.** 2002. Bax deficiency partially corrects interleukin-7 receptor alpha deficiency. *Immunity* **17**:561-573.
 374. **Kondo, M., K. Akashi, J. Domen, K. Sugamura, and I. L. Weissman.** 1997. Bcl-2 rescues T lymphopoiesis, but not B or NK cell development, in common gamma chain-deficient mice. *Immunity* **7**:155-162.
 375. **Pellegrini, M., P. Bouillet, M. Robati, G. T. Belz, G. M. Davey, and A. Strasser.** 2004. Loss of Bim increases T cell production and function in interleukin 7 receptor-deficient mice. *Journal of Experimental Medicine* **200**:1189-1195.
 376. **Rathmell, J. C., E. A. Farkash, W. Gao, and C. B. Thompson.** 2001. IL-7 enhances the survival and maintains the size of naive T cells. *Journal of Immunology* **167**:6869-6876.
 377. **Bruno, P., A. Calastretti, M. Priulla, L. Asnaghi, F. Scarlatti, A. Nicolin, and G. Canti.** 2007. Cell survival under nutrient stress is dependent on metabolic conditions regulated by Akt and not by autophagic vacuoles. *Cellular Signalling* **19**:2118-2126.
 378. **Bauer, D. E., M. H. Harris, D. R. Plas, J. J. Lum, P. S. Hammerman, J. C. Rathmell, J. L. Riley, and C. B. Thompson.** 2004. Cytokine stimulation of aerobic glycolysis in hematopoietic cells exceeds proliferative demand. *Faseb Journal* **18**:1303-+.
 379. **Wieman, H. L., J. A. Wofford, and J. C. Rathmell.** 2007. Cytokine stimulation promotes glucose uptake via phosphatidylinositol-3 kinase/Akt regulation of Glut1 activity and trafficking. *Mol.Biol.Cell* **18**:1437-1446.
 380. **Kan, O., S. A. Baldwin, and A. D. Whetton.** 1994. Apoptosis Is Regulated by the Rate of Glucose-Transport in An Interleukin-3 Dependent Cell-Line. *Journal of Experimental Medicine* **180**:917-923.
 381. **Whetton, A. D., G. W. Bazill, and T. M. Dexter.** 1984. Hematopoietic-Cell Growth-Factor Mediates Cell-Survival Via Its Action on Glucose-Transport. *Embo Journal* **3**:409-413.
 382. **Bentley, J., D. Itchayan, K. Barnes, E. McIntosh, X. W. Tang, C. P. Downes, G. D. Holman, A. D. Whetton, P. J. Owen-Lynch, and S. A. Baldwin.** 2003. Interleukin-3-mediated cell survival signals include phosphatidylinositol 3-kinase-dependent translocation of the glucose transporter GLUT1 to the cell surface. *Journal of Biological Chemistry* **278**:39337-39348.
 383. **Wofford, J. A., H. L. Wieman, S. R. Jacobs, Y. Zhao, and J. C. Rathmell.** 2007. IL-7 promotes Glut1 trafficking and glucose uptake via STAT5-mediated activation of Akt to support T cell survival. *Blood*.
 384. **Edinger, A. L. and C. B. Thompson.** 2002. Akt maintains cell size and survival by increasing mTOR-dependent nutrient uptake. *Molecular Biology of the Cell* **13**:2276-2288.
 385. **Edinger, A. L.** 2005. Growth factors regulate cell survival by controlling nutrient transporter expression. *Biochemical Society Transactions* **33**:225-227.
 386. **Edinger, A. L.** 2007. Controlling cell growth and survival through regulated nutrient transporter expression. *Biochemical Journal* **406**:1-12.
 387. **Edinger, A. L.** 2005. Growth factors regulate cell survival by controlling nutrient transporter expression. *Biochem.Soc.Trans.* **33**:225-227.
 388. **Gonin-Giraud, S., A. L. Mathieu, S. Diocou, M. Tomkowiak, G. Delorme, and J. Marvel.** 2002. Decreased glycolytic metabolism contributes to but is not the inducer of apoptosis following IL-3-starvation. *Cell Death and Differentiation* **9**:1147-1157.
 389. **Burger, C., M. Wick, S. Brusselbach, and R. Muller.** 1994. Differential Induction of Metabolic Genes After Mitogen Stimulation and During Normal-Cell Cycle Progression. *Journal of Cell Science* **107**:241-252.
 390. **Printz, R. L., S. Koch, L. R. Potter, R. M. Odoherty, J. J. Tiesinga, S. Moritz, and D. K. Granner.** 1993. Hexokinase-II Messenger-Rna and Gene Structure, Regulation by Insulin, and Evolution. *Journal of Biological*

- Chemistry **268**:5209-5219.
391. **Robey, R. B., J. F. Ma, A. V. P. Santos, O. A. Noboa, P. E. Coy, and J. M. Bryson.** 2002. Regulation of mesangial cell hexokinase activity and expression by heparin-binding epidermal growth factor-like growth factor - Epidermal growth factors and phorbol esters increase glucose metabolism via a common mechanism involving classic mitogen-activated protein kinase pathway activation and induction of hexokinase II expression. *Journal of Biological Chemistry* **277**:14370-14378.
392. **Taneja, N., P. E. Coy, I. Lee, J. M. Bryson, and R. B. Robey.** 2004. Proinflammatory interleukin-1 cytokines increase mesangial cell hexokinase activity and hexokinase II isoform abundance. *American Journal of Physiology-Cell Physiology* **287**:C548-C557.
393. **Ahmad, A., S. Ahmad, B. K. Schneider, C. B. Allen, L. Y. Chang, and C. W. White.** 2002. Elevated expression of hexokinase II protects human lung epithelial-like A549 cells against oxidative injury. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **283**:L573-L584.
394. **Azoulay-Zohar, H., A. Israelson, S. Abu-Hamad, and V. Shoshan-Barmatz.** 2004. In self-defence: Hexokinase promotes voltage-dependent anion channel closure and prevents mitochondria-mediated apoptotic cell death. *Biochemical Journal* **377**:347-355.
395. **Bryson, J. M., P. E. Coy, K. Gottlob, N. Hay, and R. B. Robey.** 2002. Increased hexokinase activity, of either ectopic or endogenous origin, protects renal epithelial cells against acute oxidant-induced cell death. *Journal of Biological Chemistry* **277**:11392-11400.
396. **Majewski, N., V. Nogueira, R. B. Robey, and N. Hay.** 2004. Akt inhibits apoptosis downstream of BID cleavage via a glucose-dependent mechanism involving mitochondrial hexokinases. *Molecular and Cellular Biology* **24**: 730-740.
397. **Conricode, K. M. and R. S. Ochs.** 1990. Epidermal Growth-Factor and 12-O-Tetradecanoylphorbol 13-Acetate Stimulate Lactate Production and the Pentose-Phosphate Pathway in Freshly Isolated Rat Hepatocytes. *Journal of Biological Chemistry* **265**:20931-20937.
398. **Bertrand, L., D. R. Alessi, J. Deprez, M. Deak, E. Viaene, M. H. Rider, and L. Hue.** 1999. Heart 6-phosphofructo-2-kinase activation by insulin results from Ser-466 and Ser-483 phosphorylation and requires 3-phosphoinositide-dependent kinase-1, but not protein kinase B. *Journal of Biological Chemistry* **274**:30927-30933.
399. **Darville, M. I., I. V. Antoine, J. R. Mertensstrijthagen, V. J. Dupriez, and G. G. Rousseau.** 1995. An E2F-Dependent Late-Serum-Response Promoter in A Gene That Controls Glycolysis. *Oncogene* **11**:1509-1517.
400. **Joaquin, M., J. L. Rosa, R. Bartrons, and A. Tauler.** 1997. Expression of the F-type 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase mRNA during liver regeneration. *Biochim.Biophys.Acta* **1334**:256-260.
401. **Lefebvre, V., M. C. Mechin, M. P. Louckx, M. H. Rider, and L. Hue.** 1996. Signaling pathway involved in the activation of heart 6-phosphofructo-2-kinase by insulin. *Journal of Biological Chemistry* **271**:22289-22292.
402. **Rider, M. H., L. Bertrand, D. Vertommen, P. A. Michels, G. G. Rousseau, and L. Hue.** 2004. 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase: Head-to-head with a bifunctional enzyme that controls glycolysis. *Biochemical Journal* **381**:561-579.
403. **Gottlieb, E., S. M. Armour, and C. B. Thompson.** 2002. Mitochondrial respiratory control is lost during growth factor deprivation. *Proceedings of the National Academy of Sciences of the United States of America* **99**:12801-12806.
404. **Khaled, A. R., D. A. Reynolds, H. A. Young, C. B. Thompson, K. Muegge, and S. K. Durum.** 2001. Interleukin-3 withdrawal induces an early increase in mitochondrial membrane potential unrelated to the Bcl-2 family - Roles of intracellular pH, ADP transport, and F0F1-ATPase. *Journal of Biological Chemistry* **276**:6453-6462.
405. **Beutner, G., A. Ruck, B. Riede, and D. Brdiczka.** 1998. Complexes between porin, hexokinase, mitochondrial creatine kinase and adenylate translocator display properties of the permeability transition pore. Implication for regulation of permeability transition by the kinases. *Biochimica et Biophysica Acta-Biomembranes* **1368**:7-18.
406. **Crompton, M.** 1999. The mitochondrial permeability transition pore and its role in cell death. *Biochemical Journal* **341**:233-249.
407. **Desagher, S. and J. C. Martinou.** 2000. Mitochondria as the central control point of apoptosis. *Trends in Cell Biology* **10**:369-377.
408. **Majewski, N., V. Nogueira, P. Bhaskar, P. E. Coy, J. E. Skeen, K. Gottlob, N. S. Chandel, C. B. Thompson, R. B. Robey, and N. Hay.** 2004. Hexokinase-mitochondria interaction mediated by Akt is required to inhibit apoptosis in the presence or absence of Bax and Bak. *Molecular Cell* **16**:819-830.
409. **Pastorino, J. G., N. Shulga, and J. B. Hoek.** 2002. Mitochondrial binding of hexokinase II inhibits Bax-induced cytochrome c release and apoptosis. *Journal of Biological Chemistry* **277**:7610-7618.
410. **Maraskovsky, E., L. A. O'Reilly, M. Teepe, L. M. Corcoran, J. J. Peschon, and A. Strasser.** 1997. Bcl-2 can rescue T lymphocyte development in interleukin-7 receptor-deficient mice but not in mutant rag-1(-/-) mice. *Cell* **89**:1011-1019.
411. **Maraskovsky, E., J. J. Peschon, H. McKenna, M. Teepe, and A. Strasser.** 1998. Overexpression of Bcl-2 does not rescue impaired B lymphopoiesis in IL-7 receptor-deficient mice but can enhance survival of mature B cells. *International Immunology* **10**:1367-1375.
412. **Pardee, A. B.** 1974. Restriction Point for Control of Normal Animal-Cell Proliferation. *Proceedings of the National Academy of Sciences of the United States of America* **71**:1286-1290.
413. **Pardee, A. B.** 1989. G1 Events and Regulation of Cell-Proliferation. *Science* **246**:603-608.
414. **Temin, H. M.** 1971. Stimulation by serum of multiplication of stationary chicken cells. *J.Cell Physiol* **78**:161-170.

415. **Olashaw, N. and W. J. Pledger.** 2002. Paradigms of growth control: relation to Cdk activation. *Sci.STKE*. **2002**: RE7.
416. **Chen, P. L., P. Scully, J. Y. Shew, J. Y. J. Wang, and W. H. Lee.** 1989. Phosphorylation of the Retinoblastoma Gene-Product Is Modulated During the Cell-Cycle and Cellular-Differentiation. *Cell* **58**:1193-1198.
417. **Helin, K., E. Harlow, and A. Fattaey.** 1993. Inhibition of E2F-1 Transactivation by Direct Binding of the Retinoblastoma Protein. *Molecular and Cellular Biology* **13**:6501-6508.
418. **Sellers, W. R., J. W. Rodgers, and W. G. Kaelin, Jr.** 1995. A potent transrepression domain in the retinoblastoma protein induces a cell cycle arrest when bound to E2F sites. *Proc.Natl.Acad.Sci.U.S A* **92**:11544-11548.
419. **Luo, R. X., A. A. Postigo, and D. C. Dean.** 1998. Rb interacts with histone deacetylase to repress transcription. *Cell* **92**:463-473.
420. **Magnaghi-Jaulin, L., R. Groisman, I. Naguibneva, P. Robin, S. Lorain, J. P. Le Villain, F. Troalen, D. Trouche, and A. Harel-Bellan.** 1998. Retinoblastoma protein represses transcription by recruiting a histone deacetylase. *Nature* **391**:601-605.
421. **Geng, Y., E. N. Eaton, M. Picon, J. M. Roberts, A. S. Lundberg, A. Gifford, C. Sardet, and R. A. Weinberg.** 1996. Regulation of cyclin E transcription by E2Fs and retinoblastoma protein. *Oncogene* **12**:1173-1180.
422. **Harbour, J. W. and D. C. Dean.** 2000. The Rb/E2F pathway: expanding roles and emerging paradigms. *Genes & Development* **14**:2393-2409.
423. **Ekholm, S. V., P. Zickert, S. I. Reed, and A. Zetterberg.** 2001. Accumulation of cyclin E is not a prerequisite for passage through the restriction point. *Molecular and Cellular Biology* **21**:3256-3265.
424. **Martinsson, H. S., M. Starborg, F. Erlandsson, and A. Zetterberg.** 2005. Single cell analysis of G(1) check points - The relationship between the restriction point and phosphorylation of pRb. *Experimental Cell Research* **305**:383-391.
425. **Sherr, C. J. and J. M. Roberts.** 2004. Living with or without cyclins and cyclin-dependent kinases. *Genes & Development* **18**:2699-2711.
426. **Zetterberg, A., O. Larsson, and K. G. Wiman.** 1995. What Is the Restriction Point. *Current Opinion in Cell Biology* **7**:835-842.
427. **Cheng, M. G., P. Olivier, J. A. Diehl, M. Fero, M. F. Roussel, J. M. Roberts, and C. J. Sherr.** 1999. The p21(Cip1) and p27(Kip1) CDK 'inhibitors' are essential activators of cyclin D-dependent kinases in murine fibroblasts. *Embo Journal* **18**:1571-1583.
428. **Sherr, C. J. and J. M. Roberts.** 1999. CDK inhibitors: positive and negative regulators of G(1)-phase progression. *Genes & Development* **13**:1501-1512.
429. **Khaled, A. R., D. V. Bulavin, C. Kittipatarin, W. Q. Li, M. Alvarez, K. Kim, H. A. Young, A. J. Fornace, and S. K. Durum.** 2005. Cytokine-driven cell cycling is mediated through Cdc25A. *Journal of Cell Biology* **169**:755-763.
430. **Greider, C., A. Chattopadhyay, C. Parkhurst, and E. Yang.** 2002. BCL-x(L) and BCL2 delay Myc-induced cell cycle entry through elevation of p27 and inhibition of G1 cyclin-dependent kinases. *Oncogene* **21**:7765-7775.
431. **Linette, G. P., Y. Li, K. Roth, and S. J. Korsmeyer.** 1996. Cross talk between cell death and cell cycle progression: BCL-2 regulates NFAT-mediated activation. *Proceedings of the National Academy of Sciences of the United States of America* **93**:9545-9552.
432. **Mazel, S., D. Burtrum, and H. T. Petrie.** 1996. Regulation of cell division cycle progression by bcl-2 expression: A potential mechanism for inhibition of programmed cell death. *Journal of Experimental Medicine* **183**:2219-2226.
433. **Vairo, G., K. M. Innes, and J. M. Adams.** 1996. Bcl-2 has a cell cycle inhibitory function separable from its enhancement of cell survival. *Oncogene* **13**:1511-1519.
434. **Vairo, G., T. J. Soos, T. M. Upton, J. Zalvide, J. A. DeCaprio, M. E. Ewen, A. Koff, and J. M. Adams.** 2000. Bcl-2 retards cell cycle entry through p27(Kip1), pRB relative p130, and altered E2F regulation. *Molecular and Cellular Biology* **20**:4745-4753.
435. **Chattopadhyay, A., C. W. Chiang, and E. Yang.** 2001. BAD/BCL-(xL) heterodimerization leads to bypass of G0/G1 arrest. *Oncogene* **20**:4507-4518.
436. **Cheng, N. L., Y. M. Janumyan, L. Didion, C. Van Hofwegen, E. Yang, and C. M. Knudson.** 2004. Bcl-2 inhibition of T-cell proliferation is related to prolonged T-cell survival. *Oncogene* **23**:3770-3780.
437. **Janumyan, Y. M., C. G. Sansam, A. Chattopadhyay, N. L. Cheng, E. L. Soucie, L. Z. Penn, D. Andrews, C. M. Knudson, and E. Yang.** 2003. Bcl-x(L)/Bcl-2 coordinately regulates apoptosis, cell cycle arrest and cell cycle entry. *Embo Journal* **22**:5459-5470.
438. **Hashimoto, A., K. Hirose, and M. Iino.** 2005. BAD detects coincidence of G2/M phase and growth factor deprivation to regulate apoptosis. *Journal of Pharmacological Sciences* **97**:124P.

Part II: Experimental Data and Results

Part II:

Experimental Data and Results

Chapter 6:

Phosphorylation of Glyoxalase I during growth factor depletion via a GSK-3 β -dependent mechanism

**Ruben Audiffret, Virginie de Hemptinne
and Katia Vancompernelle**

Paper in preparation

Chapter 6: Phosphorylation of glyoxalase I during growth factor depletion via a GSK-3 β -dependent mechanism

INTRODUCTION

The main function of the glyoxalase system, comprising glyoxalase 1 and 2 (GLO1 and 2), is to detoxify reactive and cytotoxic -oxoaldehydes, mainly methylglyoxal (MG) (1-3). After its discovery in 1913, a function of the glyoxalase system in glycolysis was hypothesized. However, this could not be proven (4) and, until today, the full biological function of the glyoxalase system has not been elucidated, but its ubiquitous nature suggests a fundamental and conserved role in life (4,5). In the 1960s and 70s the work of Szent-Gyorgyi indicated a regulatory role for GLO1 and its substrate MG in the control of cell division (for an overview see (1)), but a mechanistic link was never established. However, more and more recent evidence indicates that GLO1 may indeed play a role in the control of cellular growth. For instance, in plants it has been shown that there is a strong correlation between cell proliferation and the expression levels of GLO1, being the highest in strongly proliferating tissues (6). Also in yeast a strong correlation between growth status and GLO1 activity has been described (7). Along with an induction of cell proliferation, the plant growth hormone auxin induces the expression and activity of GLO1 (8). Gibberellin, a growth regulator in plants, induces phosphorylation of GLO1 (9), and calmodulin, a protein involved in cell growth, stimulates GLO1 activity (10).

Methylglyoxal is a cytotoxic by-product of normal metabolism. The main source is the enzymatic and nonenzymatic dephosphorylation from glyceraldehyde-3-phosphate and dihydroxyacetone phosphate, both intermediates of glycolysis (for reviews on MG see (1,3)). MG is very reactive and under physiological conditions MG reacts nonenzymatically with arginine residues mainly (11), but also with lysine and cysteine residues (12). Under normal conditions MG is detoxified by the GLO system to D-lactate using reduced glutathione (GSH) as a cofactor (13). Pathophysiological conditions can cause a rise of MG concentrations and Advanced Glycation Endproducts (AGEs) can be formed (for an overview see (14)). These AGEs are involved in the development of pathophysiological processes *in vivo*. Reports show that MG is able to inhibit growth. This growth inhibition shows a cell type selectivity and is dependent on divers mechanisms (15-22).

We have previously discovered a phosphorylated form of GLO1 and, in a tumor cell death model, we showed that the TNF-induced phosphorylation of GLO1 is required for cell death. Furthermore, phosphorylated GLO1 is not involved in the detoxification of MG, but instead mediates the cytotoxic effects of MG via a pathway that leads to MG-modification of specific target molecules (MG-derived AGE formation). However, the real biological function of phosphorylated GLO1 remains to be determined. (23,24).

Mammalian cells require stimuli from other cells in order to counter the intrinsic apoptosis machinery (25,26). This serves as a mechanism to ensure cells only survive

in the proper context. Cell death is inhibited partly by interfering directly with apoptosis. Anti-apoptotic proteins are up-regulated or activated (27-30) and pro-apoptotic proteins are down-regulated or inactivated (31-38). Another important mechanism by which growth factors may prevent apoptosis is by regulating basic cellular physiology by promoting nutrient uptake and cellular metabolism (30,39-42). A key regulator in apoptotic signaling is glycogen synthase kinase 3 β (GSK-3 β (see reviews (43,44)). Also in growth factor deprivation-induced apoptosis, GSK-3 β has an important regulatory role (45-48).

In this report we show that depletion of IL-3 induces a phosphorylation of GLO1. This phosphorylation is partly mediated by GSK-3 β as inhibition of GSK-3 β also inhibits part of the phosphorylation of GLO1. This phosphorylation doesn't show an effect on the detoxification of MG by GLO1 and it occurs before the onset of apoptosis. Also we show that specific MG derived AGEs are formed, which also can be prevented by inhibition of GSK-3 β .

MATERIALS AND METHODS

Cell Lines and Cultures

The IL-3 dependent pro-B cell line Ba/F3 (49) was purchased from DSMZ. Ba/F3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) with glutamax (Invitrogen) supplemented with heat-inactivated FCS (10% v/v) (Cambrex), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) and 5% conditioned medium from WEHI-3B cells as a source of mouse IL-3. Ba/F3 cells were cultured at 37°C in a humidified incubator under an 5% CO₂ atmosphere. HekT cells, NIH3T3 cells and L929 cells were cultured in DMEM with glutamax (Invitrogen) supplemented with heat-inactivated FCS (10% v/v), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) at 37°C in a humidified incubator under an 8% CO₂ atmosphere. The acute myeloid leukemia cell line HL-60 was a kind gift from J. Van Der Heyden. The cells were maintained in RPMI 1640 with L-glutamine (Invitrogen) supplemented with 10% heat-inactivated FBS and penicillin (100 units/ml) and streptomycin (0.1 mg/ml) and were cultured at 37°C in a humidified incubator under an 5% CO₂ atmosphere.

Cell stimulations

IL-3 was removed by washing exponentially growing cells 3 times in culture medium without conditioned WEHI-3B medium and resuspended in culture medium without conditioned WEHI-3B medium in the absence or presence of indicated concentrations of methylglyoxal (Sigma), N-acetyl-L-cysteine (NAC; Sigma), LiCl (Fluka), SB216763 (Sigma), SB203580 (Sigma) or Z-VAD-FMK (Promega).

HL-60 cells were depleted of FBS by washing exponentially growing cells 3 times in culture medium without FBS and resuspended in FBS-free culture medium at a concentration of 1×10^6 cells/ml. NIH3T3 cells were depleted by washing them 3 times with culture medium without FBS and subsequently cultured in culture medium containing 0.5% (v/v) FBS.

DNA distribution analysis

Cells were stained with propidium iodide (PI, Sigma) containing staining solution (50) and analyzed by a FACSCalibur flow cytometer (488_{Ex}/590_{Em}). Cell Quest software was used to analyze cell cycle distribution (FACS Calibur, Becton Dickinson) (50,51). Ten thousand cells were routinely analyzed.

Electrophoresis and immunoblotting

The cells were washed 3 times with ice-cold PBS buffer and lysed with cytosol extraction buffer (10 mM Tris-HCL pH7.4, 50 mM EDTA pH8.0, 25 mM NaCl, 0,7% TritonX-100, 100 mM PMSF, 1 tablet Complete Protease Inhibitor Cocktail (Roche)/50 ml cytosol extraction buffer). Cell lysates were cleared by centrifugation (14,000 x g). Protein concentrations were determined using Bradford method. Proteins were separated by SDS-PAGE and transferred onto a PVDF membrane (Hybond-P, Amersham Pharmacia Biotech). The blots were incubated with the desired antibodies followed by ECL-based detection (Amersham Pharmacia Biotech).

2-Dimensional Gel Electrophoresis

Isoelectric focusing was carried out on 18 cm IPG strips, pH 4-7 (GE Healthcare) according to the manufacturer's instructions. For the second dimension, proteins were separated by SDS-polyacrylamide gel electrophoresis (PAGE, 12%).

Assay of glyoxalase1 activity

GLO1 activity was determined by using a spectrophotometric method, which monitors the initial rate of change in absorbance at 240 nm caused by the formation of S-D-lactoylglutathione (52). The standard assay mixture contained 2-mM MG and 2-mM GSH in a sodium phosphate buffer (50 mM, pH 6.6, 20°C). The reaction mixture was allowed to stand for 10 min at 37°C before the addition of the cytosolic protein fraction to ensure the equilibration of hemithioacetal formation.




Figure 6.1. Time course of GLO1 phosphorylation in IL-3 depleted Ba/F3 cells.

Ba/F3 cells were depleted of IL-3 and lysed after indicated periods of time. Cells were lysed in CEB and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with a polyclonal anti-human GLO1 antibody (6) are shown. A reference spot α , which corresponds to the non-NO-responsive form of GLO1 is indicated (6).

Measurement of D-Lactate

A fluorimetric method was used as previously described (53). Briefly, D-Lactate formation is measured by the formation of NADH during oxidation of D-Lactate by a D-lactate dehydrogenase.


RESULTS

Glyoxalase I is phosphorylated upon IL-3 starvation.


We recently described that most cell types, analysed so far express multiple isoforms of glyoxalase1 (GLO1), including a NO-responsive form and a non-NO-responsive form, which we designated as the α -isoform (23). These isoforms can be separated by 2-dimensional gel electrophoresis (2-D) and detected by Western blotting using an anti-GLO1 antibody. We previously discovered phosphorylated GLO1 by means of cell labeling with 32 P-orthophosphate (24). Multiple phosphorylation of GLO1 can occur either on the NO-responsive form or the α -isoform and this can be monitored in 2-D gels by a shift of the isoform to the acidic side of the gel (23).

To investigate whether phosphorylation of GLO1 was related to growth factor signaling in mammalian systems, Ba/F3 cells, which are dependent on IL-3 for survival and proliferation, were depleted of IL-3 and cell lysates were prepared at the indicated time points cells. The samples were analysed by high-resolution 2-D in combination with

A.



B.



C.




Figure 6.2. Phosphorylation of GLO1 in different cell lines.

A. Phosphorylation of GLO1 in NIH3T3 cells. NIH3T3 cells were washed in culture medium without FBS and cultured in the presence of 0.5 % FBS. Cells were lysed in CEB after 1 hour and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 400 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Cells were lysed in CEB and after ethanol precipitation resuspended in 2-DE lysis buffer. Representative western blots of 2-DE gels with a polyclonal anti-human GLO1 antibody are shown. **B.** Phosphorylation of GLO1 in HL-60 cells. HL-60 cells were depleted of FBS and lysed after 1 day. Cells were lysed in CEB and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Cells were lysed in CEB and after ethanol precipitation resuspended in 2-DE lysis buffer. Representative western blots of 2-DE gels with a polyclonal anti-human GLO1 antibody are shown. **C.** Serum starvation-induced apoptosis in HL-60 cells. HL-60 cells were depleted of FBS. Apoptosis was measured by the analysis of the percentage of cells with sub-diploid DNA at the indicated time points.

Western blotting using a polyclonal antibody raised against GLO1 (Figure 6.1). As shown in the upper panel (0h), the isoform pattern of GLO1 in Ba/F3 cells is identical to the one we described previously for several other mouse and human cell types (23). The most basic isoforms corresponds to the NO-responsive form of GLO1 and the non NO-modified form of GLO1 is indicated by α . After one hour of IL-3 starvation, the most basic isoform has shifted to the acidic side, which is indicative for phosphorylation (spot indicated by an arrow). After three hours of IL-3 starvation, the relative amount of the NO-responsive form of GLO1 is reduced and had become partially dephosphorylated (shifted more to the basic side as compared to the 1h time point). After five hours IL-3 starvation, the NO-responsive form of GLO1 is completely disappeared and phosphorylation of GLO1 is induced on the non-No-responsive form (phosphorylated spots are indicated by an arrow). Note that the total amount of phosphorylated GLO1 at 5h of IL-3 starvation is relatively high as compared to the previous time points of IL-3 starvation. In several independent experiments, we also noticed that the time course and degree of phosphorylation of GLO1 induced by depletion of IL-3 could vary according to the experiment and could be induced either earlier or later as shown in Figure 6.1.

Growth Factor withdrawal induced phosphorylation of GLO1 is not exclusive for IL-3

To examine whether phosphorylation of GLO1 was specific for IL-3 depletion or whether it was a more general phenomenon, the phosphorylation of GLO1 upon GF depletion was also examined in two other cell lines.

Mouse fibroblasts are commonly used in the study of serum dependence. After serum withdrawal mouse fibroblasts reversibly exit the cell cycle (54-57) and after a sustained absence of GFs they undergo apoptosis (58,59). The mouse fibroblast cell line NIH3T3 was serum-starved as described in M&M. As shown in Figure 6.2A immunoblots of lysates of cells that were maintained in culture medium with 0.5 % FBS display GLO1 is phosphorylated on the non NO-modified form of GLO1. Immunoblots of lysates from cells that were kept in culture medium with 10 % FBS show that this spot was absent. This phosphorylation occurs after 1 hour of starvation, well before the onset of apoptosis.

Another cell line tested was the human leukemia cell line HL-60. HL-60 is a hematopoietic cell line, like Ba/F3 cells, albeit of the myeloid lineage, whereas Ba/F3 cells are lymphoid cells. It has been reported that HL-60 cells are sensitive to exogenous MG, making HL-60 very interesting to study possible effects on the GLO system. Addition of MG will cause G₁ growth arrest and induce apoptosis in HL-60 cells (19). Upon serum depletion HL-60 also arrest growth and induce apoptosis (60,61). Analysis of immunoblots detected with an anti-GLO1 antibody shows the induction of phosphorylation of GLO1 upon serum starvation (Figure 6.2.B). When compared to the control situation the spot second to the right disappeared and a more acidic new spot appears. Although lysates were made after




Figure 6.3. Inhibition of GSK-3 interferes with phosphorylation of GLO1.

A. Effect of GSK-3 inhibition on IL-3 depletion-induced apoptosis. Ba/F3 cells were depleted of IL-3 in the presence or absence of the indicated inhibitor. Apoptosis was measured by the analysis of the percentage of cells with sub-diploid DNA at the indicated time points. **B and C.** GLO1 phosphorylation in IL-3 depleted Ba/F3 cells in the presence of GSK-3 inhibitors. Ba/F3 cells were depleted of IL-3 and cultured in the presence or absence of LiCl (**A**) or SB216763 (**B**). Cells were lysed in CEB after 3 hours and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels with a polyclonal anti-human GLO1 antibody are shown. Concentrations used are LiCl: 20 mM and SB216763: 20 μ M.




Figure 6.4. Phosphorylation of GLO1 does not affect the formation of D-lactate.

A. A time course of D-lactate formation in Ba/F3 cells. Ba/F3 cells were washed in culture medium without IL-3 and subsequently cultured in the presence or absence of IL-3. At indicated time points the presence of D-lactate in the medium was measured by detection of the formation of NADH upon incubation with D-lactate dehydrogenase. **B and C.** D-lactate production in IL-3 depleted Ba/F3 cells in the presence of GSK-3 inhibitors. Ba/F3 cells were depleted of IL-3 and cultured in the presence or absence of LiCl (**B**) or SB216763 (**C**). The presence of D-lactate in the medium was measured after indicated time intervals. Concentrations used are LiCl: 20 mM and SB216763: 20 μ M.





Figure 6.5. *In vitro* GLO1 activity is not altered by phosphorylation.

A. A time course of GLO1 activity in IL-3 starved Ba/F3 cells. Ba/F3 cells were washed in culture medium without IL-3 and subsequently cultured in the presence or absence of IL-3. At indicated time points cells were lysed in CEB. Lysates were brought to equal concentrations and cell lysates were used to assay GLO1 activity. **B.** GLO1 activity in IL-3 depleted Ba/F3 cells in the presence of GSK-3 inhibitors. Ba/F3 cells were washed in culture medium without IL-3 and subsequently cultured in the presence or absence of IL-3 and LiCl was added. At indicated time points cells were lysed in CEB. Lysates were brought to equal concentrations and cell lysates were used to assay GLO1 activity. Reaction rates were determined using Ultraspec 1100 software (Amersham, BD Biosciences) and are shown relative to the reaction rate of lysates made of exponentially growing cells, which are set as 100%. LiCl was added in a concentration of 20 mM.

24 hours of serum starvation, phosphorylation is still induced before the onset of apoptosis, which is only observed after 72 hours of serum starvation (Figure 6.2.C).

In conclusion, the data presented here show that the phosphorylation of GLO1 upon GF withdrawal is not restricted to Ba/F3 cells, but can also occur in mouse fibroblasts and

A.



B.






Figure 6.6. IL-3 depletion induces the formation of specific MG-derived AGEs in Ba/F3 cells.

A. Formation of specific MG-AGEs. Ba/F3 cells were depleted of IL-3 and lysed in CEB after indicated periods of time. Equal amounts of protein were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Formation of specific MG-AGEs was analysed using the mAb6B antibody (Biologo). As a loading control, blots were subsequently stripped and reprobed with a β -actin antibody (Abcam). **B.** Inhibition of MG-AGE formation. Ba/F3 cells were depleted of IL-3, cultured in the presence of LiCl or Z-VAD-FMK and lysed in CEB after indicated periods of time. Equal amounts of protein were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Formation of specific MG-AGEs was analysed using the mAb6B antibody. As a loading control, blots were subsequently stripped and reprobed with a β -actin antibody.

A.



B.



C.




Figure 6.7. MG is synergistic with IL-3 depletion-induced apoptosis.

A. Cell death with exogenous MG. Ba/F3 cells were washed in culture medium without IL-3 and subsequently cultured in the presence or absence of IL-3 and MG was added. After 24 hours apoptosis was measured by the analysis of the percentage of cells with sub-diploid DNA. **B. and C.** Inhibition of cell death. Ba/F3 cells were depleted of IL-3 and cultured in the presence of MG and LiCl (**B**) or NAC (**C**). After 24 hours apoptosis was measured by the analysis of the percentage of cells with sub-diploid DNA. Concentrations used are LiCl: 20 mM, NAC: 20 mM, MG: 500 μ M.

human leukemic cells upon GF withdrawal. It may therefore represent a more general mechanism. This hypothesis is strengthened by the observation that GF withdrawal-induced phosphorylation takes place in several species, mouse and human, and is not confined to one cell type.

GSK-3 is involved in IL-3 starvation-induced post-translational modification of GLO1.

GSK-3 has been shown to have a central role in the regulation of cell cycle and apoptosis (43,44). Also, upon growth factor withdrawal, GSK-3 is involved in the induction of G₁ arrest by regulating the levels of cyclin D1 (62,63), and in the induction of apoptosis and inhibition of GSK-3 activity also ameliorates the growth factor-withdrawal-induced cell death (45-48,64). To investigate whether GSK-3 was involved in the phosphorylation of GLO1 upon IL-3 depletion of Ba/F3 cells, we used LiCl, a well known inhibitor of GSK-3 (65). Because it has been previously shown that activation of GSK-3 β is required for the induction of apoptosis by GF-depletion, we first tested in our experimental setting whether inhibition of GSK-3 β by LiCl also inhibited apoptosis in Ba/F3 induced by depletion of IL-3. Ba/F3 cells were cultured with or without conditioned WEHI-3B medium as a source of IL-3, and in the presence or absence of LiCl. As expected, LiCl inhibited apoptosis by approximately 80% induced by IL-3 depletion (Figure 6.3.A). Next, we examined whether LiCl had an effect on the phosphorylation of GLO1 induced by IL-3 depletion from Ba/F3 cells. As shown in Figure 6.3.B, the phosphorylation of GLO1 (indicated by an arrow) was completely prevented in the presence of LiCl. In addition, the phosphorylation of GLO1 induced by depletion of IL-3 was also completely prevented in the presence of a more specific inhibitor of GSK-3 β namely SB216763 (Figure 6.3.C). SB216763 also inhibited apoptosis induced by IL-3 starvation to the same extent as LiCl (Figure 6.3.A). These results indicate that GSK-3 β is involved in the pathway that leads to phosphorylation of GLO1 induced by IL-3 starvation. It is however unlikely that GLO1 is a direct substrate for GSK-3 β , because coexpression of the catalytic subunit of GSK-3 β with GLO1 in serum-starved HEK293 cells did not result in phosphorylation of GLO1 as studied by means of labelling cells with ³²P-orthophosphate. Nor could GSK-3 β phosphorylate GLO1 in vitro (see chapter 7). By contrast, coexpression of the catalytic subunit of CaMKII with GLO1 in HEK293 did result in phosphorylation of GLO1 (de Hemptinne and Vancompernelle, unpublished results) and GLO1 is also an in vitro substrate for CaMKII (see chapter 7).

Methylglyoxal detoxification is not impaired after IL-3 depletion

Given the cytostatic and cytotoxic properties of MG, a likely hypothesis could be that upon IL-3 depletion, the MG-detoxification capacity of the GLO system could be inhibited by for instance phosphorylation of GLO1. This would cause a rise in the intracellular MG concentration, a condition that could induce growth arrest and if the absence of IL-3 persists, ultimately apoptosis. To investigate whether the MG-detoxification capacity of the glyoxalase system was impaired upon IL-3 depletion, we measured GLO1 activities as well as D-lactate concentrations, the end product of the glyoxalase system. The formation of D-lactate was measured by a fluorimetric enzyme assay as described in (53). The

concentration of D-lactate was determined by the formation of NADH during oxidation of D-Lactate by D-lactate dehydrogenase. As shown in Figure 6.4.A, there is no significant difference in formation of D-lactate between IL-3 starved cells and control cells. GLO1 activity in cell lysates derived from Ba/F3 cells in the presence or absence of IL-3 was determined by a spectrophotometric method, which monitors the initial rate of the GLO1-catalyzed production of S-D-lactoylglutathione from the hemithioacetal that is formed spontaneously by incubating MG in glutathione (see M&M). As shown in Figure 6.5.A, in several separate experiments there was no significant change in GLO1 activity in cell lysates from IL-3-depleted cells compare to control cells. These findings indicate that the MG-detoxification activity of the glyoxalase system is not altered upon GF withdrawal. These data also suggest that phosphorylation of GLO1, which is induced upon GF withdrawal, is not accompanied by changes in the GLO1 activity, which is in agreement with our previous findings in L929 cells (24). This finding is further strengthened by the observation that inhibition of GLO1 phosphorylation with LiCl or SB216763 upon IL-3 depletion, did not result in a change of the D-lactate production or the *in vitro* GLO1 activity (Figure 6.4.B and 6.4.C, and Figure 6.5.B respectively).

Specific MG-derived AGEs are formed during IL-3 withdrawal.

MG has been reported to be involved in several signaling pathways (24,66-69). Recent findings indicate MG has a distinct role as a modulator of cell death (24,70-72). Previous results indicated that phosphorylated GLO1 was involved, directly or indirectly, in a pathway that leads to MG-derived AGE modification of specific proteins. Therefore, we tested whether MG-derived AGE modification was also induced upon GF withdrawal. For this purpose, we used a monoclonal antibody (mAb6B) raised against *in vitro* MG-modified keyhole limpet hemocyanin. This antibody recognizes MG-modified proteins in arterial walls of diabetic kidneys (73) and we have previously shown that this antibody recognizes specific MG-derived AGE epitopes induced during TNF-induced cell death in L929 cells (24). A Western blot of cell lysates derived from IL-3 depleted Ba/F3 cells developed with this antibody is shown in Figure 6.6.A. MG-derived AGE modification is induced on a distinct protein band, with an apparent M.W. 16 kDa (indicated by an arrow), after 22h of IL-3 depletion. The MG-derived AGE formation was already observed after eight hours of IL-3 starvation (data not shown).

Next, we investigated whether phosphorylated GLO1 was involved in the induction of MG-derived AGE-modification of the 16 kDa protein. As already shown in Figure 6.3, inhibition of GSK-3 by LiCl efficiently inhibited the phosphorylation of GLO1, therefore we used LiCl to test whether inhibition of GSK-3 also influenced the MG-derived AGE-formation. As shown in Figure 6.6.B, MG-derived AGE-modification of the 16 kDa protein (indicated by an arrow) was significantly reduced in the presence of LiCl.

Since GF-withdrawal also induces caspase activation (74), we also tested the




Figure 6.8. Activation of p38 by exogenous MG.

A. Activation of p38. Ba/F3 cells were washed in culture medium without IL-3 and subsequently cultured in the presence or absence of IL-3 and MG was added. Cells were lysed in CEB after indicated periods of time. Equal amounts of protein were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Activation of p38 was determined using an antibody directed against phosphor-p38 (Cell Signaling). Subsequently blots were stripped and reprobed with an anti-p38 antibody (Cell Signaling). **B.** Inhibition of p38 activation. Ba/F3 cells were depleted of IL-3 and cultured in the presence of MG and LiCl or NAC. Cells were lysed in CEB after indicated periods of time. Equal amounts of protein were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Activation of p38 was determined using an antibody directed against phosphor-p38. Subsequently blots were stripped and reprobed with an anti-p38 antibody. **C.** MG-dependent cell death. Ba/F3 cells were depleted of IL-3 and MG and SB203580 were added. After 24 hours apoptosis was measured by the analysis of the percentage of cells with sub-diploid DNA. Concentrations used are LiCl: 20 mM, NAC: 20 mM, MG: 500 μ M and SB203580: 10 μ M. Protein band intensities were measured using Genetools software (Westburg) and phosphor-p38 to p38 ratios were determined.

possibility whether the 16 kDa MG-AGE-modified protein was formed by caspase-cleavage of a larger pre-formed MG-AGE-modified protein. As shown in Figure 6.6.B, the 16 kDa MG-AGE-modified protein was still formed in the presence of the pan-caspase inhibitor Z-VAD-FMK.

Exogenous MG reinforces the induction of apoptosis after IL-3 withdrawal

To further examine whether MG played a role in the induction of apoptosis in Ba/F3 cells induced by IL-3 depletion, we tested the effect of exogenously added MG. The percentage of apoptotic cells was measured by flow cytometry using the percentage of cells that had a hypoploid DNA profile (expressed as sub-G₁). As shown in Figure 6.7.A, in the presence of 0.5 mM exogenously added MG and in the absence of IL-3, the percentage of apoptotic cells was increased by approximately 55 % compared to the percentage of apoptotic cells induced by IL-3 depletion alone. As a control, addition of 0.5 mM MG did not induce apoptosis in a Ba/F3 cell culture in the presence of IL-3. In the presence of IL-3, exogenous addition of 0.5 mM MG could also not induce formation of the specific 16 kDa MG-AGE-modified protein (data not shown).

We next sought to determine whether GSK-3 β was involved in this synergistic effect induced by exogenous MG. As shown in Figure 6.7.B, the synergistic effect by exogenous MG (0.5 mM) on the induction of apoptosis induced by IL-3 depletion was completely abolished in the presence of LiCl. Moreover, addition of LiCl also prevented the synergistic effect on the formation of the specific 16 kDa MG-AGE-modified protein (data not shown). These data suggest that GSK-3 β plays a crucial role in this process and that the role of MG and presumably the specific MG-derived AGE-modified proteins is downstream of GSK-3 β .

Effects of MG on cells are often dependent on the formation of reactive oxygen species (ROS; (15,16,21,75-78)). Even though the reaction of MG with amino acids by itself leads to superoxide formation (79-81), MG also hampers the activities of the antioxidant mechanisms in cells (21,82-84) and directly modifies components of respiratory complex III in mitochondria, the major site of superoxide production (85,86). In order to investigate the role of ROS in the induction of apoptosis, the effect of N-acetyl-cysteine (NAC) was examined on IL-3 starvation and MG-induced apoptosis. Co-incubation of Ba/F3 cells with NAC strongly impedes the outcome of addition of MG on apoptosis (Figure 6.7.C).

MG-induced effects are not mediated by the p38 MAPK pathway

Several reports implicate the p38 and JNK MAPK pathways in MG-induced signaling to cell death. MG induces a rise in intracellular ROS, hereby leading to the activation of apoptosis signal-regulating kinase 1 (ASK1), and subsequently the p38 α MAPK pathway (20). When Ba/F3 cells are deprived of IL-3 in the presence of MG, a strong and rapid

phosphorylation of p38 is induced (Figure 6.8.A). A time-course experiment shows that p38 is phosphorylated within 0.5 hour and peaks after 2 hours. Together with the induction of phosphorylation, an increase in the total amount p38 is also observed. But when the ratios of phosphorylated p38 to total p38 are calculated it is clear that the rise in activated p38 is mainly caused by phosphorylation. However, also in the presence of IL-3 addition of MG leads to a strong activation of p38 (Figure 6.9.A), although MG is not cytotoxic at the concentration used (Figure 6.7.A). This suggest that exogenous MG is able to activate p38 MAPK independent of when IL-3 is present or not.

Next, we tested whether the GSK-3 β inhibitor LiCl could prevent the phosphorylation of p38 induced by MG. As shown in Figure 6.8.B, this was not the case although LiCl strongly abrogates the induction of apoptosis by MG. In contrast, NAC could prevent the MG-induced phosphorylation of p38 and also MG-induced cell death (Figure 6.7.C and Figure 6.8.B). Next, we used a specific p38 MAPK inhibitor SB203580, to examine whether p38 played a role in the synergistic effect of MG on the induction of apoptosis. Pre-incubation with SB203508 had no significant effect on MG-induced cell death (Figure 6.8.C). Of note is the observation that NAC by itself is able to induce p38 phosphorylation to a certain degree (Figure 6.8.B). This might be a consequence of the observed pro-oxidant activity of NAC under certain conditions (87). In summary, these data suggest that the synergistic effect of MG on the induction of apoptosis induced by IL-3 depletion occurs independent of p38 MAPK, but is dependent on GSK-3 β .

DISCUSSION

MG is a by-product of glycolysis, the most fundamental metabolic pathway. It therefore occurs in every cell and despite the presence of several detoxifying mechanisms Mg can reach high intracellular levels (88). Approximately 10% of total protein content has been estimated to be modified by MG under physiological conditions. Previous reports from our group have implicated MG in TNF-induced necrosis (24). GLO1 is phosphorylated in a PKA-dependent manner and a concomitant rise in MG is observed. As a consequence of these events specific MG-derived AGEs are formed. Inhibition of these events efficiently provides protection against TNF-induced necrosis in L929 cells.

In this study we have demonstrated that depletion of IL-3 induces phosphorylation of GLO1. A possible kinase to be involved in this process would be GSK-3 β . GSK-3 β is a ubiquitously expressed serine/threonine kinase and is a key regulator in diverse biological processes, including regulation of metabolism and apoptosis (reviewed in (43,44)). A remarkable characteristic of this kinase is the fact that it GFs lead to its inactivation by phosphorylation, and that inactivation of GSK-3 β is correlated with survival whereas activation correlates with apoptosis (89,90). Inhibition of GSK-3 β with LiCl or the specific inhibitor SB216763 also inhibits phosphorylation of GLO1 (Figure 6.3), showing the

involvement of GSK-3 β . The phosphorylation of GLO1 does not seem to influence the activity of GLO1 as measured *in vitro* nor is there a change in detoxification of MG as determined by the production of D-lactate (Figure 6.4 and 5). In spite of the unaltered capacities of GLO1 to detoxify MG, specific MG-AGEs are formed. We hereby provide a possible physiological function for the ubiquitous GLO system and its primary substrate MG. Mammalian cells are dependent on GFs for growth and survival (91). An important mechanism used by GFs is the maintenance of cellular metabolism and glycolysis in particular (41,42,92,93). The intracellular concentration of MG is determined by the rate of glycolysis, and could therefore function as a signal molecule to elicit a response when drastic changes occur in the rate of glycolysis, since this would be reflected in intracellular MG concentration. IL-3 depletion would indeed induce a rapid decline in glycolysis (42,94).

Since *in vivo* induced MG-AGEs are heterogeneous, the choice of the antibody was based on the proven efficacy of the antibody to detect argpyrimidine in immunohistochemical samples of diabetic patients (73). The specific nature of the MG-derived AGEs was substantiated by the observation that addition of exogenous MG was insufficient to induce their formation (Figure 6.6.C). Only after IL-3 depletion MG-AGEs were detected, but this could be inhibited by inhibition of GSK-3 β and the consequent phosphorylation of GLO1. This strongly supports the hypothesis that both phosphorylated GLO1 and a rise in MG levels are required for the formation of specific MG-derived protein modification.

MG is known to be a strong inducer of ROS and this ROS formation is often implied in the detrimental activity of MG (15,16,21,77,85). We also found the synergy of MG with IL-3 depletion could be abrogated by addition of NAC, an anti-oxidant which has been shown to be able to quench MG-induced ROS production (76). The concentrations of MG used in this report are within the range commonly used when the biological effects of MG are studied (19,21,75,85,95,96). Moreover, MG levels do not exceed the levels found in living cells (88), and are not cytotoxic by themselves (Figure 6.7.A). Several studies show the p38 and JNK MAPK pathways are activated upon MG-induced ROS formation and mediate the subsequent signaling events (15,16,20,75,95). However, this doesn't seem to be the case in our system. Although a strong activation of p38 was induced by MG, this was not sufficient to induce cell death (Figure 6.8). Nor could it be inhibited by LiCl, even though this strongly inhibited cell death and MG modification of protein. This observation is even strengthened by the inability of the specific p38 inhibitor SB203580 to inhibit the synergistic effect seen between IL-3 starvation and exogenous MG. Phosphorylation of JNK was also observed, but to a much lesser extent than p38 activation (data not shown). Further examination of the specific MG-modified protein or the possible pathways activated by ROS are needed to clarify the action of MG and phosphorylated GLO1 in IL-3 depletion-induced apoptosis.

In conclusion this report shows that upon IL-3 depletion in Ba/F3 cells GLO1 is phosphorylated in a GSK-3 β dependent manner. Subsequently specific MG-modification

of proteins occurs. The formation of these MG-modified proteins is not the consequence of a simple rise in intracellular MG concentrations, since neither GLO1 activity or MG detoxification is impaired, nor could they be induced by addition of exogenous MG. Our results could provide a system that allows cells to integrate external signals from GFs with metabolism.

Reference List

1. **Kalapos, M. P.** 1999. Methylglyoxal in living organisms - Chemistry, biochemistry, toxicology and biological implications. *Toxicology Letters* **110**:145-175.
2. **Thornalley, P. J.** 1990. The Glyoxalase System - New Developments Towards Functional-Characterization of A Metabolic Pathway Fundamental to Biological Life. *Biochemical Journal* **269**:1-11.
3. **Thornalley, P. J.** 1996. Pharmacology of methylglyoxal: Formation, modification of proteins and nucleic acids, and enzymatic detoxification - A role in pathogenesis and antiproliferative chemotherapy. *General Pharmacology* **27**:565-573.
4. **RACKER, E.** 1951. The mechanism of action of glyoxalase. *J.Biol.Chem.* **190**:685-696.
5. **Hopkins, F. G. and E. J. Morgan.** 1945. On the distribution of glyoxalase and glutathione. *Biochem.J.* **39**:320-324.
6. **Deswal, R., T. N. Chakaravarty, and S. K. Sopory.** 1993. The Glyoxalase System in Higher-Plants - Regulation in Growth and Differentiation. *Biochemical Society Transactions* **21**:527-530.
7. **Dudani, A. K., L. K. Srivastava, and R. Prasad.** 1984. Glyoxalase-I Activity and Cell-Cycle Regulation in Yeast. *Biochemical and Biophysical Research Communications* **119**:962-967.
8. **Paulus, C., B. Kollner, and H. J. Jacobsen.** 1993. Physiological and Biochemical-Characterization of Glyoxalase-I, A General Marker for Cell-Proliferation, from A Soybean Cell-Suspension. *Planta* **189**:561-566.
9. **Khan, M. M. K., A. Jan, H. Karibe, and S. Komatsu.** 2005. Identification of phosphoproteins regulated by gibberellin in rice leaf sheath. *Plant Molecular Biology* **58**:27-40.
10. **Deswal, R. and S. K. Sopory.** 1999. Glyoxalase I from *Brassica juncea* is a calmodulin stimulated protein. *Biochimica et Biophysica Acta-Molecular Cell Research* **1450**:460-467.
11. **Thornalley, P. J., S. Battah, N. Ahmed, N. Karachalias, S. Agalou, R. Babaei-Jadidi, and A. Dawney.** 2003. Quantitative screening of advanced glycation endproducts in cellular and extracellular proteins by tandem mass spectrometry. *Biochemical Journal* **375**:581-592.
12. **Ravichandran, V., T. Seres, T. Moriguchi, J. A. Thomas, and R. B. Johnston.** 1994. S-Thiolation of Glyceraldehyde-3-Phosphate Dehydrogenase Induced by the Phagocytosis-Associated Respiratory Burst in Blood Monocytes. *Journal of Biological Chemistry* **269**:25010-25015.
13. **Jowett, M. and J. H. Quastel.** 1933. The glyoxalase activity of the red blood cell: The function of glutathione. *Biochem.J.* **27**:486-498.
14. **Singh, R., A. Barden, T. Mori, and L. Beilin.** 2001. Advanced glycation end-products: a review. *Diabetologia* **44**:129-146.
15. **Du, J., H. Suzuki, F. Nagase, A. A. Akhand, T. Yokoyama, T. Miyata, K. Kurokawa, and I. Nakashima.** 2000. Methylglyoxal induces apoptosis in Jurkat leukemia T cells by activating c-jun N-terminal kinase. *Journal of Cellular Biochemistry* **77**:333-344.
16. **Du, J., H. Suzuki, F. Nagase, A. A. Akhand, X. Y. Ma, T. Yokoyama, T. Miyata, and I. M. Nakashima.** 2001. Superoxide-mediated early oxidation and activation of ASK1 are important for initiating methylglyoxal-induced apoptosis process. *Free Radical Biology and Medicine* **31**:469-478.
17. **Du, J., S. H. Cai, H. Suzuki, A. A. Akhand, X. Y. Ma, Y. Takagi, T. Miyata, L. Nakashima, and F. Nagase.** 2003. Involvement of MEKK1/ERK/P21(Waf1/Cip1) signal transduction pathway in inhibition of IGF-I-mediated cell growth response by methylglyoxal. *Journal of Cellular Biochemistry* **88**:1235-1246.
18. **Halder, J., M. Ray, and S. Ray.** 1993. Inhibition of Glycolysis and Mitochondrial Respiration of Ehrlich Ascites-Carcinoma Cells by Methylglyoxal. *International Journal of Cancer* **54**:443-449.
19. **Kang, Y. B., L. G. Edwards, and P. J. Thornalley.** 1996. Effect of methylglyoxal on human leukaemia 60 cell growth: Modification of DNA, G(1) growth arrest and induction of apoptosis. *Leukemia Research* **20**:397-405.
20. **Liu, B. F., S. Miyata, Y. Hirota, S. Higo, H. Miyazaki, M. Fukunaga, Y. Hamada, S. Ueyama, O. Muramoto, A. Uriuhara, and M. Kasuga.** 2003. Methylglyoxal induces apoptosis through activation of p38 mitogen-activated protein kinase in rat mesangial cells. *Kidney International* **63**:947-957.
21. **Okado, A., Y. Kawasaki, Y. Hasuike, M. Takahashi, T. Teshima, J. Fujii, and N. Taniguchi.** 1996. Induction of apoptotic cell death by methylglyoxal and 3-deoxyglucosone in macrophage-derived cell lines. *Biochemical and Biophysical Research Communications* **225**:219-224.
22. **Ray, M., J. Halder, S. K. Dutta, and S. Ray.** 1991. Inhibition of Respiration of Tumor-Cells by Methylglyoxal and Protection of Inhibition by Lactaldehyde. *International Journal of Cancer* **47**:603-609.
23. **de, H., V. D. Rondas, J. Vandekerckhove, and K. Vancompernelle.** 2007. Tumour necrosis factor induces phosphorylation primarily of the nitric-oxide-responsive form of glyoxalase I. *Biochem.J.* **407**:121-128.
24. **Van Herreweghe, F., J. Q. Mao, F. W. R. Chaplen, J. Grooten, K. Gevaert, J. Vandekerckhove, and K. Vancompernelle.** 2002. Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methyl glyoxal-derived AGE. *Proceedings of the National Academy of Sciences of the United States of America* **99**:949-954.
25. **Raff, M. C.** 1992. Social Controls on Cell-Survival and Cell-Death. *Nature* **356**:397-400.
26. **Raff, M. C., B. A. Barres, J. F. Burne, H. S. R. Coles, Y. Ishizaki, and M. D. Jacobson.** 1994. Programmed Cell-Death and the Control of Cell-Survival. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **345**:265-268.
27. **Dumon, S., S. C. R. Santos, F. Debierre-Grockiego, V. Gouilleux-Gruart, L. Cocault, C. Boucheron, P. Mollat, S. Gisselbrecht, and F. Gouilleux.** 1999. IL-3 dependent regulation of Bcl-x(L) gene expression by STAT5 in a bone marrow derived cell line. *Oncogene* **18**:4191-4199.
28. **Leverrier, Y., J. Thomas, G. R. Perkins, M. Mangeney, M. K. L. Collins, and J. Marvel.** 1997. In bone marrow

- derived Baf-3 cells, inhibition of apoptosis by IL-3 is mediated by two independent pathways. *Oncogene* **14**:425-430.
29. **Leverrier, Y., J. Thomas, A. L. Mathieu, W. Low, B. Blanquier, and J. Marvel.** 1999. Role of PI3-kinase in Bcl-X induction and apoptosis inhibition mediated by IL-3 or IGF-1 in Baf-3 cells. *Cell Death and Differentiation* **6**:290-296.
30. **Plas, D. R., S. Talapatra, A. L. Edinger, J. C. Rathmell, and C. B. Thompson.** 2001. Akt and Bcl-x(L) promote growth factor-independent survival through distinct effects on mitochondrial physiology. *Journal of Biological Chemistry* **276**:12041-12048.
31. **Allan, L. A., N. Morrice, S. Brady, G. Magee, S. Pathak, and P. R. Clarke.** 2003. Inhibition of caspase-9 through phosphorylation at Thr 125 by ERK MAPK. *Nature Cell Biology* **5**:647-U45.
32. **delPeso, L., M. GonzalezGarcia, C. Page, R. Herrera, and G. Nunez.** 1997. Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt. *Science* **278**:687-689.
33. **Ekert, P. G., A. M. Jabbour, A. Manoharan, J. E. Heraud, J. Yu, M. Pakusch, E. M. Michalak, P. N. Kelly, B. Callus, T. Kiefer, A. Verhagen, J. Silke, A. Strasser, C. Borner, and D. L. Vaux.** 2006. Cell death provoked by loss of interleukin-3 signaling is independent of Bad, Bim, and PI3 kinase, but depends in part on Puma. *Blood* **108**:1461-1468.
34. **Ekoff, M., T. Kaufmann, M. Engstrom, N. Motoyama, A. Villunger, J. I. Jonsson, A. Strasser, and G. Nilsson.** 2007. The BH3-only protein Puma plays an essential role in cytokine deprivation-induced apoptosis of mast cells. *Blood*.
35. **Harada, H., B. Quearry, A. Ruiz-Vela, and S. J. Korsmeyer.** 2004. Survival factor-induced extracellular signal-regulated kinase phosphorylates BIM, inhibiting its association with BAX and proapoptotic activity. *Proceedings of the National Academy of Sciences of the United States of America* **101**:15313-15317.
36. **Low, W., G. Olmos-Centenera, C. Madsen, Y. Leverrier, and M. Collins.** 2001. Role of Bax in apoptosis of IL-3-dependent cells. *Oncogene* **20**:4476-4483.
37. **Tan, Y., M. R. Demeter, H. Ruan, and M. J. Comb.** 2000. BAD Ser-155 phosphorylation regulates BAD/Bcl-XL interaction and cell survival. *Journal of Biological Chemistry* **275**:25865-25869.
38. **Zhou, X. M., Y. M. Liu, G. Payne, R. J. Lutz, and T. Chittenden.** 2000. Growth factors inactivate the cell death promoter BAD by phosphorylation of its BH3 domain on Ser(155). *Journal of Biological Chemistry* **275**:25046-25051.
39. **Danial, N. N., C. F. Gramm, L. Scorrano, C. Y. Zhang, S. Krauss, A. M. Ranger, S. R. Datta, M. E. Greenberg, L. J. Licklider, B. B. Lowell, S. P. Gygi, and S. J. Korsmeyer.** 2003. BAD and glucokinase reside in a mitochondrial complex that integrates glycolysis and apoptosis. *Nature* **424**:952-956.
40. **Edinger, A. L.** 2005. Growth factors regulate cell survival by controlling nutrient transporter expression. *Biochemical Society Transactions* **33**:225-227.
41. **Gottlob, K., N. Majewski, S. Kennedy, E. Kandel, R. B. Robey, and N. Hay.** 2001. Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. *Genes & Development* **15**:1406-1418.
42. **Vander Heiden, M. G., D. R. Plas, J. C. Rathmell, C. J. Fox, M. H. Harris, and C. B. Thompson.** 2001. Growth factors can influence cell growth and survival through effects on glucose metabolism. *Molecular and Cellular Biology* **21**:5899-5912.
43. **Forde, J. E. and T. C. Dale.** 2007. Glycogen synthase kinase 3: A key regulator of cellular fate. *Cellular and Molecular Life Sciences* **64**:1930-1944.
44. **Joje, R. S. and G. V. W. Johnson.** 2004. The glamour and gloom of glycogen synthase kinase-3. *Trends in Biochemical Sciences* **29**:95-102.
45. **Hashimoto, M., Y. Sagara, D. Langford, I. P. Everall, M. Mallory, A. Everson, M. Digicaylioglu, and E. Masliah.** 2002. Fibroblast growth factor 1 regulates signaling via the glycogen synthase kinase-3 beta pathway - Implications for neuroprotection. *Journal of Biological Chemistry* **277**:32985-32991.
46. **Sinha, D., K. Ruchalski, Z. Y. Wang, J. S. Levine, S. C. Borkan, and J. H. Schwartz.** 2003. Lithium activation of the WNT signaling and p13 kinase-Akt pathways promote survival of growth. factors deprived cultured proximal tubular cells. *Journal of the American Society of Nephrology* **14**:325A.
47. **Somervaille, T. C. P., D. C. Linch, and A. Khwaja.** 2000. Survival signalling in primary human erythroblasts occurs through PI3-kinase/Akt/GSK3 and not the classical MAP kinase pathway. *Blood* **96**:79A.
48. **Somervaille, T. C. P., D. C. Linch, and A. Khwaja.** 2001. Growth factor withdrawal from primary human erythroid progenitors induces apoptosis through a pathway involving glycogen synthase kinase-3 and Bax. *Blood* **98**:1374-1381.
49. **Palacios, R. and M. Steinmetz.** 1985. IL3-Dependent Mouse Clones That Express B-220 Surface-Antigen, Contain Ig Genes in Germ-Line Configuration, and Generate Lymphocytes-B In vivo. *Cell* **41**:727-734.
50. **Krishan, A.** 1990. Rapid DNA content analysis by the propidium iodide-hypotonic citrate method. *Methods Cell Biol.* **33**:121-125.
51. **Nicoletti, I., G. Migliorati, M. C. Pagliacci, F. Grignani, and C. Riccardi.** 1991. A Rapid and Simple Method for Measuring Thymocyte Apoptosis by Propidium Iodide Staining and Flow-Cytometry. *Journal of Immunological Methods* **139**:271-279.
52. **Oray, B. and S. J. Norton.** 1982. Glyoxalase-I from Mouse-Liver. *Methods in Enzymology* **90**:542-546.
53. **McLellan, A. C., S. A. Phillips, and P. J. Thornalley.** 1992. Fluorometric Assay of D-Lactate. *Analytical Biochemistry* **206**:12-16.
54. **Campisi, J., G. Morreo, and A. B. Pardee.** 1984. Kinetics of G1 Transit Following Brief Starvation for Serum Factors. *Experimental Cell Research* **152**:459-466.

55. **Yen, A. and A. B. Pardee.** 1978. Exponential 3T3-Cells Escape in Mid-G1 from Their High Serum Requirement. *Experimental Cell Research* **116**:103-113.
56. **Yen, A. and A. B. Pardee.** 1977. Location of 3T3 Cell-Cycle Arrests Induced by Low Serum and Isoleucine Deprivation. *Journal of Cell Biology* **75**:A2.
57. **Zetterberg, A. and O. Larsson.** 1985. Kinetic-Analysis of Regulatory Events in G1 Leading to Proliferation Or Quiescence of Swiss 3T3 Cells. *Proceedings of the National Academy of Sciences of the United States of America* **82**:5365-5369.
58. **Jimenez, B., M. Arends, P. Esteve, R. Perona, R. Sanchez, S. R. Cajal, A. Wyllie, and J. C. Lacal.** 1995. Induction of Apoptosis in Nih3T3 Cells After Serum Deprivation by Overexpression of Rho-P21, A Gtpase Protein of the Ras Superfamily. *Oncogene* **10**:811-816.
59. **Kulkarni, G. V. and C. A. G. McCulloch.** 1994. Serum Deprivation Induces Apoptotic Cell-Death in A Subset of Balb/C 3T3 Fibroblasts. *Journal of Cell Science* **107**:1169-1179.
60. **Liu, W. H. and B. Y. M. Yung.** 1998. Mortalization of human promyelocytic leukemia HL-60 cells to be more susceptible to sodium butyrate induced apoptosis and inhibition of telomerase activity by down-regulation of nucleophosmin/B23. *Oncogene* **17**:3055-3064.
61. **Notarbartolo, M., M. Cervello, L. Dusonchet, A. Cusimano, and N. D'Alessandro.** 2002. Resistance to diverse apoptotic triggers in multidrug resistant HL60 cells and its possible relationship to the expression of P-glycoprotein, Fas and of the novel anti-apoptosis factors IAP (inhibitory of apoptosis proteins). *Cancer Letters* **180**:91-101.
62. **Alt, J. R., J. L. Cleveland, M. Hannink, and J. A. Diehl.** 2000. Phosphorylation-dependent regulation of cyclin D1 nuclear export and cyclin D1-dependent cellular transformation. *Genes & Development* **14**:3102-3114.
63. **Diehl, J. A., M. G. Cheng, M. F. Roussel, and C. J. Sherr.** 1998. Glycogen synthase kinase 3 beta regulates cyclin D1 proteolysis and subcellular localization. *Genes & Development* **12**:3499-3511.
64. **Cross, D. A. E., A. A. Culbert, K. A. Chalmers, L. Facci, S. D. Skaper, and A. D. Reith.** 2001. Selective small-molecule inhibitors of glycogen synthase kinase-3 activity protect primary neurones from death. *Journal of Neurochemistry* **77**:94-102.
65. **Klein, P. S. and D. A. Melton.** 1996. A molecular mechanism for the effect of lithium on development. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8455-8459.
66. **Aguilera, J. and J. A. Prieto.** 2001. The *Saccharomyces cerevisiae* aldose, reductase is implied in the metabolism of methylglyoxal in response to stress conditions. *Current Genetics* **39**:273-283.
67. **Aguilera, J. and J. A. Prieto.** 2004. Yeast cells display a regulatory mechanism in response to methylglyoxal. *Fems Yeast Research* **4**:633-641.
68. **Yao, D., T. Taguchi, T. Matsumura, R. Pestell, D. Edelstein, I. Giardino, G. Suske, N. Rabbani, P. J. Thornalley, V. P. Sarthy, H. P. Hammes, and M. Brownlee.** 2007. High glucose increases angiotensin-2 transcription in microvascular endothelial cells through methylglyoxal modification of mSin3A. *Journal of Biological Chemistry* **282**:31038-31045.
69. **Yao, D. C., T. Taguchi, T. Matsumura, R. Pestell, D. Edelstein, I. Giardino, G. Suske, N. Ahmed, P. J. Thornalley, V. P. Sarthy, H. P. Hammes, and M. Brownlee.** 2006. Methylglyoxal modification of mSin3A links glycolysis to angiotensin-2 transcription. *Cell* **124**:275-286.
70. **Oya-Ito, T., B. F. Liu, and R. H. Nagaraj.** 2006. Effect of methylglyoxal modification and phosphorylation on the chaperone and anti-apoptotic properties of heat shock protein 27. *Journal of Cellular Biochemistry* **99**:279-291.
71. **Sakamoto, H., T. Mashima, K. Yamamoto, and T. Tsuruo.** 2002. Modulation of heat-shock protein 27 (Hsp27) anti-apoptotic activity by methylglyoxal modification. *Journal of Biological Chemistry* **277**:45770-45775.
72. **van Heijst, J. W. J., H. W. M. Niessen, R. J. Musters, V. W. M. van Hinsbergh, K. Hoekman, and C. G. Schalkwijk.** 2006. Argpyrimidine-modified Heat Shock Protein 27 in human non-small cell lung cancer: A possible mechanism for evasion of apoptosis. *Cancer Letters* **241**:309-319.
73. **Oya, T., N. Hattori, Y. Mizuno, S. Miyata, S. Maeda, T. Osawa, and K. Uchida.** 1999. Methylglyoxal modification of protein - Chemical and immunochemical characterization of methylglyoxal-arginine adducts. *Journal of Biological Chemistry* **274**:18492-18502.
74. **Cornelis, S., Y. Bruynooghe, G. Van Loo, X. Saelens, P. Vandenabeele, and R. Beyaert.** 2005. Apoptosis of hematopoietic cells induced by growth factor withdrawal is associated with caspase-9 mediated cleavage of Raf-1. *Oncogene* **24**:1552-1562.
75. **Akhand, A. A., K. Hossain, H. Mitsui, M. Kato, T. Miyata, R. Inagi, J. Du, K. Takeda, Y. Kawamoto, H. Suzuki, K. Kurokawa, and I. Nakashima.** 2001. Glyoxal and methylglyoxal trigger distinct signals for map family kinases and caspase activation in human endothelial cells. *Free Radical Biology and Medicine* **31**:20-30.
76. **Che, W. Y., M. Asahi, M. Takahashi, H. Kaneto, A. Okado, S. Higashiyama, and N. Taniguchi.** 1997. Selective induction of heparin-binding epidermal growth factor-like growth factor by methylglyoxal and 3-deoxyglucosone in rat aortic smooth muscle cells - The involvement of reactive oxygen species formation and a possible implication for atherogenesis in diabetes. *Journal of Biological Chemistry* **272**:18453-18459.
77. **Di Loreto, S., V. Caracciolo, S. Colafarina, P. Sebastiani, A. Gasbarri, and F. Amicarelli.** 2004. Methylglyoxal induces oxidative stress-dependent cell injury and up-regulation of interleukin-10 and nerve growth factor in cultured hippocampal neuronal cells. *Brain Research* **1006**:157-167.
78. **Du, X. L., D. Edelstein, L. Rossetti, I. G. Fantus, H. Goldberg, F. Ziyadeh, J. Wu, and M. Brownlee.** 2000. Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proceedings of the*

- National Academy of Sciences of the United States of America **97**:12222-12226.
79. **Lee, C., M. B. Yim, P. B. Chock, H. S. Yim, and S. O. Kang.** 1998. Oxidation-reduction properties of methylglyoxal-modified protein in relation to free radical generation. *Journal of Biological Chemistry* **273**:25272-25278.
80. **Ortwerth, B. J., H. James, G. Simpson, and M. Linetsky.** 1998. The generation of superoxide anions in glycation reactions with sugars, ozones, and 3-deoxyosones. *Biochemical and Biophysical Research Communications* **245**:161-165.
81. **Yim, H. S., S. O. Kang, Y. C. Hah, P. B. Chock, and M. B. Yim.** 1995. Free-Radicals Generated During the Glycation Reaction of Amino-Acids by Methylglyoxal - A Model Study of Protein-Cross-Linked Free-Radicals. *Journal of Biological Chemistry* **270**:28228-28233.
82. **Choudhary, D., D. Chandra, and R. K. Kale.** 1997. Influence of methylglyoxal on antioxidant enzymes and oxidative damage. *Toxicology Letters* **93**:141-152.
83. **Di Loreto, S., V. Zimmitti, P. Sebastiani, C. Cervelli, S. Falone, and F. Amicarelli.** 2007. Methylglyoxal causes strong weakening of detoxifying capacity and apoptotic cell death in rat hippocampal neurons. *Int.J.Biochem.Cell Biol.*
84. **Park, Y. S., Y. H. Koh, M. Takahashi, Y. Miyamoto, K. Suzuki, N. Dohmae, K. Takio, K. Honke, and N. Taniguchi.** 2003. Identification of the binding site of methylglyoxal on glutathione peroxidase: Methylglyoxal inhibits glutathione peroxidase activity via binding to glutathione binding sites Arg 184 and 185. *Free Radical Research* **37**:205-211.
85. **de Arriba, S. G., G. Stuchbury, J. Yarin, J. Burnell, C. Loske, and G. Munch.** 2007. Methylglyoxal impairs glucose metabolism and leads to energy depletion in neuronal cells - protection by carbonyl scavengers. *Neurobiology of Aging* **28**:1044-1050.
86. **Rosca, M. G., T. G. Mustata, M. T. Kinter, A. M. Ozdemir, T. S. Kern, L. I. Szweda, M. Brownlee, V. M. Monnier, and M. F. Weiss.** 2005. Glycation of mitochondrial proteins from diabetic rat kidney is associated with excess superoxide formation. *American Journal of Physiology-Renal Physiology* **289**:F420-F430.
87. **Sagrista, M. L., A. F. Garcia, M. A. De Madariaga, and M. Mora.** 2002. Antioxidant and pro-oxidant effect of the thiolic compounds N-acetyl-L-cysteine and glutathione against free radical-induced lipid peroxidation. *Free Radical Research* **36**:329-340.
88. **Chaplen, F. W. R., W. E. Fahl, and D. C. Cameron.** 1998. Evidence of high levels of methylglyoxal in cultured Chinese hamster ovary cells. *Proceedings of the National Academy of Sciences of the United States of America* **95**:5533-5538.
89. **Pap, M. and G. M. Cooper.** 1998. Role of glycogen synthase kinase-3 in the phosphatidylinositol 3-kinase/Akt cell survival pathway. *Journal of Biological Chemistry* **273**:19929-19932.
90. **Renaud, F., S. Desset, L. Oliver, G. GimenezGallego, E. VanObberghen, Y. Courtois, and M. Laurent.** 1996. The neurotrophic activity of fibroblast growth factor 1 (FGF1) depends on endogenous FGF1 expression and is independent of the mitogen-activated protein kinase cascade pathway. *Journal of Biological Chemistry* **271**:2801-2811.
91. **Conlon, I. and M. Raff.** 1999. Size control in animal development. *Cell* **96**:235-244.
92. **Bauer, D. E., M. H. Harris, D. R. Plas, J. J. Lum, P. S. Hammerman, J. C. Rathmell, J. L. Riley, and C. B. Thompson.** 2004. Cytokine stimulation of aerobic glycolysis in hematopoietic cells exceeds proliferative demand. *Faseb Journal* **18**:1303-+.
93. **Lum, J. J., D. E. Bauer, M. Kong, M. H. Harris, C. Li, T. Lindsten, and C. B. Thompson.** 2005. Growth factor regulation of autophagy and cell survival in the absence of apoptosis. *Cell* **120**:237-248.
94. **Kan, O., S. A. Baldwin, and A. D. Whetton.** 1994. Apoptosis Is Regulated by the Rate of Glucose-Transport in An Interleukin-3 Dependent Cell-Line. *Journal of Experimental Medicine* **180**:917-923.
95. **Akhand, A. A., K. Hossain, M. Kato, T. Miyata, J. Du, H. Suzuki, K. Kurokawa, and I. Nakashima.** 2001. Glyoxal and methylglyoxal induce aggregation and inactivation of ERK in human endothelial cells. *Free Radical Biology and Medicine* **31**:1228-1235.
96. **Portero-Otin, M., R. Pamplona, M. J. Bellmunt, M. C. Ruiz, J. Prat, R. Salvayre, and A. Negre-Salvayre.** 2002. Advanced glycation end product precursors impair epidermal growth factor receptor signaling. *Diabetes* **51**:1535-1542.

Part II:

Experimental Data and Results

Chapter 7:

Phosphorylation of Glyoxalase I during the cell cycle

Chapter 7: Glyoxalase I is a direct substrate for Ca^{2+} /Calmodulin Dependent Kinase II (CaMKII)

INTRODUCTION

Despite the ample research on the glyoxalase (GLO) system and GLO1 in particular, most of the focus has been on the expression levels and activity, and relatively little is known about posttranslational modification. For human GLO1 it has been described that NO, in cooperation with glutathione, is able to induce modification of GLO1, resulting in a decreased activity and an isoelectric shift towards the basic end on two-dimensional gels (1,2). Similar effects have been described later to occur in yeast and *in vitro* as well (3). Phosphorylation has been described in yeast (4) and mammalian cells ((5,6); chapters 6 and 8 of this thesis), and in plants as well (7,8). Although several different kinases, such as PKA (5,6), GSK-3 β (chapter 6) and the PI3K/Akt/mTOR pathway (see chapter 8), have been shown to be involved in the phosphorylation of GLO1 in mammalian cells, to date, it has not been described which kinase can directly phosphorylate GLO1.

In *Arabidopsis*, GLO1 was found to be phosphorylated upon overexpression of SNF1-related protein kinase 2.8 (SnRK2.8 (8)). *In vitro* kinase assays shows that SnRK2.8 directly phosphorylated GLO1 and a concomitant increase of GLO1 activity was found upon phosphorylation. The SnRK family members are the plant homologues of AMPK (AMP-activated protein kinase) and play a broad role in growth and metabolic responses to cellular stress (9). Quite interestingly, SnRK2.8 has been shown to be important in the response to drought and salinity (10), a property shared with GLO1 (11-13). Also, in rice leaf, GLO1 was identified as a phosphoprotein (7). Phosphorylation was induced by gibberellin, which is known to be an important regulator of cell growth, and was mediated by a putative 54-kDa Ca^{2+} dependent protein kinase (CDPK). Furthermore it was shown earlier that Ca^{2+} and calmodulin (CaM) were involved in light-mediated GLO1 activity in plants (14,15) and inhibitors of CaM showed an inhibitory effect on GLO1 activity (16). The data presented in this chapter provide evidence for the phosphorylation of mammalian GLO1 by Ca^{2+} /calmodulin-dependent kinase II (CaMKII). Of note is also the shared consensus sequence for phosphorylation among CaMKII (17) and SnRK2.8 (18,19).

The goal of this study was to find the kinase that directly phosphorylates GLO1. Our data demonstrate that GLO1 is a direct substrate for CaMKII *in vitro*, and CaMKII is able to phosphorylate GLO1 *in vivo*. We could however not yet establish a role for CaMKII in GLO1 phosphorylation induced by IL-3 starvation of Ba/F3 cells, as KN-93, a specific inhibitor for CaMKII, was unable to inhibit IL-3 induced apoptosis.

RESULTS AND DISCUSSION

GLO1 is phosphorylated upon coexpression with GSK-3 β in Hek293 cells

As described in chapter 6 of this thesis, we showed earlier that inhibition of GSK-3 β inhibited the phosphorylation of GLO1 which is induced by IL-3 starvation of Ba/F3 cells. To further study the phosphorylation of GLO1, we cotransfected Hek293 cells with an expression vector encoding wild type GLO1 and a vector containing GSK-3 β . Cells were lysed 24 hours after transfection and phosphorylation was examined by 2-dimensional electrophoresis (2-DE) using an antibody raised against rabbit GLO1. Overexpression of GLO1 in itself already induced multiple phosphorylations, as shown by the many isoforms observed in Figure 7.1.A (middle panel; the non-phosphorylated, non-NO-responsive form is designated as the α -isoform). However, when GLO1 was cotransfected with a GSK-3 β cDNA, a strong phosphorylation was induced on the NO-responsive isoforms (Figure 7.1.A, indicated with an arrow). Subsequently the same cotransfection was performed as described, but 24 hours after transfection the serum was depleted for 2 hours before cell lysis in order to fully activate GSK-3 β and further mimic the conditions from chapter 6. Under these conditions, however, GSK-3 β only showed a minor effect on the phosphorylation of the NO-responsive isoform of GLO1 (Figure 7.1.B; indicated with an arrow).

CamKII phosphorylates GLO1 *in vitro*

Next, we studied whether GLO1 is a direct substrate for GSK-3 β . Analysis of the amino acid sequence shows the presence of the typical GSK-3 β consensus site (consecutive Ser/Thr residues, separated by three amino acids), with Ser94 and Thr98 being the potential sites for phosphorylation by GSK-3 β . This site is located at the interface of the GLO1 dimer and is conserved between human and mouse (Figure 7.2.A; the GSK-3 β consensus site is underlined; for more information about potential phosphorylation sites see also supplementary figure page 114). We therefore performed *in vitro* kinase assays using purified recombinant GLO1 from *E. coli*. However, as shown in figure 7.2.B, we were unable to detect phosphorylation of GLO1 by GSK-3 β *in vitro*.

The consensus sequence for GSK-3 β phosphorylation however includes a serine or threonine residue that is already phosphorylated (primed) by another kinase (S/T-X-X-X-phospho-S/T) (20). Analysis of the amino acid sequence from GLO1 reveals the presence of a predicted CaMKII consensus site at Thr98 ((R-X-X-S/T) (17); Figure 7.2.A; indicated by a box). This CaMKII consensus site overlaps with the predicted consensus site for GSK-3 β and could therefore serve as a priming site for GSK-3 β . Of note, the CaMKII consensus site is identical to the consensus site for the plant kinase SnRK2.8, which has been shown to be a direct kinase for GLO1 in *Arabidopsis* (8).

To evaluate whether CaMKII could phosphorylate GLO1 we first performed *in vitro*




Figure 7.1. Coexpression of GLO1 with GSK-3 β .

A. Coexpression of GLO1 with kinases in Hek293 cells. Hek293 cells were cotransfected with expression vectors encoding GLO1 and expression vectors containing GSK-3 β . Cells were lysed in CEB after 24 hours and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 10 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with a polyclonal anti-human GLO1 antibody are shown. A reference spot α , which corresponds to the non-NO-responsive form of GLO1 is indicated (5). **B.** Coexpression of GLO1 with kinases in serum-starved Hek293 cells. Hek293 cells were co-transfected with expression vectors encoding GLO1 and expression vectors containing GSK-3 β . 24 hours after transfection cells were depleted of serum for 2 hours. Cells were lysed in CEB and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 10 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with a polyclonal anti-human GLO1 antibody are shown.

kinase assays using purified recombinant GLO1 from *E. coli*. When recombinant GLO1 was incubated in reaction buffer alone in the presence of 32 P-ATP, no phosphorylation could be observed (Figure 7.3.A). The presence of contaminating kinases or a possible autophosphorylation activity could thus be excluded. Coincubation of GLO1 with activated CaMKII, on the other hand, clearly induced phosphorylation (Figure 7.3.A).

We subsequently overexpressed and purified GLO1 from Hek293 cells for *in vitro* kinase assays. As shown in Figure 7.3.B, purified GLO1 from Hek293 cells could also be efficiently phosphorylated by CaMKII *in vitro*. Of note is the appearance of a high molecular weight band on the autoradiograph (indicated by a white arrowhead). Overlay of the autoradiograph with a Western blot performed with an anti-GLO1 antibody however revealed this represents a GLO1-dimer. Mammalian GLO1 requires dimerization for its activity (21) and this dimer is known to be extremely stable and highly resistant against




Figure 7.2. GLO1 is no direct target for GSK-3 β *in vitro*.

A. Kinase consensus sequences. Alignment of the sequence of human and mouse GLO1 shows the presence of the conserved consensus sequences for GSK-3 β (S/T-X-X-X-phospho-S/T; underlined) and CaMKII phosphorylation (R-X-X-S/T; box). Potential targets for phosphorylation are indicated in bold (phosphorylation sites are determined using Netphos software). **B.** Recombinant GLO1 is not phosphorylated by GSK-3 β . 2 μ g of recombinant GLO1 was incubated with GSK-3 β as described in methods. Phosphorylation was detected by autoradiography, followed by western blot with an anti-GLO1 antibody.

reducing conditions (22). These results suggest that phosphorylation does not inhibit dimerization. Inhibition of dimerization would lead to inactivation of GLO1 (21,23,24).

The *in vitro* phosphorylation assays described above were performed on GLO1 which was purified using hexyl-glutathione linked beads and subsequently eluted using 10 mM s-hexylglutathione, dissolved in CaMKII reaction buffer. However, for the required sequential phosphorylation reactions, the priming with CaMKII and the phosphorylation with GSK-3 β , the phosphorylation activity of the first kinase needs to be eliminated. This can be achieved by heat inactivation of the enzyme. Alternatively, GLO1 which is still bound to the hexyl-glutathione beads, can be used for the *in vitro* assays, with the advantage that the first kinase can be washed away. An additional benefit would be the possibility to use different reaction buffers for the sequential reactions. We therefore tested an alternative protocol to examine whether GLO1 which was bound to beads could also serve as a substrate for our assays.

When protein-loaded beads were coincubated with activated CaMKII, GLO1 was phosphorylated by CaMKII (Figure 7.3.C left panel). This demonstrates that the binding of GLO1 to beads does not interfere with phosphorylation. Of note, when lysis buffer in which phosphatase inhibitors were omitted was used to lyse cells and wash the beads, phosphorylation of GLO1 by CaMKII could further be enhanced, most likely because any phosphorylation that might have occurred in the cells was removed or reduced (Figure





Figure 7.3. GLO1 is a direct substrate for CaMKII *in vitro*.

A. Recombinant GLO1 is phosphorylated by CaMKII. 2 μg of recombinant wild type GLO1 was incubated with activated CaMKII as described in methods. Phosphorylation was detected by autoradiography, followed by western blot with an anti-GLO1 antibody. **B.** Purified GLO1 from Hek293 is phosphorylated with CaMKII. Hek293 cells were transfected with expression vectors containing wild type GLO1. After 24 hours, transfected cells were lysed in CEB and GLO1 was purified as described in methods. Purified GLO1 was subsequently incubated with activated CaMKII. Phosphorylation was detected by autoradiography, followed by western blot with an anti-GLO1 antibody. Monomer GLO1 is indicated with a black arrowhead, dimer GLO1 is indicated with a white arrowhead. **C.** GLO1-loaded beads are phosphorylated by CaMKII. Hek293 cells were transfected with expression vectors containing wild type GLO1. After 24 hours, transfected cells were lysed in CEB (left panel) or in CEB without phosphatase inhibitors (right panel) and GLO1 was purified as described in methods. Protein-loaded beads were subsequently incubated with activated CaMKII. Phosphorylation was detected by radiography, followed by western blot with an anti-GLO1 antibody.

7.3.C right panel).

Subsequently we tested the hypothesis that CaMKII-dependent phosphorylation could serve as a priming phosphorylation for GSK-3 β . Recombinant GLO1 from *E. coli* was first coincubated with activated CaMKII in the absence of ^{32}P -ATP for 30 minutes. Then CaMKII was heat inactivated and the reaction mixture was incubated with GSK-3 β , in the presence of ^{32}P -ATP. Concomitantly unprimed GLO1 was coincubated with GSK-3 β . As shown in Figure 7.4.A, no phosphorylation could be observed with GSK-3 β , not even when GLO1 was first primed with CaMKII. However, CaMKII itself significantly phosphorylated GLO1. Also when GLO1 was purified from Hek293 cells and GLO1 bound to beads was used

A.



B.




Figure 7.4. Primed GLO1 is no direct target for GSK-3 β *in vitro*.

A. Recombinant GLO1 is not phosphorylated by GSK-3 β after priming with CaMKII. 2 μ g of recombinant wild type GLO1 was incubated with activated CaMKII in the absence of ³²P-ATP as described. After heat inactivation, the samples were incubated with GSK-3 β in the presence of ³²P-ATP. As a control 2 μ g of recombinant GLO1 was directly incubated with either activated CaMKII or GSK-3 β . Phosphorylation was detected by autoradiography, followed by western blot with an anti-GLO1 antibody. **B.** Recombinant GLO1 is not phosphorylated by PKA. 2 μ g of recombinant GLO1 was incubated with PKA as described in methods. Phosphorylation was detected by autoradiography, followed by western blot with an anti-GLO1 antibody.

for *in vitro* phosphorylation, no phosphorylation could be observed with GSK-3 β (data not shown).

Another kinase that has been described to be involved in the phosphorylation of GLO1 is PKA (5,6). In spite of the presence of several consensus sites in the amino acid sequence for PKA (not shown), we were unable to detect *in vitro* phosphorylation of GLO1 with PKA (Figure 7.4.B).




Figure 7.5. Inhibition of CaMKII does not inhibit IL-3 starvation-induced apoptosis.

Effect of CaMKII inhibition on IL-3 depletion-induced apoptosis. Ba/F3 cells were depleted of IL-3 and KN-93 (10 μ M) was added. Apoptosis was measured by the analysis of the percentage of cells with sub-diploid DNA at the indicated time points.

Involvement of CaMKII in IL-3 depletion-induced apoptosis

To assess whether CaMKII was involved in IL-3 starvation-induced apoptosis, Ba/F3 cells were depleted of IL-3 in the presence of KN-93, a specific inhibitor of CaMKII. Apoptosis was subsequently determined by measuring DNA breakdown. However, inhibition of CaMKII had no effect on apoptosis induced by IL-3 depletion (Figure 7.5).

CONCLUSIONS

Although the GLO system has been discovered almost a hundred years ago, the only known function is the detoxification of reactive α -oxoaldehydes (25). Identification of the kinases involved in the modification of GLO1 might shed a light over the precise role of the GLO system. GLO1 has been shown to be phosphorylated in mammals, plants and in yeast (5-8,26). In spite of the involvement of several signaling pathways, to date, the kinase responsible for the phosphorylation of GLO1 has not been described in mammals. In Chapter 6 it was described how inhibition of GSK-3 β also inhibited the IL-3 depletion-induced phosphorylation of GLO1. These observations were corroborated here by the observation that coexpression of GLO1 with GSK-3 β also induces phosphorylation of GLO1. On the other hand, *in vitro* phosphorylation assays revealed that GLO1 is no direct substrate for GSK-3 β . However, our data shows that GLO1 is a direct substrate for CaMKII. Although the predicted consensus sites for CaMKII and GSK-3 β seem to overlap, CaMKII does not seem to be a priming kinase for GSK-3 β phosphorylation. It can not be ruled out however that GSK-3 β can directly phosphorylate GLO1, but that there is a different priming kinase.

CaMKII is a ubiquitously expressed kinase family, which needs Ca^{2+} and calmodulin for its activation (27). CaMKII is involved in a wide range of cellular processes, among which cell growth and division (28-30), and cell death (28-31), making it a likely candidate to be involved in TNF-induced necrosis and IL-3 starvation-induced apoptosis. However, inhibition of CaMKII did not exert an effect in the induction of apoptosis induced by IL-3 depletion of Ba/F3 cells. Nevertheless, the involvement of CaMKII in the phosphorylation of GLO1 *in vivo* was demonstrated in TNF-induced necrosis in the mouse fibroblast cell line L929 (de Hemptinne et al., in preparation).

CaMKII is probably not the only kinase capable of phosphorylating GLO1. From the 2-DE pattern of GLO1 it could be concluded that, for instance TNF can induce multiple phosphorylations of GLO1 (5). Further research is therefore still needed to fully elucidate all the kinases involved in modification of GLO1.

MATERIALS AND METHODS

Cell Lines and Cultures

The IL-3 dependent pro-B cell line Ba/F3 (32) was purchased from DSMZ. Ba/F3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) with glutamax (Invitrogen) supplemented with heat-inactivated FBS (10% v/v) (Cambrex), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) and 5% conditioned medium from WEHI-3B cells as a source of mouse IL-3. Ba/F3 cells were cultured at 37°C in a humidified incubator under an 5% CO_2 atmosphere. Hek293 cells, were cultured in DMEM with glutamax (Invitrogen) supplemented with heat-inactivated FBS (10% v/v), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) at 37°C in a humidified incubator under an 8% CO_2 atmosphere.

Cell stimulations

IL-3 was removed by washing exponentially growing cells 3 times in culture medium without conditioned WEHI-3B medium and resuspended in culture medium in the absence or presence of conditioned WEHI-3B.

Chemicals and inhibitors used are KN-93 (calbiochem). Duration and concentrations used as described.

Hek293 cells were transiently transfected by Ca precipitation. Vectors used are pCAGGS vectors containing wild type GLO1 and an expression vector for GSK-3 β , kindly donated by Dr Jim Woodgett (Department of Medical Biophysics, Samuel Lunenfeld Research Institute, Toronto, Ontario, Canada).

DNA distribution analysis

Cells were stained with propidium iodide (PI, Sigma) containing staining solution (33) and analyzed by a FACSCalibur flow cytometer (488_{Ex}/590_{Em}). Cell Quest software was used to analyze cell cycle distribution (FACS Calibur, Becton Dickinson) (33,34). Ten thousand cells were routinely analyzed.

Electrophoresis and immunoblotting

The cells were washed 3 times with ice-cold PBS buffer and lysed with cytosol extraction buffer (10 mM Tris-HCL pH7.4, 50 mM EDTA pH8.0, 25 mM NaCl, 0,7% TritonX-100, 100 mM PMSF, 1 tablet Complete Protease Inhibitor Cocktail (Roche)/50 ml cytosol extraction buffer). Cell lysates were cleared by centrifugation (14,000 x g). Protein concentrations were determined using Bradford method. Proteins were separated by SDS-PAGE and transferred onto a PVDF membrane (Hybond-P, Amersham Pharmacia Biotech). The blots were incubated with the desired antibodies followed by ECL-based detection (Amersham Pharmacia Biotech).

2-Dimensional Gel Electrophoresis

Isoelectric focusing was carried out on 18 cm IPG strips, pH 4-7 (GE Healthcare) according to the manufacturer's instructions. For the second dimension, proteins were separated by SDS-polyacrylamide gel electrophoresis (PAGE, 12%).

Recombinant expression and purification of GLO1 in E. coli

Wild type GLO1 was subcloned into pGEX-6P-1 GST fusion vector (Amersham Pharmacia Biotech) and wild type GLO1 was purified according to manufacturers instructions. Protein concentrations were determined using Bradford method. Aliquots were stored at -80°C until further use.

Purification of GLO1 from Hek293 cells for *in vitro* phosphorylation

Cell cultures were transfected and lysed as described above. The cell lysates were incubated with S-hexylglutathione-agarose beads (Sigma) for 30 minutes. The cell lysates were then removed and the beads were washed 4 times with PBS buffer. GLO1 was eluted from the beads using 10 mM s-hexylglutathione, dissolved in CaMKII reaction buffer.

Phosphorylation reactions

2 µg of recombinant GLO1 or 10 µL of purified GLO1 was incubated with activated CaMKII (New England Biolabs), GSK-3β (New England Biolabs) or PKA (New England Biolabs) in the respective kinase reaction buffer (New England Biolabs) in the presence of 5 µCi ³²P-γ-ATP. Reactions were stopped after 1 hour by adding 5x laemmli buffer and boiling at 95°C for 5 minutes.

Alternatively cells were lysed as described above. The cell lysates were incubated with S-hexylglutathione-agarose beads for 30 minutes. The cell lysates were then removed and the beads were washed 4 times with PBS buffer and twice in kinase reaction buffer. Protein-loaded beads were resuspended in kinase reaction buffer and incubated with activated CaMKII or GSK-3β in the presence of 5 µCi ³²P-γ-ATP at 30°C for 1 hour. Reactions were stopped by washing the beads in cytosol extraction buffer containing phosphatase inhibitors. Beads were resuspended in 5x laemmli and boiled for 5 minutes.

The phosphorylated GLO1 was analyzed by SDS-PAGE and autoradiography.

Reference List

1. **Mitsumoto, A., K. R. Kim, G. Oshima, M. Kunimoto, K. Okawa, A. Iwamatsu, and Y. Nakagawa.** 1999. Glyoxalase I is a novel nitric-oxide-responsive protein. *Biochemical Journal* **344**:837-844.
2. **Mitsumoto, A., K. R. Kim, G. Oshima, M. Kunimoto, K. Okawa, A. Iwamatsu, and Y. Nakagawa.** 2000. Nitric oxide inactivates glyoxalase I in cooperation with glutathione. *Journal of Biochemistry* **128**:647-654.
3. **Sahoo, R., R. Sengupta, and S. Ghosh.** 2003. Nitrosative stress on yeast: inhibition of glyoxalase-I and glyceraldehyde-3-phosphate dehydrogenase in the presence of GSNO. *Biochemical and Biophysical Research Communications* **302**:665-670.
4. **Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura.** 1989. Sexual-Response in *Saccharomyces-Cerevisiae* - Alteration of Enzyme-Activity in the Glyoxalase System by Mating Factor. *Biochemical and Biophysical Research Communications* **165**:1091-1095.
5. **de, H., V. D. Rondas, J. Vandekerckhove, and K. Vancompernelle.** 2007. Tumour necrosis factor induces phosphorylation primarily of the nitric-oxide-responsive form of glyoxalase I. *Biochem.J.* **407**:121-128.
6. **Van Herreweghe, F., J. Q. Mao, F. W. R. Chaplen, J. Grooten, K. Gevaert, J. Vandekerckhove, and K. Vancompernelle.** 2002. Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methyl glyoxal-derived AGE. *Proceedings of the National Academy of Sciences of the United States of America* **99**:949-954.
7. **Khan, M. M. K., A. Jan, H. Karibe, and S. Komatsu.** 2005. Identification of phosphoproteins regulated by gibberellin in rice leaf sheath. *Plant Molecular Biology* **58**:27-40.
8. **Shin, R., S. Alvarez, A. Y. Burch, J. M. Jez, and D. P. Schachtman.** 2007. Phosphoproteomic identification of targets of the Arabidopsis sucrose nonfermenting-like kinase SnRK2.8 reveals a connection to metabolic processes. *Proceedings of the National Academy of Sciences of the United States of America* **104**:6460-6465.
9. **Hardie, D. G., D. Carling, and M. Carlson.** 1998. The AMP-activated/SNF1 protein kinase subfamily: Metabolic sensors of the eukaryotic cell? *Annual Review of Biochemistry* **67**:821-855.
10. **Boudsocq, M., H. Barbier-Brygoo, and C. Lauriere.** 2004. Identification of nine sucrose nonfermenting 1-related protein kinases 2 activated by hyperosmotic and saline stresses in Arabidopsis thaliana. *Journal of Biological Chemistry* **279**:41758-41766.
11. **Singla-Pareek, S. L., M. K. Reddy, and S. K. Sopory.** 2003. Genetic engineering of the glyoxalase pathway in tobacco leads to enhanced salinity tolerance. *Proc.Natl.Acad.Sci.U.S.A* **100**:14672-14677.
12. **Yadav, S. K., S. L. Singla-Pareek, M. Ray, M. K. Reddy, and S. K. Sopory.** 2005. Methylglyoxal levels in plants under salinity stress are dependent on glyoxalase I and glutathione. *Biochemical and Biophysical Research Communications* **337**:61-67.
13. **Yadav, S. K., S. L. Singla-Pareek, M. K. Reddy, and S. K. Sopory.** 2005. Transgenic tobacco plants overexpressing glyoxalase enzymes resist an increase in methylglyoxal and maintain higher reduced glutathione levels under salinity stress. *FEBS Lett.* **579**:6265-6271.
14. **Chakravarty, T. N. and S. K. Sopory.** 1998. Blue light stimulation of cell proliferation and glyoxalase I activity in callus cultures of *Amaranthus paniculatus*. *Plant Science* **132**:63-69.
15. **Das, R., S. Bagga, and S. K. Sopory.** 1987. Involvement of Phosphoinositides, Calmodulin and Glyoxalase-I in Cell-Proliferation in Callus-Cultures of *Amaranthus-Paniculatus*. *Plant Science* **53**:45-51.
16. **Bagga, S., R. Das, and S. K. Sopory.** 1987. Inhibition of Cell-Proliferation and Glyoxalase-I Activity by Calmodulin Inhibitors and Lithium in Brassica-Oleracea. *Journal of Plant Physiology* **129**:149-153.
17. **Kennelly, P. J. and E. G. Krebs.** 1991. Consensus Sequences As Substrate-Specificity Determinants for Protein-Kinases and Protein Phosphatases. *Journal of Biological Chemistry* **266**:15555-15558.
18. **Furihata, T., K. Maruyama, Y. Fujita, T. Umezawa, R. Yoshida, K. Shinozaki, and K. Yamaguchi-Shinozaki.** 2006. Absciscic acid-dependent multisite phosphorylation regulates the activity of a transcription activator AREB1. *Proceedings of the National Academy of Sciences of the United States of America* **103**:1988-1993.
19. **Kobayashi, Y., M. Murata, H. Minami, S. Yamamoto, Y. Kagaya, T. Hobo, A. Yamamoto, and T. Hattori.** 2005. Absciscic acid-activated SNRK2 protein kinases function in the gene-regulation pathway of ABA signal transduction by phosphorylating ABA response element-binding factors. *Plant Journal* **44**:939-949.
20. **Forde, J. E. and T. C. Dale.** 2007. Glycogen synthase kinase 3: A key regulator of cellular fate. *Cellular and Molecular Life Sciences* **64**:1930-1944.
21. **Cameron, A. D., B. Olin, M. Ridderstrom, B. Mannervik, and T. A. Jones.** 1997. Crystal structure of human glyoxalase .1. Evidence for gene duplication and 3D domain swapping. *Embo Journal* **16**:3386-3395.
22. **Clapper, M. L., S. J. Hoffman, and K. D. Tew.** 1991. Glutathione S-Transferases in Normal and Malignant Human Colon Tissue. *Biochimica et Biophysica Acta* **1096**:209-216.
23. **Garciaíniguez, L., L. Powers, B. Chance, S. Sellin, B. Mannervik, and A. S. Mildvan.** 1984. X-Ray Absorption Studies of the Zn-2+ Site of Glyoxalase-I. *Biochemistry* **23**:685-689.
24. **Ridderstrom, M., A. D. Cameron, T. A. Jones, and B. Mannervik.** 1998. Involvement of an active-site Zn2+ ligand mechanism of human glyoxalase I. *Journal of Biological Chemistry* **273**:21623-21628.
25. **Thornalley, P. J.** 2003. Glyoxalase I - structure, function and a critical role in the enzymatic defence against glycation. *Biochemical Society Transactions* **31**:1343-1348.
26. **Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura.** 1990. Sexual-Response of *Saccharomyces-Cerevisiae* - Phosphorylation of Yeast Glyoxalase-I by A Cell Extract of Mating Factor Treated-Cells. *Journal of Biochemistry* **108**:4-6.
27. **Braun, A. P. and H. Schulman.** 1995. The Multifunctional Calcium Calmodulin-Dependent Protein-Kinase - from Form to Function. *Annual Review of Physiology* **57**:417-445.

28. **Karcher, R. L., J. T. Roland, F. Zappacosta, M. J. Huddleston, R. S. Annan, S. A. Carr, and V. Gelfand.** 2001. Cell cycle regulation of myosin-V by calcium/calmodulin-dependent protein kinase II. *Science* **293**:1317-1320.
29. **Ohta, Y., T. Ohba, K. Fukunaga, and E. Miyamoto.** 1988. Serum and Growth-Factors Rapidly Elicit Phosphorylation of the Ca-2+/Calmodulin-Dependent Protein Kinase-II in Intact Quiescent Rat 3Y1 Cells. *Journal of Biological Chemistry* **263**:11540-11547.
30. **Planassilva, M. D. and A. R. Means.** 1992. Expression of A Constitutive Form of Calcium Calmodulin Dependent Protein Kinase-II Leads to Arrest of the Cell-Cycle in G2. *Embo Journal* **11**:507-517.
31. **Mishra, S., J. P. Mishra, K. Gee, D. C. McManus, E. C. LaCasse, and A. Kumar.** 2005. Distinct role of calmodulin and calmodulin-dependent protein kinase-II in lipopolysaccharide and tumor necrosis factor-alpha-mediated suppression of apoptosis and antiapoptotic c-IAP2 gene expression in human monocytic cells. *Journal of Biological Chemistry* **280**:37536-37546.
32. **Palacios, R. and M. Steinmetz.** 1985. IL3-Dependent Mouse Clones That Express B-220 Surface-Antigen, Contain Ig Genes in Germ-Line Configuration, and Generate Lymphocytes-B *In vivo*. *Cell* **41**:727-734.
33. **Krishan, A.** 1990. Rapid DNA content analysis by the propidium iodide-hypotonic citrate method. *Methods Cell Biol.* **33**:121-125.
34. **Nicoletti, I., G. Migliorati, M. C. Pagliacci, F. Grignani, and C. Riccardi.** 1991. A Rapid and Simple Method for Measuring Thymocyte Apoptosis by Propidium Iodide Staining and Flow-Cytometry. *Journal of Immunological Methods* **139**:271-279.

Part II:

Experimental Data and Results

Chapter 8:

Phosphorylation of glyoxalase I during the cell cycle

Chapter 8: Phosphorylation of Glyoxalase I during the cell cycle

INTRODUCTION

The glyoxalase (GLO) system is comprised of two enzymes, GLO1 and GLO2. In spite of the fact that it has been discovered almost a hundred years ago (1,2), the only function conferred to the GLO system is the detoxification of reactive α -oxoaldehydes, such as MG (3). The GLO systems ubiquitous nature (4,5), however, suggest a more fundamental and conserved role. Moreover, several other enzyme systems exist that are capable of detoxifying α -oxoaldehydes (6), rendering the GLO system partially redundant for detoxification. Since the work of Albert Szent-Györgyi (5,7-9), the GLO system has often been connected to proliferation. A strong correlation has been observed between high GLO1 activity and the rate of proliferation (10,11). Elevated levels of GLO1 expression or activity have been documented as important for the progression of several tumors and have been proposed to be indicative for the enhanced proliferative status (12-20), although some reports disprove this (21). Also in plants GLO1 activity strongly correlates with cell growth (22-25) and stimulation of cell growth involves an increase in GLO1 activity (26).

The availability of growth factors (GFs) is determining in survival, proliferation and growth. The phosphatidylinositol 3-kinase (PI3K)/Akt pathway is involved in many cellular functions in response to GFs (27-30). A requisite for cell growth and division is the presence of glucose and a tight regulation of glucose metabolism (31,32). The involvement of the PI3K/Akt signaling pathway in the regulation of glucose uptake and the regulation of glycolysis has been well-documented (33-36). An important downstream target by which Akt exerts its influence on metabolism is mTOR (37-41). Furthermore, PI3K/Akt/mTOR signaling also directly regulates GF controlled cell cycle progression, primarily by controlling the expression of G₁ cyclins (42-45), but also at the end of G₂/M (46).

In order to study the possible role of GLO1 in the regulation of the cell cycle we analyzed the phosphorylation of GLO1 during the cell cycle. We show here that during progression through the cell cycle different phosphorylations arise. In line with previous reports (47), these phosphorylations could not be associated with a change in GLO1 activity *in vitro*. Inhibition of the PI3K/Akt partially inhibits the phosphorylation of GLO1. Also inhibition of mTOR, one of the downstream targets of the PI3K/Akt pathway involved in control of cellular growth, in part inhibits the phosphorylation of GLO1. These data implicate the PI3K/Akt/mTOR pathway, an important regulator of cell cycle in the modification of GLO1.

RESULTS

Cell cycle dependent phosphorylation of GLO1

Previous studies in our group show that most cell types express multiple isoforms of GLO1, including a NO-modified form and a non-NO-modified form, which was denominated the α -isoform (48). These different forms of GLO1 can be multifariously phosphorylated on the α -isoform as well as on the NO-modified form. These changes in isoform pattern can easily be studied using 2-dimensional gel electrophoresis (2-DE) and immunoblotting with an anti-GLO1 antibody (48). Phosphorylation has been shown to yield a shift of the isoform to the acidic side of the gel, whereas NO-induced modification results in a shift to the basic side.




Figure 8.1. Cell cycle dependent phosphorylation GLO1.

GLO1 was phosphorylated during G₁ (A) and at G₁/S transition (B). Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by re-addition of IL-3. This time point was designated 0 hours. Cells were lysed in CEB at indicated time points, and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with a polyclonal anti-human GLO1 antibody (48) are shown. A reference spot α , which corresponds to the non-NO-responsive form of GLO1 is indicated (48). C and D. Analysis of the cell cycle. The different phases of the cell cycle were monitored by analysis of the DNA content by flow cytometry.

In *Arabidopsis* and rice leaf, regulators of cell cycle have been reported to induce phosphorylation of GLO1 (49-51). To date, in mammalian cells, no posttranslational modifications of GLO1 have been reported in the cell cycle. In chapter 6 we have described how GLO1 is phosphorylated during GF withdrawal induced G_1 arrest. To study phosphorylation during cell cycle, the cell cycle of Ba/F3 cultures was synchronised in the G_1 phase by IL-3 depletion overnight and subsequently released from this block by re-addition of IL-3. Cells were lysed 1 hour, 3 hours and 5 hours after re-addition of IL-3 for the analysis of the posttranslational modifications of GLO1 in G_1 phase using 2-dimensional gel electrophoresis (2-DE) (Figure 8.1). Analysis of the DNA-distribution showed Ba/F3 cells resided in the G_1 phase after all three time intervals (Figure 8.1.C) In lysates made after 1 hour, only a small number of isoforms is displayed. The α -form could not be observed. Instead, the non-NO-modified form is strongly phosphorylated, as can be observed by the localisation of the spot more to the acidic side (Figure 8.1.A, upper panel, arrowhead). At this time, only a very limited presence of the NO-modified form of GLO1 could be detected. In lysates made 3 hours after the release of the cell cycle block, the non-NO-modified form of GLO1 is strongly dephosphorylated, as shown in a shift of the spot to the basic side. Concomitantly, NO-mediated modification of GLO1 occurs, as witnessed by the appearance of a more basic spot (Figure 8.1.A, middle panel, arrow). In the lysates made 5 hours after the re-addition of IL-3, both the NO-modified form of GLO1 and the non-NO-modified form of GLO1 were further dephosphorylated (Figure 8.1.A, bottom panel). The α -form is now visible.

Subsequently, we studied the phosphorylation of GLO1 during G_1 -S transition using a same approach (Figure 8.1.B). The percentage of cells in the different phases of the cell cycle was determined by analysis of the DNA-distribution on FACS, indicating at the chosen time points, cells pass from the G_1 phase into the S phase (Figure 8.1.D). Cells were lysed 11 hours, 12 hours and 13 hours after the release from G_1 arrest. 11 hours after the release from G_1 arrest, when 65 % of the cells is still in the G_1 phase (Figure 8.1.D, black bars for the percentage in the G_1 phase and white bars for the percentage in the S phase), we observed a strong presence of several phosphorylated isoforms on the NO-responsive form (Figure 8.1.C, upper panel, arrows). After 12 hours 50 % of the cell population was in the G_1 phase and 50 % of the cells was in the S phase (Figure 8.1.D). At this time point, a hyperphosphorylation of the NO-responsive isoforms could be detected (Fig 1.C, middle panel, arrow). After 13 hours, when 70 % of the cells were in the S phase and only 30 % still resides in the G_1 phase, the NO-responsive form was partially dephosphorylated again (Figure 8.1.C, lower panel, arrow).

All together, our data shows that GLO1 is strongly modified during the early stages of the cell cycle. Both phosphorylation and NO-modification occur. Of note is the observation that the G_1 -S transition coincides with a hyperphosphorylated state of the NO-modified

form of GLO1. This hyperphosphorylation is rapidly removed when cells pass on into the S phase.

GLO1 activity does not change throughout the cell cycle

Next, we determined whether the posttranslational modification of GLO1 was accompanied by changes in the activity of GLO1. *In vitro* GLO1 activity in lysates was assayed using a spectrophotometric method, which monitors the GLO1-catalyzed production of S-D-lactoylglutathione from the hemithioacetal that is spontaneously formed when MG is incubated with glutathione (see materials and methods).

Cells were synchronized by overnight IL-3 starvation and subsequent re-addition of IL-3. Cell lysates were made after the indicated time intervals after re-addition of IL-3 and *in vitro* GLO1 activity was assayed (Figure 8.2.A). The chosen time points are situated at the G_1 -S transition (Figure 8.1.D). Although hyperphosphorylation and consequent




Figure 8.2. *In vitro* GLO1 activity is not altered during the cell cycle.

A. GLO1 activity at G_1 /S transition. Ba/F3 cells were synchronized by IL-3 starvation over night. The cell cycle was induced by re-addition of IL-3. This time point was designated 0 hours. Cells were lysed in CEB at indicated time points after the re-addition of IL-3, lysates were brought to equal concentrations and cell lysates were used to assay GLO1 activity. *In vitro* GLO1 reaction rates were determined using Ultraspec 1100 software (Amersham, BD Biosciences) and are shown relative to the reaction rate of lysates made of exponentially growing cells, which are set as 100%. **B.** GLO1 activity in different phases of the cell cycle. G_1 arrest was induced by IL-3 starvation of Ba/F3 cells and G_2/M arrest was induced by addition of nocodazole. Cells were lysed in CEB, lysates were brought to equal concentrations and cell lysates were used to assay GLO1 activity. **C.** Analysis of the cell cycle. The different phases of the cell cycle were monitored by analysis of the DNA content by flow cytometry.

dephosphorylation of GLO1 can be observed at the G_1 -S transition (Figure 8.1.C), a concomitant change in GLO1 activity could not be observed (Figure 8.2.A). Some fluctuation can be noticed in GLO1 activity. However, this was not significant and is probably caused by experimental variability.

In order to further compare GLO1 activity in different phases of the cell cycle, we arrested Ba/F3 cells in the G_1 phase and in G_2 /M, respectively by IL-3 depletion or addition of nocodazole for 8 hours. Nocodazole inhibits microtubule polymerization and blocks mitotic spindle formation, resulting in the accumulation of cells in M. The arrest in respective cell cycle phases was verified by analysis of the DNA distribution (Figure 8.2.C). Cell lysates were prepared and *in vitro* GLO1 activity subsequently assayed (Figure 8.2.B). Neither G_1 arrest or G_2 /M arrest was accompanied by a change in *in vitro* GLO1 activity when compared to a culture of asynchronously growing cells.

We therefore conclude that GLO1 activity as assayed by its ability to catalyze the formation of S-D-lactoylglutathione, is not modulated by post-translational modification. These results are in line with our previous observations in Ba/F3 cells (chapter 6), but also in L929 cells (47) and bovine aortic endothelial cells (BAEC; Laga et al., submitted). Our findings however contradict the reported rise in GLO1 activity observed during G_2 /M in carrot cell suspension (52).

Phosphorylation of GLO1 is mediated by the PI3K/Akt pathway

IL-3 has been shown to induce activation of several signaling pathways, including Jak-2/STAT-5, ras/raf/MEK/ERK, and PI3K/Akt kinase cascades (53-55). The PI3K/Akt signaling pathway has been implicated in regulating metabolism and proliferation (56-59). We therefore first investigated whether the phosphorylation of GLO1 was dependent on PI3K/Akt signaling.

Ba/F3 cells were synchronized as described previously. 8 hours after the release from cell cycle arrest, three dominant spots can be observed. The α -form of GLO1, and two more acidic spots. Only little NO-responsive GLO1 is present. 11 hours after re-addition of IL-3, the α -form of GLO1 and the middle isoform both are phosphorylated (Figure 8.3.A, second panel, black arrowhead and white arrowhead respectively). Also NO-mediated modification of GLO1 is induced at that time (Figure 8.3.A, second panel, arrow). One hour later, the most acidic spot shifts to the left, indicating this form of GLO1 is subject to additional phosphorylation (Figure 8.3.A, third panel, arrowhead), and the phosphorylation on the α -form has disappeared again. The NO-responsive form is hyperphosphorylated at this time (Figure 8.3.A, third panel, arrow).

However, upon addition of wortmannin, a specific inhibitor for the PI3K/Akt pathway,




Figure 8.3. Inhibition of PI3K/Akt interferes with phosphorylation of GLO1 during the cell cycle.

A. Phosphorylation of GLO1 with PI3K/Akt inhibitors. Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by re-addition of IL-3. This time point was designated 0 hours. Wortmannin was added after 8 hours. Cells were lysed in CEB after indicated time intervals, and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with an anti-GLO1 antibody are shown. Differential spots are indicated with an arrow or an arrowhead. **B.** Effect of PI3K/AKT inhibitors on *in vitro* GLO1 activity. Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by re-addition of IL-3. This time point was designated 0 hours. Wortmannin was added after 8 hours. Cells were lysed in CEB after indicated time points, lysates were brought to equal concentrations and cell lysates were used to assay GLO1 activity. *In vitro* GLO1 reaction rates were determined using Ultraspec 1100 software (Amersham, BDBiosciences) and are shown relative to the reaction rate of lysates made of exponentially growing cells, which are set as 100%.

strong inhibition of phosphorylation could be observed. 11 hours after re-addition of IL-3 (and three hours after addition of wortmannin), both the phosphorylation on the acidic isoform of GLO1 and the phosphorylation on the α -form were severely impaired (Figure 8.3.A, fourth panel, black arrowhead and white arrowhead respectively). NO-modification still occurred, but the NO-responsive form was phosphorylated to a lesser extent than in the absence of wortmannin (Figure 8.3.A, fourth panel, arrow). When the 2-DE pattern of lysates made 12 hours after initiation of the cell cycle was studied, the hyperphosphorylation of the NO-modified form of GLO1 was reduced in the presence of wortmannin, when compared to the absence of the inhibitor (Figure 8.3.A, fifth panel, arrow). Also the phosphorylation on the more acidic isoform was absent (Figure 8.3.A, fifth panel, arrowhead).

Also the *in vitro* activity of GLO1 was determined. In line with the observations discussed above, no differences were observed in the level of GLO1 activity at the G_1 -S phase transition and the addition of wortmannin, which did show a drastic effect on phosphorylation, did not alter the activity of GLO1 *in vitro* (Figure 8.3.B).

In order to corroborate our findings in the cell cycle, wortmannin was also added to an asynchronous exponentially growing Ba/F3 culture for 3 hours (Figure 8.5.A). Whereas a phosphorylation could be observed under control conditions, possibly caused by the manipulation of the cells, this was not the case when PI3K/Akt was inhibited with wortmannin (Figure 8.5.A, arrowhead). Next, we assayed lysates from asynchronous

Ba/F3 cells that were cultured in the presence or absence of wortmannin. In spite of the differences in phosphorylation between the different conditions, no change in GLO1 activity was induced (Figure 8.5.C).

In summary, our data show that cell cycle dependent phosphorylation of GLO1 is at least in part dependent on the PI3K/Akt pathway. Our data also strengthens the observation that phosphorylation of GLO1 can not be correlated with a difference in *in vitro* GLO1 activity.

Phosphorylation of GLO1 is partially inhibited by rapamycin

Although inhibition of PI3K strongly influences the phosphorylation status of GLO1, phosphorylation of GLO1 still occurs (cf. Figure 8.3.A, fifth panel, where the NO-responsive form is still phosphorylated, albeit less). An important downstream target of PI3K/Akt is mTOR. mTOR and its target p70S6K1 have been shown to be important mediators of the effects of PI3K on cell cycle regulation and proliferation (42,45,60-62). Furthermore, mTOR is considered to be a central regulator in cellular growth and is involved in the coordination of cell growth with cell cycle (63-65).

Ba/F3 cells were synchronized as described above and cells were lysed at the indicated time points after re-addition of IL-3. Analysis of cells lysed after 8 hours show the presence of multiple phosphorylations on both the non-NO-modified form and the




Figure 8.4. Inhibition of mTOR interferes with phosphorylation of GLO1 during the cell cycle.

Phosphorylation of GLO1 with mTOR inhibitors. Ba/F3 cells were synchronized by IL-3 starvation over night. The cell cycle was induced by re-addition of IL-3. This time point was designated 0 hours. Rapamycin was added after 8 hours. Cells were lysed in CEB at indicated time points, and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 µg of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with an anti-GLO1 antibody are shown. Differential spots are indicated with arrows on the NO-modified form and with an arrowhead on the non-NO-modified form.




Figure 8.5. Inhibition of PI3K/Akt or mTOR inhibits phosphorylation of GLO1 in asynchronously growing cells.

A. Phosphorylation of GLO1 with PI3K/Akt inhibitors. Wortmannin was added to exponentially growing Ba/F3 cells. Cells were lysed in CEB after 3 hours and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with an anti-GLO1 antibody are shown. **B.** Phosphorylation of GLO1 with rapamycin. Rapamycin was added to exponentially growing Ba/F3 cells. Cells were lysed in CEB after 3 hours and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with an anti-GLO1 antibody are shown. **C.** *In vitro* GLO1 activity of exponentially growing Ba/F3 cells. Wortmannin or rapamycin was added to exponentially growing Ba/F3 cells. Cells were lysed in CEB after 5 hours, lysates were brought to equal concentrations and cell lysates were used to assay GLO1 activity. Reaction rate of lysates made of exponentially growing cells are set as 100%. Differential spots are indicated with an arrowhead.

NO-modified form (Figure 8.4, upper panel). After 12 hours, the number of different phosphorylations was reduced, with only a phosphorylated NO-modified form, and a phosphorylated non-NO-modified form remaining (Figure 8.4, second panel). After 13 hours, additional phosphorylation can be observed on the non-NO-modified form (Figure 8.4, third panel, arrowhead), as well as on the NO-modified form (Figure 8.4, third panel, arrows). Although the phosphorylations induced at the 13 hour time point, both on the NO-modified form and the non-NO-modified form, can be inhibited with rapamycin (Figure 8.4, fifth panel, arrow and arrowhead respectively), this was not the case after 12 hours. In contrast, at 12 hours, addition of rapamycin seemingly enhanced phosphorylation

on the NO-modified forms (Figure 8.4, fourth panel, arrows). On the other hand, the phosphorylation on the non-NO-modified form was inhibited (Figure 8.4, fourth panel, arrowhead)

Likewise, incubation of asynchronously growing Ba/F3 cells with rapamycin for 3 hours, strongly changed the phosphorylation of GLO1. Whereas multiple phosphorylations could be observed in the absence of rapamycin, these were abolished upon the addition of rapamycin (Figure 8.5.B, second and third panel, arrowheads). As well as in the case of inhibition with wortmannin, no effects on the activity of GLO1 could be found after incubation of Ba/F3 cells in the presence of rapamycin (Figure 8.5.C).

In conclusion, these observations show that phosphorylation of GLO1 in the cell cycle occurs in part in a mTOR dependent manner.

DISCUSSION

GLO1 expression levels and activity have been shown to associate strongly with proliferation. This has been reported to be the case in animal cells (10,11,66) as well as in plants (22,23,25) and yeast (67). In general it is observed that strongly dividing cells and tissues display high GLO1 activity, whereas GLO1 levels and activity drop in more quiescent or matured cells and tissues (11,22,68,69). Furthermore, under various pathophysiological conditions GLO1 is regulated and higher levels of GLO1 are associated with increased proliferation of tumors (14,16-19). The data presented above show that during the cell cycle, GLO1 is strongly subjected to phosphorylation. This is in agreement with the findings described in chapter 6, where was shown how phosphorylation of GLO1 occurred when G_1 arrest was induced by GF withdrawal in a GSK-3 β dependent manner. This took place in several cell lines, suggesting a general mechanism. Of note is the apparent absence of the NO-responsive isoform in early G_1 . It has been described that phosphorylation primarily occurs on the NO-responsive isoforms of GLO1, but both types of posttranslational modification are independent and neither type of modification is required for the other type to occur (48). Phosphorylation could be severely altered by use of wortmannin, a PI3K inhibitor, and rapamycin, an inhibitor for mTOR. This suggests the involvement of the PI3K/Akt/mTOR signaling cascade.

Interestingly, in spite of the strong effects observed with both wortmannin and rapamycin, neither one could however completely inhibit GLO1 phosphorylation. Possibly this is the consequence of feedback mechanisms or compensatory mechanisms that exist in this pathway. Elaborate evidence exists for such mechanisms. Although mTOR activation is downstream of PI3K/Akt, mTOR reciprocally phosphorylates Akt, hereby

either further activating Akt, when mTOR is part of the mTOR complex 2 with rictor (70), or serving as a negative feedback mechanism when mTOR is bound in the raptor containing mTOR complex 1 (71,72). Alternatively, there might be several kinases involved in phosphorylation of GLO1, depending on the stimulus, or acting in a redundant manner. It would also be plausible to hypothesize that inhibition of either Akt or mTOR would result in (partial) activation of GSK-3 β since both kinases are able to phosphorylate and inhibit GSK-3 β *in vivo* (73-77). Moreover, we have shown that GLO1 is a direct substrate for CaMKII (chapter 7).

Cycling cells require a dynamic regulation of glucose metabolism. The presence of glucose is a requisite for cells to proliferate and it has been observed that cells upregulate glucose uptake and glucose metabolism during G₁ and S phase and downregulate glucose uptake during G₂/M phase (31,32,78). GFs play an essential role by promoting cell metabolism and determining the rate of cell division. The PI3K/Akt pathway is critically involved in GF signaling in hematopoietic cells and directly affects the glucose metabolism (34,36,39,79) as well as the cell cycle (42,80-82). It would therefore be reasonable to hypothesize that GFs would attach a dynamic regulation of the GLO system to the cell cycle determined regulation of glycolysis, since a rise in the rate of glycolysis would be expected to automatically implicate higher intracellular levels of the cytotoxic MG. Our data show, however, that different phases in the cell cycle did not display different levels of *in vitro* GLO1 activity (Figure 8.2.A), in contrast with the described rise in G₂/M, which has been described in carrot cell suspension cultures (52). Also at the G₁-S transition, no higher levels of GLO1 were observed *in vitro* (Figure 8.2.B), in spite of the fact that from literature we would expect a rise in glucose uptake and glycolysis (32,83-86). On the other hand, it has been shown that higher GLO1 activity, even when adequate concentrations of reduced glutathione are present, not necessarily suffices to keep intracellular concentrations of MG low (87), which could make a rise in GLO1 activity redundant. Furthermore, addition of rapamycin or wortmannin did not have an effect on GLO1 activity *in vitro*, despite the drastic changes that were induced on phosphorylation. These data should however be interpreted cautiously since they do not exclude the possibility that in the cells GLO1 activity might differ locally, hereby creating microenvironments with different levels of MG. Alternatively GLO1 isoforms have been described that are not involved in detoxification of MG (88), and possibly these isoforms are phosphorylated during the cell cycle.

In summary, our data shows that GLO1 is subject to several phosphorylations during the cell cycle, but this does not seem to influence the activity of GLO1. These phosphorylations are in part mediated by the PI3K/Akt/mTOR signaling cascade.

MATERIALS AND METHODS

Cell Lines and Cultures

The IL-3 dependent pro-B cell line Ba/F3 (89) was purchased from DSMZ. Ba/F3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) with glutamax (Invitrogen) supplemented with heat-inactivated FBS (10% v/v) (Cambrex), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) and 5% conditioned medium from WEHI-3B cells as a source of mouse IL-3. Ba/F3 cells were cultured at 37°C in a humidified incubator under an 5% CO₂ atmosphere.

Cell stimulations

Ba/F3 cells were synchronized by washing exponentially growing cells 3 times in culture medium without conditioned WEHI-3B medium and resuspension in culture medium without conditioned WEHI-3B medium for 8 hours or overnight. At the end of the synchronisation conditioned WEHI-3B medium (5% v/v) was re-added. This time point was designated 0 hours. Alternatively was added for 8 hours or overnight. At the end of the synchronisation nocodazole was removed by washing cells 2 times in fresh culture medium. This time point was designated 0 hours.

Chemicals and inhibitors used are nocodazole (Sigma aaldrich), wortmannin and rapamycin (Calbiochem). Duration and concentrations used as described.

DNA distribution analysis

Cells were stained with propidium iodide (PI, Sigma) containing staining solution (90) and analyzed by a FACSCalibur flow cytometer (488_{Ex}/590_{Em}). Cell Quest software was used to analyze cell cycle distribution (FACS Calibur, Becton Dickinson) (90,91). Ten thousand cells were routinely analyzed.

Electrophoresis and immunoblotting

The cells were washed 3 times with ice-cold PBS buffer and lysed with cytosol extraction buffer (10 mM Tris-HCL pH7.4, 50 mM EDTA pH8.0, 25 mM NaCl, 0,7% TritonX-100, 100 mM PMSF, 1 tablet Complete Protease Inhibitor Cocktail (Roche)/50 ml cytosol extraction buffer). Cell lysates were cleared by centrifugation (14,000 x g). Protein concentrations were determined using Bradford method. Proteins were separated by SDS-PAGE and transferred onto a PVDF membrane (Hybond-P, Amersham Pharmacia Biotech). The blots were incubated with the desired antibodies followed by ECL-based

detection (Amersham Pharmacia Biotech).

2-Dimensional Gel Electrophoresis

Isoelectric focusing was carried out on 18 cm IPG strips, pH 4-7 (GE Healthcare) according to the manufacturer's instructions. For the second dimension, proteins were separated by SDS-polyacrylamide gel electrophoresis (PAGE, 12%).

Determination of GLO1 activity

GLO1 activity was determined as described earlier (47), using the spectrophotometric method developed by Oray and Norton (92). Equal concentrations of cell lysates were used.

Reference List

1. **Dakin, H. D. and H. W. Dudley.** 1913. AN ENZYME CONCERNED WITH THE FORMATION OF HYDROXY ACIDS FROM KETONIC ALDEHYDES. *Journal of Biological Chemistry* **14**:155-157.
2. **Neuberg, C.** The destruction of lactic aldehyde and methylglyoxal by animal organs. *Biochemische Zeitschrift* **49**, 502-506. 1913.
3. **Thornalley, P. J.** 2003. Glyoxalase I - structure, function and a critical role in the enzymatic defence against glycation. *Biochemical Society Transactions* **31**:1343-1348.
4. **Gale, C. P. and P. J. Grant.** 2004. The characterisation and functional analysis of the human glyoxalase-1 gene using methods of bioinformatics. *Gene* **340**:251-260.
5. **Hopkins, F. G. and E. J. Morgan.** 1945. On the distribution of glyoxalase and glutathione. *Biochem.J.* **39**:320-324.
6. **Vanderjagt, D. L., B. Robinson, K. K. Taylor, and L. A. Hunsaker.** 1992. Reduction of Trioses by NADPH-Dependent Aldo-Keto Reductases - Aldose Reductase, Methylglyoxal, and Diabetic Complications. *Journal of Biological Chemistry* **267**:4364-4369.
7. **Egyud, L. G. and A. Szent-Gyorgyi.** 1966. On the regulation of cell division. *Proc.Natl.Acad.Sci.U.S.A* **56**:203-207.
8. **Egyud, L. G. and A. Szent-Gyorgyi.** 1966. Cell division, SH, ketoaldehydes, and cancer. *Proc.Natl.Acad.Sci.U.S.A* **55**:388-393.
9. **Szent-Gyorgyi, A., A. HEGYELI, and J. A. McLaughlin.** 1963. Growth and cellular constituents. *Proc.Natl.Acad.Sci.U.S.A* **49**:878-879.
10. **Amicarelli, F., P. Sacchetta, S. Colafarina, S. Angelucci, M. Miranda, and C. Di Ilio.** 1998. Glyoxalases activity during Bufo bufo embryo development. *Mechanisms of Ageing and Development* **100**:261-267.
11. **Principato, G. B., M. Bodo, M. G. Biagioni, G. Rosi, and F. S. Liotti.** 1982. Glyoxalases and glutathione reductase activity changes in chicken liver during embryo development and after hatching. *Acta Embryol.Morphol.Exp.* **3**:173-179.
12. **Antognelli, C., F. Baldracchini, V. N. Talesa, E. Costantini, A. Zucchi, and E. Mearini.** 2006. Overexpression of glyoxalase system enzymes in human kidney tumor. *Cancer Journal* **12**:222-228.
13. **Ayoub, F., M. Zaman, P. Thornalley, and J. Masters.** 1993. Glyoxalase activities in human tumour cell lines in vitro. *Anticancer Res.* **13**:151-155.
14. **Davidson, S. D., D. M. Milanese, C. Mallouh, M. S. Choudhury, H. Tazaki, and S. Konno.** 2002. A possible regulatory role of glyoxalase I in cell viability of human prostate cancer. *Urol.Res.* **30**:116-121.
15. **Mearini, E., R. Romani, L. Mearini, C. Antognelli, A. Zucchi, T. Baroni, M. Porena, and V. N. Talesa.** 2002. Differing expression of enzymes of the glyoxalase system in superficial and invasive bladder carcinomas. *Eur.J.Cancer* **38**:1946-1950.
16. **Ranganathan, S. and K. D. Tew.** 1993. Analysis of glyoxalase-I from normal and tumor tissue from human colon. *Biochim.Biophys.Acta* **1182**:311-316.
17. **Ranganathan, S., E. S. Walsh, and K. D. Tew.** 1995. Glyoxalase-I in Detoxification - Studies Using A Glyoxalase-I Transfectant Cell-Line. *Biochemical Journal* **309**:127-131.
18. **Rulli, A., L. Carli, R. Romani, T. Baroni, E. Giovannini, G. Rosi, and V. Talesa.** 2001. Expression of glyoxalase I and II in normal and breast cancer tissues. *Breast Cancer Res.Treat.* **66**:67-72.
19. **Sakamoto, H., T. Mashima, S. Sato, Y. Hashimoto, T. Yamori, and T. Tsuruo.** 2001. Selective activation of apoptosis program by S-p-bromobenzylglutathione cyclopentyl diester in glyoxalase I-overexpressing human lung cancer cells. *Clinical Cancer Research* **7**:2513-2518.
20. **Selicharova, I., K. Smutna, M. Sanda, K. Ubik, E. Matouskova, E. Bursikova, M. Brozova, J. Vydra, and J. Jiracek.** 2007. 2-DE analysis of a new human cell line EM-G3 derived from breast cancer progenitor cells and comparison with normal mammary epithelial cells. *Proteomics* **7**:1549-1559.
21. **Di Ilio, C., S. Angelucci, A. Pennelli, A. Zezza, R. Tenaglia, and P. Sacchetta.** 1995. Glyoxalase activities in tumor and non-tumor human urogenital tissues. *Cancer Lett.* **96**:189-193.
22. **Deswal, R., T. N. Chakaravarty, and S. K. Sopory.** 1993. The Glyoxalase System in Higher-Plants - Regulation in Growth and Differentiation. *Biochemical Society Transactions* **21**:527-530.
23. **Paulus, C., B. Kollner, and H. J. Jacobsen.** 1993. Physiological and Biochemical-Characterization of Glyoxalase-I, A General Marker for Cell-Proliferation, from A Soybean Cell-Suspension. *Planta* **189**:561-566.
24. **Ramaswamy, O., S. Pal, S. Guhamukherjee, and S. K. Sopory.** 1984. Correlation of Glyoxalase-I Activity with Cell-Proliferation in Datura-Callus Culture. *Plant Cell Reports* **3**:121-124.
25. **Sethi, U., A. Basu, and S. Guhamukherjee.** 1988. Control of Cell-Proliferation and Differentiation by Regulating Polyamine Biosynthesis in Cultures of Brassica and Its Correlation with Glyoxalase-I Activity. *Plant Science* **56**:167-175.
26. **Chakravarty, T. N. and S. K. Sopory.** 1998. Blue light stimulation of cell proliferation and glyoxalase I activity in callus cultures of *Amaranthus paniculatus*. *Plant Science* **132**:63-69.
27. **Chan, T. O., S. E. Rittenhouse, and P. N. Tsichlis.** 1999. AKT/PKB and other D3 phosphoinositide-regulated kinases: Kinase activation by phosphoinositide-dependent phosphorylation. *Annual Review of Biochemistry* **68**:965-1014.
28. **Duronio, V., M. P. Scheid, and S. Ettinger.** 1998. Downstream signalling events regulated by phosphatidylinositol 3-kinase activity. *Cellular Signalling* **10**:233-239.
29. **Kandel, E. S. and N. Hay.** 1999. The regulation and activities of the multifunctional serine/threonine kinase Akt/PKB. *Experimental Cell Research* **253**:210-229.

30. **Scheid, M. P., R. W. Lauener, and V. Duronio.** 1995. Role of Phosphatidylinositol 8-OH-Kinase Activity in the Inhibition of Apoptosis in Hematopoietic-Cells - Phosphatidylinositol 3-OH-Kinase Inhibitors Reveal A Difference in Signaling Between Interleukin-3 and Granulocyte-Macrophage Colony-Stimulating Factor. *Biochemical Journal* **312**:159-162.
31. **Greiner, E. F., M. Guppy, and K. Brand.** 1994. Glucose Is Essential for Proliferation and the Glycolytic Enzyme-Induction That Provokes A Transition to Glycolytic Energy-Production. *Journal of Biological Chemistry* **269**: 31484-31490.
32. **Kansara, M. and M. V. Berridge.** 2004. Hemopoietic cell transformation is associated with failure to downregulate glucose uptake during the G2/M phase of the cell cycle. *Experimental Cell Research* **293**:321-330.
33. **Hajdуч, E., D. R. Alessi, B. A. Hemmings, and H. S. Hundal.** 1997. Overexpression of wild-type protein kinase B increases glucose transport and amino acid transport in L6 skeletal muscle cells. *Diabetes* **46**:313.
34. **Hajdуч, E., G. J. Litherland, and H. S. Hundal.** 2001. Protein kinase B (PKB/Akt) - a key regulator of glucose transport? *Febs Letters* **492**:199-203.
35. **Plas, D. R., S. Talapatra, A. L. Edinger, J. C. Rathmell, and C. B. Thompson.** 2001. Akt and Bcl-x(L) promote growth factor-independent survival through distinct effects on mitochondrial physiology. *Journal of Biological Chemistry* **276**:12041-12048.
36. **Rathmell, J. C., C. J. Fox, D. R. Plas, P. S. Hammerman, R. M. Cinalli, and C. B. Thompson.** 2003. Akt-directed glucose metabolism can prevent Bax conformation change and promote growth factor-independent survival. *Molecular and Cellular Biology* **23**:7315-7328.
37. **Edinger, A. L. and C. B. Thompson.** 2002. Akt maintains cell size and survival by increasing mTOR-dependent nutrient uptake. *Molecular Biology of the Cell* **13**:2276-2288.
38. **Edinger, A. L.** 2005. Growth factors regulate cell survival by controlling nutrient transporter expression. *Biochemical Society Transactions* **33**:225-227.
39. **Edinger, A. L.** 2007. Controlling cell growth and survival through regulated nutrient transporter expression. *Biochemical Journal* **406**:1-12.
40. **Wiernan, H. L., J. A. Wofford, and J. C. Rathmell.** 2007. Cytokine stimulation promotes glucose uptake via phosphatidylinositol-3 kinase/Akt regulation of Glut1 activity and trafficking. *Molecular Biology of the Cell* **18**: 1437-1446.
41. **Woodland, R. T., C. J. Fox, M. R. Schmidt, P. S. Hammerman, J. T. Opferman, S. J. Korsmeyer, D. M. Hilbert, and C. B. Thompson.** 2008. Multiple signaling pathways promote B lymphocyte stimulator dependent B-cell growth and survival. *Blood* **111**:750-760.
42. **Gao, N., D. C. Flynn, Z. Zhang, X. S. Zhong, V. Walker, K. J. Liu, X. L. Shi, and B. H. Jiang.** 2004. G(1) cell cycle progression and the expression of G(1) cyclins are regulated by PI3K/AKT/mTOR/p70S6K1 signaling in human ovarian cancer cells. *American Journal of Physiology-Cell Physiology* **287**:C281-C291.
43. **Garcia-Morales, P., E. Hernando, E. Carrasco-Garcia, M. P. Menendez-Gutierrez, M. Saceda, and I. Martinez-Lacaci.** 2006. Cyclin D3 is down-regulated by rapamycin in HER-2-overexpressing breast cancer cells. *Molecular Cancer Therapeutics* **5**:2172-2181.
44. **Ikezoe, T., C. Nishioka, K. Bandobashi, Y. Yang, Y. Kuwayama, Y. Adachi, T. Takeuchi, H. P. Koeffler, and H. Taguchi.** 2007. Longitudinal inhibition of PI3K/Akt/mTOR signaling by LY294002 and rapamycin induces growth arrest of adult T-cell leukemia cells. *Leukemia Research* **31**:673-682.
45. **Shah, S. A., M. W. Potter, R. Ricciardi, R. A. Perugini, and M. P. Callery.** 2001. FRAP-p70s6K signaling is required for pancreatic cancer cell proliferation. *Journal of Surgical Research* **97**:123-130.
46. **Kuo, P. L., Y. L. Hsu, and C. Y. Cho.** 2006. Plumbagin induces G(2)-M arrest and autophagy by inhibiting the AKT/mammalian target of rapamycin pathway in breast cancer cells. *Molecular Cancer Therapeutics* **5**:3209-3221.
47. **Van Herreweghe, F., J. Q. Mao, F. W. R. Chaplen, J. Grooten, K. Gevaert, J. Vandekerckhove, and K. Vancompernelle.** 2002. Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methyl glyoxal-derived AGE. *Proceedings of the National Academy of Sciences of the United States of America* **99**:949-954.
48. **de, H., V. D. Rondas, J. Vandekerckhove, and K. Vancompernelle.** 2007. Tumour necrosis factor induces phosphorylation primarily of the nitric-oxide-responsive form of glyoxalase I. *Biochem.J.* **407**:121-128.
49. **Khan, M. M. K., A. Jan, H. Karibe, and S. Komatsu.** 2005. Identification of phosphoproteins regulated by gibberellin in rice leaf sheath. *Plant Molecular Biology* **58**:27-40.
50. **Shin, R., S. Alvarez, A. Y. Burch, J. M. Jez, and D. P. Schachtman.** 2007. Phosphoproteomic identification of targets of the Arabidopsis sucrose nonfermenting-like kinase SnRK2.8 reveals a connection to metabolic processes. *Proceedings of the National Academy of Sciences of the United States of America* **104**:6460-6465.
51. **Usui, Y., M. Nakase, H. Hotta, A. Urisu, N. Aoki, K. Kitajima, and T. Matsuda.** 2001. A 33-kDa allergen from rice (*Oryza sativa* L. Japonica) - cDNA cloning, expression, and identification as a novel glyoxalase I. *Journal of Biological Chemistry* **276**:11376-11381.
52. **Ghosh, S., J. Sen, S. Kalia, and S. Guha-Mukherjee.** 1999. Establishment of synchronization in carrot cell suspension culture and studies on stage specific activation of glyoxalase I. *Methods Cell Sci.* **21**:141-148.
53. **Blalock, W. L., C. Weinstein-Oppenheimer, F. Chang, P. E. Hoyle, X. Y. Wang, P. A. Algate, R. A. Franklin, S. M. Oberhaus, L. S. Steelman, and J. A. McCubrey.** 1999. Signal transduction, cell cycle regulatory, and anti-apoptotic pathways regulated by IL-3 in hematopoietic cells: possible sites for intervention with anti-neoplastic drugs. *Leukemia* **13**:1109-1166.
54. **de Groot, R. P., P. J. Coffer, and L. Koenderman.** 1998. Regulation of proliferation, differentiation and survival

- by the IL-3/IL-5/GM-CSF receptor family. *Cellular Signalling* **10**:619-628.
55. **Shelton, J. G., P. W. Moye, L. S. Steelman, W. L. Blalock, J. T. Lee, R. A. Franklin, M. McMahon, and J. A. McCubrey.** 2003. Differential effects of kinase cascade inhibitors on neoplastic and cytokine-mediated cell proliferation. *Leukemia* **17**:1765-1782.
 56. **Baixeras, E., S. Jeay, P. A. Kelly, and M. C. Postel-Vinay.** 2001. The proliferative and antiapoptotic actions of growth hormone and insulin-like growth factor-1 are mediated through distinct signaling pathways in the pro-B Ba/F3 cell line. *Endocrinology* **142**:2968-2977.
 57. **Craddock, B. L., E. A. Orchiston, H. J. Hinton, and M. J. Welham.** 1999. Dissociation of apoptosis from proliferation, protein kinase B activation, and BAD phosphorylation in interleukin-3-mediated phosphoinositide 3-kinase signaling. *Journal of Biological Chemistry* **274**:10633-10640.
 58. **Gottlob, K., N. Majewski, S. Kennedy, E. Kandel, R. B. Robey, and N. Hay.** 2001. Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. *Genes & Development* **15**:1406-1418.
 59. **Jeay, S., G. E. Sonenshein, P. A. Kelly, M. C. Postel-Vinay, and E. Baixeras.** 2001. Growth hormone exerts antiapoptotic and proliferative effects through two different pathways involving nuclear factor-kappa B and phosphatidylinositol 3-kinase. *Endocrinology* **142**:147-156.
 60. **Martin, K. A. and J. Blenis.** 2002. Coordinate regulation of translation by the PI 3-kinase and mTOR pathways. *Advances in Cancer Research*, Vol 86 **86**:1-39.
 61. **Gera, J. F., I. K. Mellingshoff, Y. J. Shi, M. B. Rettig, C. Tran, J. H. Hsu, C. L. Sawyers, and A. K. Lichtenstein.** 2004. AKT activity determines sensitivity to mammalian target of rapamycin (mTOR) inhibitors by regulating cyclin D1 and c-myc expression. *Journal of Biological Chemistry* **279**:2737-2746.
 62. **Shi, Y. J., H. J. Yan, P. Frost, J. Gera, and A. Lichtenstein.** 2005. Mammalian target of rapamycin inhibitors activate the AKT kinase in multiple myeloma cells by up-regulating the insulin-like growth factor receptor/insulin receptor substrate-1/phosphatidylinositol 3-kinase cascade. *Molecular Cancer Therapeutics* **4**:1533-1540.
 63. **Coqueret, O.** 2002. Linking cyclins to transcriptional control. *Gene* **299**:35-55.
 64. **Jorgensen, P. and M. Tyers.** 2004. How cells coordinate growth and division. *Current Biology* **14**:R1014-R1027.
 65. **Schmelzle, T. and M. N. Hall.** 2000. TOR, a central controller of cell growth. *Cell* **103**:253-262.
 66. **Principato, G. B., P. Locci, G. Rosi, V. Talesa, and E. Giovannini.** 1983. Activity Changes of Glyoxalase-I Glyoxalase-II and Glutathione-Reductase in Regenerating Rat-Liver. *Biochemistry International* **6**:249-255.
 67. **Dudani, A. K., L. K. Srivastava, and R. Prasad.** 1984. Glyoxalase-I Activity and Cell-Cycle Regulation in Yeast. *Biochemical and Biophysical Research Communications* **119**:962-967.
 68. **McLellan, A. C. and P. J. Thornalley.** 1989. Glyoxalase Activity in Human Red Blood-Cells Fractioned by Age. *Mechanisms of Ageing and Development* **48**:63-71.
 69. **Haik, G. M., T. W. C. Lo, and P. J. Thornalley.** 1994. Methylglyoxal Concentration and Glyoxalase Activities in the Human Lens. *Experimental Eye Research* **59**:497-500.
 70. **Sarbasov, D. D., S. M. Ali, S. Sengupta, J. H. Sheen, P. P. Hsu, A. F. Bagley, A. L. Markhard, and D. M. Sabatini.** 2006. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Molecular Cell* **22**:159-168.
 71. **Findlay, G. M., L. S. Harrington, and R. F. Lamb.** 2005. TSC1-2 tumour suppressor and regulation of mTOR signalling: linking cell growth and proliferation? *Current Opinion in Genetics & Development* **15**:69-76.
 72. **Harrington, L. S., G. M. Findlay, and R. F. Lamb.** 2005. Restraining P13K: mTOR signalling goes back to the membrane. *Trends in Biochemical Sciences* **30**:35-42.
 73. **Cross, D. A. E., D. R. Alessi, P. Cohen, M. Andjelkovich, and B. A. Hemmings.** 1995. Inhibition of Glycogen-Synthase Kinase-3 by Insulin-Mediated by Protein-Kinase-B. *Nature* **378**:785-789.
 74. **Peyrollier, K., E. Hajdich, A. S. Blair, R. Hyde, and H. S. Hundal.** 2000. L-leucine availability regulates phosphatidylinositol 3-kinase, p70 S6 kinase and glycogen synthase kinase-3 activity in L6 muscle cells: evidence for the involvement of the mammalian target of rapamycin (mTOR) pathway in the L-leucine-induced up-regulation of System A amino acid transport. *Biochemical Journal* **350**:361-368.
 75. **Sutherland, C., I. A. Leighton, and P. Cohen.** 1993. Inactivation of Glycogen-Synthase Kinase-3-Beta by Phosphorylation - New Kinase Connections in Insulin and Growth-Factor Signaling. *Biochemical Journal* **296**:15-19.
 76. **Sutherland, C. and P. Cohen.** 1994. The Alpha-Isoform of Glycogen-Synthase Kinase-3 from Rabbit Skeletal-Muscle Is Inactivated by P70 S6 Kinase Or Map Kinase-Activated Protein Kinase-1 In-Vitro. *Febs Letters* **338**:37-42.
 77. **Zhang, H. H., A. I. Lipovsky, C. C. Dibble, M. Sahin, and B. D. Manning.** 2006. S6K1 regulates GSK3 under conditions of mTOR-dependent feedback inhibition of Akt. *Molecular Cell* **24**:185-197.
 78. **Ishii, T., M. Fujishiro, M. Masuda, K. Okudela, H. Kitamura, S. Teramoto, and T. Matsuse.** 2004. Nutritional deficiency affects cell cycle status and viability in A549 cells: role of p27Kip1. *Cancer Letters* **213**:99-109.
 79. **Barata, J. T., A. Silva, J. G. Brandao, L. M. Nadler, A. A. Cardoso, and V. A. Boussiotis.** 2004. Activation of P13K is indispensable for interleukin 7-mediated viability, proliferation, glucose use, and growth of T cell acute lymphoblastic leukemia cells. *Journal of Experimental Medicine* **200**:659-669.
 80. **Grewe, M., F. Gansauge, R. M. Schmid, G. Adler, and T. Seufferlein.** 1999. Regulation of cell growth and cyclin D1 expression by the constitutively active FRAP-p70(s6K) pathway in human pancreatic cancer cells. *Cancer Research* **59**:3581-3587.
 81. **Muise-Helmericks, R. C., H. L. Grimes, A. Bellacosa, S. E. Malstrom, P. N. Tsichlis, and N. Rosen.** 1998. Cyclin D expression is controlled post-transcriptionally via a phosphatidylinositol 3-kinase Akt-dependent

- pathway. *Journal of Biological Chemistry* **273**:29864-29872.
82. **Rossig, L., C. Badorff, Y. Holzmann, A. M. Zeiher, and S. Dimmeler.** 2002. Glycogen synthase kinase-3 couples AKT-dependent signaling to the regulation of p21(Cip1) degradation. *Journal of Biological Chemistry* **277**:9684-9689.
83. **Bushart, G. B., U. Vetter, and W. Hartmann.** 1993. Glucose-Transport During Cell-Cycle in Im9 Lymphocytes. *Hormone and Metabolic Research* **25**:210-213.
84. **Hedekov, C. J.** 1968. Early Effects of Phytohaemagglutinin on Glucose Metabolism of Normal Human Lymphocytes. *Biochemical Journal* **110**:373-&.
85. **Plagemenn, P. G., D. P. Richey, J. M. Zylka, and J. Erbe.** 1975. Cell cycle and growth stage-dependent changes in the transport of nucleosides, hypoxanthine, choline, and deoxyglucose in cultured Novikoff rat hepatoma cells. *J. Cell Biol.* **64**:29-41.
86. **Roos, D. and J. A. Loos.** 1970. Changes in Carbohydrate Metabolism of Mitogenically Stimulated Human Peripheral Lymphocytes .1. Stimulation by Phytohaemagglutinin. *Biochimica et Biophysica Acta* **222**:565-&.
87. **Staniszewska, M. M. and R. H. Nagaraj.** 2006. Upregulation of glyoxalase I fails to normalize methylglyoxal levels: A possible mechanism for biochemical changes in diabetic mouse lenses. *Molecular and Cellular Biochemistry* **288**:29-36.
88. **Larsen, K., A. C. Aronsson, E. Marmstal, and B. Mannervik.** 1985. Immunological Comparison of Glyoxalase-I from Yeast and Mammals and Quantitative-Determination of the Enzyme in Human-Tissues by Radioimmunoassay. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* **82**:625-638.
89. **Palacios, R. and M. Steinmetz.** 1985. II3-Dependent Mouse Clones That Express B-220 Surface-Antigen, Contain Ig Genes in Germ-Line Configuration, and Generate Lymphocytes-B Invivo. *Cell* **41**:727-734.
90. **Krishan, A.** 1990. Rapid DNA content analysis by the propidium iodide-hypotonic citrate method. *Methods Cell Biol.* **33**:121-125.
91. **Nicoletti, I., G. Migliorati, M. C. Pagliacci, F. Grignani, and C. Riccardi.** 1991. A Rapid and Simple Method for Measuring Thymocyte Apoptosis by Propidium Iodide Staining and Flow-Cytometry. *Journal of Immunological Methods* **139**:271-279.
92. **Oray, B. and S. J. Norton.** 1982. Glyoxalase-I from Mouse-Liver. *Methods in Enzymology* **90**:542-546.

A.

```

Homo sapiens GLO1 1 MAEPQPSGGLTDEAALSCCSDADPSTKDFLLQQTMLRVKDPKKSLDFYTRVL 53
                        +                *
Mus musculus GLO1 1 MAEPQPASSGLTDETAFSCCSDPDPSTKDFLLQQTMLRIKDPKKSLDFYTRVL 53
                        +                *

Homo sapiens GLO1 54 GMTLIQKCDFIMKFSLYFLAYEDKNDIPKEKDEKIAWALSRKATLELTHNWG 106
                                                *
                                                *
                                                *
Mus musculus GLO1 54 GLTLLQKLDFPAMKFSLYFLAYEDKNDIPKDKSEKTAWTFSRKATLELTHNWG 106
                                                *
                                                *
                                                *
                                                *

Homo sapiens GLO1 107 TEDDETQSYHNGNSDPRGFGHIGIAVPDVYSACKRFEELGVKFVKKPDDGKMK 159
Mus musculus GLO1 107 TEDDETQSYHNGNSDPRGFGHIGIAVPDVYSACKRFEELGVKFVKKPDDGKMK 159

Homo sapiens GLO1 160 GLAFIQDPDGYWIEILNPNKMATLM 185
Mus musculus GLO1 160 GLAFIQDPDGYWIEILNPNKIATII 185

```

B.

Phosphorylation sites predicted:

Ser: 4 Thr: 2

Serine predictions				
Name	Pos	Context	Score	Pred
v				
Sequence	8	PQPPSGGLT	0.664	*S*
Sequence	18	EAALSCCSD	0.441	.
Sequence	21	LSCCSDADP	0.329	.
Sequence	26	DADPSTKDF	0.997	*S*
Sequence	45	DPKKS L DFY	0.645	*S*
Sequence	69	IMKFSLYFL	0.032	.
Sequence	94	AWAL S RKAT	0.980	*S*
Sequence	114	DETQSYHNG	0.431	.
Sequence	120	HNGNSDPRG	0.193	.
Sequence	137	PDVYSACKR	0.009	.
^				

Threonine predictions				
Name	Pos	Context	Score	Pred
v				
Sequence	12	SGGLTDEAA	0.152	.
Sequence	27	ADPSTKDFL	0.033	.
Sequence	35	LLQQTMLRV	0.053	.
Sequence	50	LDFYTRVLG	0.013	.
Sequence	56	VLGMTLIQK	0.041	.
Sequence	98	SRKATLELT	0.726	*T*
Sequence	102	TLELTHNWG	0.041	.
Sequence	107	HNWGTEDDE	0.768	*T*
Sequence	112	EDDETQSYH	0.013	.
Sequence	182	NKMATLM--	0.041	.
^				

C.

Phosphorylation sites predicted:

Ser: 6 Thr: 2

Serine predictions				
Name	Pos	Context	Score	Pred
v				
Sequence	8	PQPASSGLT	0.482	.
Sequence	9	QPASSGLTD	0.640	*S*
Sequence	18	ETAFSCCSD	0.695	*S*
Sequence	21	FSCCSDPDP	0.267	.
Sequence	26	DPDPSTKDF	0.997	*S*
Sequence	45	DPKKS L DFY	0.645	*S*
Sequence	69	AMKFSLYFL	0.023	.
Sequence	86	PKDKSEKTA	0.986	*S*
Sequence	94	AWTF S RKAT	0.963	*S*
Sequence	114	DETQSYHNG	0.431	.
Sequence	120	HNGNSDPRG	0.193	.
Sequence	137	PDVYSACKR	0.009	.
Sequence	201	PDVYSACKR	0.009	.
^				

Threonine predictions				
Name	Pos	Context	Score	Pred
v				
Sequence	12	SSGLTDETA	0.054	.
Sequence	15	LTDETA F SC	0.046	.
Sequence	27	PDPSTKDFL	0.092	.
Sequence	35	LLQQTMLRI	0.013	.
Sequence	50	LDFYTRVLG	0.013	.
Sequence	56	VLGLTLLQK	0.025	.
Sequence	89	KSEKTAW T F	0.132	.
Sequence	92	KTAWTF S RK	0.089	.
Sequence	98	SRKATLELT	0.726	*T*
Sequence	102	TLELTHNWG	0.041	.
Sequence	107	HNWGTEDDE	0.768	*T*
Sequence	112	EDDETQSYH	0.013	.
Sequence	182	NKIATIIDP	0.286	.
Sequence	246	NKMATLM--	0.041	.
^				

Supplementary figure 1. Prediction of phosphorylation sites in glyoxalase I.

A. Alignment of the amino acid sequence of human and mouse GLO1 as found in UniProtKB/Swiss-Prot. Potential phosphorylation sites as predicted by Netphos software are indicated in bold. The potential phosphorylation sites that were described for human GLO1 in (1) are indicated within boxes. Predicted phosphorylation sites for CaMKII (*), GSK-3 (+), PKA (✱) and Akt (✕) were determined using Motifscan software (low stringency).

B and C. Scores of predicted phosphorylation sites as obtained using Netphos software for human (**A**) and mouse (**B**) GLO1.

1. Ranganathan, S., E. S. Walsh, A. K. Godwin, and K. D. Tew. 1993. Cloning and Characterization of Human Colon Glyoxalase-I. *Journal of Biological Chemistry* **268**:5661-5667.

Overview of known phosphorylations of GLO1.

mammalian cell lines	Stimulus	Kinase	Cell line	In vitro		Reference	Of note
				phosphorylation	Change in GLO1 activity		
	TNF	PKA	L929	no	no	(1)	established using specific inhibitors
			L929, Hek293	yes	/	de Hemptinne, unpublished results; chapter 7 of this thesis	established using specific inhibitors confirmed with phospho-specific antibody
		CaMKII	BaF3	/	no	chapter 8 of this thesis	established using specific inhibitors
	cell cycle	PI3K/Akt	BaF3	/	no	chapter 8 of this thesis	established using specific inhibitors
		mTOR	BaF3	/	no	chapter 8 of this thesis	established using specific inhibitors
	IL-3 starvation	GSK-3 β	BaF3	no	no	chapter 6 and chapter 7 of this thesis	established using specific inhibitors confirmed using coexpression in HEK293
	Serum starvation	/	HL-60	/	/	chapter 6 of this thesis	
		/	NIH3T3	/	/	chapter 6 of this thesis	
	addition of mating factor	S. cerevisiae		/	+	(2,3)	
	addition of gibberelin	CDPK	rice leaf	/	/	(4)	
yeast	overexpression of SnRK2.8	SnRK2.8	Arabidopsis	yes	+	(5)	shares its consensus sequence with CaMKII SnRK family are plant homologues of AMPK
plants							

Reference List

1. Van Herreweghe, F., J. Q. Mao, F. W. R. Chaplen, J. Grooten, K. Gevaert, J. Vandekerckhove, and K. Vancompermolle. 2002. Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methyl glyoxal-derived AGE. Proceedings of the National Academy of Sciences of the United States of America **99**:949-954.
2. Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura. 1989. Sexual-Response in Saccharomyces-Cerevisiae - Alteration of Enzyme-Activity in the Glyoxalase System by Mating Factor. Biochemical and Biophysical Research Communications **165**:1091-1095.
3. Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura. 1990. Sexual-Response of Saccharomyces-Cerevisiae - Phosphorylation of Yeast Glyoxalase-I by A Cell Extract of Mating Factor Treated-Cells. Journal of Biochemistry **108**:4-6.
4. Khan, M. M. K., A. Jan, H. Karibe, and S. Komatsu. 2005. Identification of phosphoproteins regulated by gibberellin in rice leaf sheath. Plant Molecular Biology **58**:27-40.
5. Shin, R., S. Alvarez, A. Y. Burch, J. M. Jez, and D. P. Schachtman. 2007. Phosphoproteomic identification of targets of the Arabidopsis sucrose nonfermenting-like kinase SnRK2.8 reveals a connection to metabolic processes. Proceedings of the National Academy of Sciences of the United States of America **104**:6460-6465.

Part II:

Experimental Data and Results

Chapter 9:

The Receptor for Advanced Glycation Endproducts (RAGE) is regulated at several levels during the cell cycle

A novel proteolytically generated form of
RAGE

Chapter 9: The Receptor for Advanced Glycation Endproducts (RAGE) is regulated at several levels during the cell cycle

A novel proteolytically generated form of RAGE

INTRODUCTION

The receptor for advanced glycation endproducts (RAGE) was first isolated as a receptor for advanced glycation endproducts (AGEs), which are adducts formed by glycation and subsequent oxidation of proteins (1,2). RAGE is a member of the immunoglobulin superfamily of receptors and has been suggested to be involved in embryonic development, since it has been reported to be expressed during development, repressed in adulthood and re-expressed in several pathologies (3). Although the original ligand to be identified for RAGE were AGEs, a remarkable feature of RAGE is its ability to be activated by different other specific ligands such as amphotericin (often referred to as high mobility group box I (HMGB1) DNA-binding protein) (4), several members of the S100 Ca²⁺-binding protein family (5,6), and β -sheet fibrils and amyloid- β -peptide (7,8). Multifarious signaling pathways are induced upon RAGE activation, including the reactive oxygen species (ROS)-dependent activation of mitogen-activated protein kinases (MAPKs) (9-12), other MAPK pathways, such as the p38 pathway and the stress-activated protein/c-Jun N-terminal kinase (SAPK/JNK) (11,13,14), Janus kinase/signal transducer and activator of transcription (JAK/STAT) pathways (15,16), the small GTPases cdc42 and rac (17) and the phosphoinositol-3-kinase (PI3K) pathway (18,19). To date, the full range of different signaling pathways initiated by RAGE activation has not been elucidated. Also the intracellular factors that bind to the cytoplasmic tail of RAGE remain to be determined. The outcome of RAGE signaling is highly dependent on the cell type, the available ligands and their concentrations, and the different levels and isoforms of RAGE, all of this adding further to the complexity of RAGE signaling.

RAGE activation has been implicated in cell cycle related signaling. Although proliferation and differentiation are sequential and mutually exclusive events, RAGE has been involved in both, depending on the ligand and cell type. RAGE activation is often considered to promote tumor growth and metastasis, and the level of RAGE expression is regarded as a measure for metastatic potential of tumor cells (13,20-29). In neuronal cells it was reported that both AGEs as well as S100B could elicit a mitogenic response through RAGE, hereby upregulating cyclin D₁ expression and urging cells into S phase in a p42/44 MAPK dependent manner (30,31). Also in other cell types, such as fibroblasts (29), osteoblasts (9), smooth muscle cells (26) and mesangial cells (23), activation of RAGE induced MAPK activation and subsequent cell growth.

However, in spite of the elaborate evidence for a role of RAGE in proliferation, this has been contradicted by several reports. In melanoma and neuroblastoma cells RAGE activation by amphotericin led to inhibition of metastasis *in vivo* (32). Further it has been reported that overexpression of full-length RAGE (FLRAGE) did not promote metastasis of lung cancer cells *in vivo*, but instead showed a diminished proliferation *in vitro* (33,34). It was also shown that higher tumor stages and several other metastatic tissues displayed a downregulation of RAGE (35). The same was observed after overexpression of RAGE in rhabdomyosarcoma cells, where binding of amphotericin to RAGE stimulated differentiation in a cdc42 and p38 dependent manner (36). In the development of diabetic nephropathy it was observed that AGE interaction with RAGE activated STAT5-mediated upregulation of p21^{waf} and consequent growth arrest (37). Furthermore, both amphotericin and AGE-BSA are able to stimulate maturation of dendritic cells (38,39).

The ambiguous function of RAGE in the regulation of cell cycle may find its basis partly in the presence of different isoforms of RAGE. RAGE signaling is not mediated by the action of a single type of RAGE receptor, but relies on the interplay of different factors, such as the level of RAGE expression, the presence and relative concentrations of different ligands, the large number of different isoforms and their possible splice variants, and possible posttranslational modifications of RAGE (reviewed in (40,41)). Furthermore, the outcome of RAGE signaling is, above all, highly dependent on the cell type. We commenced the study for the involvement of RAGE in the regulation of the cell cycle by studying the differential expression of RAGE isoforms. A twofold approach was used. Cells were synchronized and the different phases and phase transitions were studied. Alternatively, G₁ arrest was induced in Ba/F3 cells or Hek293 cells by IL-3 depletion or serum starvation respectively.

RESULTS

Ba/F3 cells exhibit different isoforms of RAGE

First, we studied the expression of the different isoforms of RAGE. Several different isoforms have already been described, the major isoforms being FLRAGE, the soluble RAGE (sRAGE) which lacks the transmembrane domain and the cytoplasmic domain and the dominant negative RAGE (DNRAGE) which lacks the cytoplasmic domain (reviewed in (40)). Several other isoforms have also been described, but their expression is cell type dependent (41,42). The antibody used was an antibody raised against the C-terminus of RAGE (C-t-Ab; Abcam). Soluble isoforms of RAGE would therefore be missed in our study since they have alternative C-termini (43,44). However, the specificity of this antibody was confirmed by the specific recognition of RAGE on Western blots from wild type mouse lung




Figure 9.1. RAGE isoforms in Ba/F3 cells.

Isoforms of RAGE in Ba/F3 cells. Exponentially growing Ba/F3 cells were lysed in CEB. 40 µg of protein was separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-terminus-directed anti-RAGE antibody (C-t-Ab; Abcam).

extracts, whereas RAGE was not detected in lung extracts from RAGE^{-/-} mice (Salliau et al., submitted).

In total lysates from asynchronously growing Ba/F3 cells several isoforms are present (Figure 9.1). The band migrating at 50 kDa is ^{FL}RAGE (44,45). ^{FL}RAGE can be severely glycosylated, which is known to result in bands of higher molecular weight, such as the 52 kDa, the 55 kDa and the two bands near the 75 kDa marker (44,46). Further we observed the presence of several high molecular weight bands. These high molecular weight bands are hypothesized to result from oligomerization or complexes with other proteins (43,47). Also a band of 42 kDa is present in Ba/F3 cells. This band has been described as a N-terminal truncated form of RAGE (44). Further bands of 37 kDa and 27 kDa can be observed, which are hypothesized to be processed and truncated forms of RAGE ((43); Salliau et al., submitted). Furthermore, a band of approximately 17 kDa can be seen. This is a novel proteolytic processed isoform and will be discussed below.

It should be noted that RAGE not only appears on the plasmamembrane. Some isoforms, among which ^{FL}RAGE, also appear in the cytosol and the nucleus (Salliau et al., submitted).

RAGE expression is altered during the cell cycle

We then analysed whether different phases are associated with differences in the expressed RAGE isoforms. G₁ arrest was induced by IL-3 depletion and G₂/M arrest by addition of nocodazole. The analysis of DNA distribution on FACS confirmed the respective phases arrests (Figure 9.2.B). Lysates of cells arrested in either G₁ or G₂/M were compared with lysates of asynchronously growing cells. Nocodazole-treated cells showed an upregulation of the expression of a band with an apparent molecular weight of 42 kDa (Figure 9.2.A lane 2, indicated with an arrowhead). This band corresponds to the band that has been described to be a N-terminal truncated isoform, which lacks the N-terminal V-




Figure 9.2. Cell cycle phase specific RAGE expression.

A. Analysis of RAGE in different phases. G_1 arrest was induced by IL-3 starvation of Ba/F3 cells and G_2/M arrest was induced by addition of nocodazole. Cells were lysed in CEB, equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-t-Ab. **B.** Analysis of the cell cycle. Induction of the different phase arrests were monitored by analysis of the DNA content by flow cytometry.

type immunoglobulin-like domain (43,44). This upregulation was not observed in the cells that were arrested in the G_1 phase (Figure 9.2.A, lane 3). Also the high molecular weight isoforms displayed different levels of expression when the G_1 and G_2/M arrested cells were compared with asynchronously growing cells. These high molecular weight bands were specific for the cytosolic fraction and could not be observed in nuclear fractions (data not shown). In the lysate of the asynchronously growing culture, two intense protein bands are observed between with molecular weights of approximately 80 kDa and 100 kDa, with in between them a less intense band of approximately 90 kDa (Figure 9.2.A, indicated with arrows). The 80 kDa band was downregulated in the G_2/M -arrested cells, whereas in the G_1 -arrested cells the inverse was observed, and the 90 kDa band and the 100 kDa band were expressed substantially less.

In this context, we additionally studied whether the change of the expressed




Figure 9.3. Differential expression of RAGE at phase transitions.

A. Isoforms of RAGE at G₁-S transition. **B.** Isoforms of RAGE at S-G₂/M transition. Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by readdition of IL-3. This time point was designated 0 hours. Cells were lysed in CEB at indicated time points and equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-t-Ab. Blots were subsequently stripped and reprobed with an anti-cyclin E antibody (Abcam) (**A**) or an anti-cyclin A antibody (Santa Cruz) (**B**). Protein band intensities were measured using Genetools software (Westburg) and corrected for equal concentrations. The relative intensities of the 100 kDa isoform and the 80 kDa isoform are represented by black and white bars respectively in (**A**) and the 90 kDa isoform and the 80 kDa isoform are represented by black and white bars respectively in (**B**).

isoforms coincided with the transition of different phases. Hence, cells were synchronized by IL-3 withdrawal and lysates were made after the appropriate time intervals after re-addition of IL-3 to release cell cycle arrest (indicated by hours after re-addition of IL-3). The transgression of the cells through the different phases of the cell cycle was observed by analysis of the DNA distribution on FACS and the transition of G_1 to S was confirmed by the rise of the level of cyclin E (Figure 9.3.A, lower panel indicated with an arrow), a marker for late G_1 and S phase. Immunoblotting of the obtained lysates endorsed the observation that differential expression of RAGE isoforms coincides with specified phases (Figure 9.3). When the progression through G_1 phase towards S phase was studied (Figure 9.3.A), we observed a strong decline of the 80 kDa isoform of RAGE throughout G_1 , reaching its lowest level at 7 hours, where it only retains 15 % of its initial intensity (Figure 9.3.A, upper panel indicated with an arrow and chart, white bars). 9 hours after the re-addition of IL-3 to remove the cell cycle block, this 80 kDa isoform is then upregulated to approximately 40 % of its initial level. This time corresponds to the beginning of the S phase. Concomitantly with the downregulation of 80 kDa band, the 100 kDa isoform of RAGE (Figure 9.3.A) is upregulated throughout G_1 , reaching its maximum level at the end of G_1 , when expression levels of the 80 kDa isoform is lowest, and its expression level slightly decreases after the onset of S, 9 hours after the release of the cell cycle block (Figure 9.3.A, upper panel indicated with an arrowhead and chart, black bars).

Also at the transition of the S to the G_2 phase was studied. While 100 kDa RAGE isoform remained constant throughout this study, a rapid decline was observed in the




Figure 9.4. Cell surface staining for RAGE expression.

Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by readdition of IL-3. This time point was designated 0 hours. Staining occurred at indicated timepoints using a N-terminus-directed anti-RAGE antibody (Santa Cruz). Gray filled histograms represent isotype control and black histograms represent RAGE expression.

90 kDa band, reaching its lowest levels of approximately 20 % of the initial intensity 16 or 17 hours after the release of the cell cycle arrest. This time point coincides with the transgression from the S phase to the G₂ phase (Figure 9.3.B, upper panel indicated with an arrowhead and chart, black bars). The 80 kDa RAGE isoform also decreased during S phase, and reached minimum levels after 19 hours in late G₂ phase (Figure 9.3.B, upper panel indicated with an arrow and chart, white bars). S to G₂ transition was marked by the decrease of the level of cyclin A, a marker for S phase (Figure 9.3.B, lower panel indicated with an arrow).

Subsequently, we measured the membrane expression levels of RAGE on FACS. In spite of the strong alterations in high molecular weight isoforms of RAGE during the cell cycle, we could not correlate these observations to differences in the membrane expression levels (Figure 9.4).

In summary, we show that expression of RAGE is strongly regulated throughout the cell cycle. This regulation occurs mainly intracellularly and is not related to different levels of localization of RAGE on the plasmamembrane.

RAGE is processed into a 17 kDa isoform during G₁ arrest

As shown in Figure 9.1 we detected a novel isoform of RAGE of approximately 17 kDa in asynchronously growing Ba/F3 cells. This isoform, which will be designated 17 kDa RAGE, has not been reported to date. Cellular fractionation revealed this isoform was present predominantly in the nuclear and membrane fractions, and could be distinguished much less in cytosolic fractions (data not shown).

To examine whether this novel isoform could be connected to a particular phase, we compared lysates of asynchronously growing Ba/F3 cells with lysates from Ba/F3 cells that were arrested in G₁ by IL-3 depletion and nocodazole-induced G₂/M arrested Ba/F3 cells. As shown in Figure 9.5.A, a strong induction of 17 kDa RAGE was observed specifically in the G₁ arrested cells (Figure 9.5.A, arrow). This induction was specific, since no upregulation of FLRAGE was observed (Figure 9.5.A, arrowhead). The induction of G₂/M arrest and G₁ arrest by nocodazole and IL-3 depletion respectively were confirmed by analysis of the DNA-distribution on FACS (Figure 9.5.B). Subsequently Ba/F3 cells were synchronized by IL-3 depletion and re-addition of IL-3 and RAGE expression was studied throughout the cell cycle. The 17 kDa RAGE isoform could also be detected in the S phase (Figure 9.5.C), where it seemed to be up- and downregulated in an alternating fashion (Figure 9.5.C, arrow), in contrast to FLRAGE, which appears to remain constant (Figure 9.5.C, arrowhead).

Together we show the existence of a novel isoform of RAGE, which is strongly induced during G_1 arrest, although it also occurs during the cell cycle.

Kinetics of 17 kDa RAGE generation

Since prolonged absence of IL-3 also leads to the induction of apoptosis in Ba/F3




Figure 9.5. Novel 17 kDa isoform of RAGE.

A. Analysis of 17 kDa RAGE in different phases. G_1 arrest was induced by IL-3 starvation of Ba/F3 cells and G_2/M arrest was induced by addition of nocodazole. Cells were lysed in CEB, equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-t-Ab. **B** Analysis of the cell cycle. Induction of the different phase arrests were monitored by analysis of the DNA content by flow cytometry. **C.** 17 kDa RAGE at G_1 -S transition. Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by re-addition of IL-3. This time point was designated 0 hours. Cells were lysed in CEB at indicated time points equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-t-Ab.

cells we next studied the kinetics of 17 kDa induction in Ba/F3 cells. In the experiment shown, induction of 17 kDa RAGE was detected after 6 hours of IL-3 depletion (Figure 9.6), well before the onset of apoptosis and further increased at later time points.

We were also interested in whether addition of exogenous methylglyoxal (MG) could influence the formation of 17 kDa RAGE. MG is a very potent glycating agent of proteins and the most important precursor of advanced glycation endproducts (AGEs) (48,49). RAGE is known to be upregulated in the presence of its ligands, providing a positive feedback loop when activated (50,51). Moreover, it has been shown that exogenous MG is strongly synergistic with the depletion of IL-3 in the induction of apoptosis in Ba/F3 cells (Audiffret et al., submitted, see also chapter 6). IL-3 depletion-induced generation of 17 kDa RAGE was already observed after 3 hours upon addition of exogenous MG (Figure 9.6). Furthermore, the level of 17 kDa RAGE, induced by IL-3 depletion, was substantially higher than in the absence of exogenous MG.

17 kDa RAGE is also generated in Hek293 cells

To corroborate our findings in Ba/F3 cells we tested if 17 kDa RAGE could also be detected in other cell lines. Hek293 cells were therefore transiently transfected with a vector expressing $_{FL}$ RAGE and cells were depleted of serum 24 hours after transfection. Cells were lysed 24 hours after depletion and analyzed by immunoblotting with C-t-Ab (Figure 9.7.A). In control cells and mock transfected cells, the 17 kDa RAGE could not be detected. However, when cells were transfected with $_{FL}$ RAGE and cultured in the absence of serum, the formation of 17 kDa RAGE was induced. This shows that the 17 kDa RAGE is not restricted to a specific cell line or species (mouse and human), hereby suggesting induction of 17 kDa RAGE is a general mechanism. In addition, 17 kDa RAGE was also observed in the mouse β -cell line MIN6 (Toepoel et al., unpublished results).

Several mechanisms can be responsible for the formation of 17 kDa RAGE, the two most plausible are alternative splicing and proteolytic cleavage. To examine these




Figure 9.6. 17kD RAGE induction by exogenous MG.

G_1 arrest was induced by depletion of IL-3 and exogenous MG was added. Cells were lysed in CEB after indicated time intervals, equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-t-Ab.

mechanisms, Hek293 cells were transiently transfected with vectors containing the cDNA encoding C-terminal flagged RAGE and N-terminal flagged RAGE (designated RAGE-C-flag and N-flag-RAGE respectively). As expected, analysis of immunoblots with an antibody directed against the flag-tag showed high levels of overexpression of RAGE with either construct. However, the generation of 17 kDa RAGE could be observed using an antibody raised against the flag-tag, only when this tag was attached at the C-terminus of RAGE, hereby validating the observations made with the C-terminus directed anti-RAGE antibody (Figure 9.7.B). Most isoforms described to date have been reported to result from alternative splicing and posses different C-termini (reviewed in (40,41)). However, since 17 kDa RAGE can also be generated by overexpression of RAGE using a cDNA, alternative splicing can be excluded.

In the lysates of the cells transfected with N-flag-RAGE an additional band could be observed with a molecular mass of approximately 45 kDa (Figure 9.7.B, arrowhead). This band could very well be the remainder of the cleaved RAGE. The molecular weight differs from the band seen near the 75 kDa marker, which probably represents the glycosylated full length flag-tagged RAGE, and the difference in molecular weight could be explained by the removal of the 17 kDa RAGE part. This band was absent in the lysates of cells transfected with the RAGE-C-flag, endorsing that this isoform had an alternative C-terminus, but an intact N-terminus. Several reports indicate the involvement of protein cleavage in the formation of a soluble form of RAGE in mice (43,52). Moreover, recently it has been reported that limited proteolysis of RAGE with trypsin also gives rise to low molecular weight bands with comparable migratory properties (47), hereby further emphasizing the hypothesized proteolytic cleavage of RAGE. It should be noted however that 17 kDa was present even when serum was not depleted.

Inhibition of caspases partially inhibits the generation of 17 kDa RAGE

Our data excludes alternative splicing as a mechanism to generate 17 kDa RAGE and suggest the involvement of proteolytic cleavage. Proteolytic cleavage was subsequently further explored by using several specific inhibitors. Initial experiments with different protease inhibitors showed no effects (data not shown). Since IL-3 starvation of Ba/F3 cells also leads to apoptosis, we sought to determine if the inhibition of caspase activation could inhibit formation of 17 kDa RAGE. When Ba/F3 cells were starved of IL-3, 17 kDa RAGE formation was induced, as observed by analysis of lysates using immunoblotting with C-t-Ab. In the presence of the pan-caspase inhibitor Z-VAD-FMK however generation of 17 kDa RAGE was partially inhibited (Figure 9.8.A, arrow). Other isoforms, such as the 50 kDa from were unaffected (Figure 9.8.A, arrowhead). Subsequently Hek293 cells were transiently transfected with pCAGGS-RAGE-C-flag and depleted of serum in the presence of Z-VAD-FMK. As observed by immunoblotting with an antibody against flag, inhibition of

caspace activation led to a decrease in 17 kDa RAGE (Figure 9.8.B). Formation of 17 kDa RAGE was inhibited by Z-VAD-FMK to the same extent in Hek293 cells as in Ba/F3 cells.

Interestingly, in the modelsystem with RAGE overexpression in Hek293, a strong induction of 17 kDa was observed, even in the presence of serum (Figure 9.7.B and 9.8.B).




Figure 9.7. Generation of 17 kDa RAGE in Hek293 cells.

A. Presence of 17 kDa RAGE in Hek293 cells. Hek293 cells were transfected with expression vectors containing full length RAGE. After 24 hours, transfected cells were serum-starved over night. Lysates were made in CEB and equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-t-Ab.

B. Formation of 17 kDa RAGE in Hek293 cells. Hek293 cells were transfected with expression vectors containing either N-flag-RAGE or RAGE-C-flag. After 24 hours, transfected cells were serum-starved for 1 or 2 days. Lysates were made in CEB and equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using an anti-flag antibody (Sigma Aldrich).

A possible hypothesis would be it is induced by the G_1 arrest caused by contact inhibition (53-56). In the presence of serum, however, addition of Z-VAD-FMK led to a strong rise in the level of 17 kDa RAGE.

Analysis of posttranslational modification of RAGE during the cell cycle using 2-dimensional gel electrophoresis

In order to further examine the possible involvement of RAGE in the cell cycle we also analyzed whether posttranslational modification of RAGE occurs during the cell cycle, using 2-dimensional gel electrophoresis (2-DE). The cell cycle of Ba/F3 cultures was synchronised in the G_1 phase by IL-3 depletion over night and subsequently released from this block by re-addition of IL-3. Cells were lysed at different time points after the re-addition of IL-3, when cells were in different phases of the cell cycle.

Immunoblots are shown in Figure 9.9.A which were detected with C-t-Ab. Although the most acidic isoform remains constant, the displayed isoforms strongly alternate. The upper panel shows lysates of a Ba/F3 culture in G_1 . Three isoforms can be distinguished at this time, the two most acidic however are not well separated and appear almost fused. In the S phase, the middle spot disappeared, and seems to have merged with the most acidic spot. In the G_2 /M phase the pattern of the two most acidic isoforms strongly resembles the pattern in G_1 , but the basic isoform had vanished. Early in the G_1 phase of the following cycle, the most basic isoform reappears and in late G_1 two new spots arise, more basic than the basic spot observed so far. The newly formed basic spots then seem to shift again towards the more acidic side during the S phase.

To date the only posttranslational modification described for RAGE is glycosylation (43-45). Differences in glycosylation have however been shown to result in differences in molecular weight, which are not observed here. Unpublished data from Salliau et al. suggest phosphorylation might play a part in the alternating pattern observed. The short cytoplasmic domain of RAGE contains 2 predicted sites for phosphorylation, and when RAGE was purified from metabolically labeled L929 cells, specific spots were observed on autoradiographs from 2D gels (Salliau et al., unpublished data). Since these patterns observed on 2-dimensional electrophoresis strongly resemble the patterns observed for GLO1 and arise analogous with the modifications of GLO1 (see chapter 6 and data not shown), it is conceivable that phosphorylation indeed is responsible for at least part of the modifications observed.

To further substantiate the cell cycle-dependent modification of RAGE, we then investigated if the induction of G_1 arrest by the depletion of IL-3 also induced alterations in the pattern of displayed isoforms of RAGE (Figure 9.9.B). We depleted Ba/F3 cells of IL-3




Fig. 9.8. Inhibition of the generation of 17 kDa RAGE.

A. 17kDa RAGE generation in Ba/F3 cells. Ba/F3 cells were IL-3-depleted and Z-VAD-FMK (20 μ M) was added. After 24 hours, cells were lysed in CEB and equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using the C-terminus-directed anti-RAGE antibody. **B.** Generation of 17 kDa in Hek293 cells. Hek293 cells were transfected with expression vectors containing RAGE-C-flag. After 24 hours, transfected cells were serum-starved over night and Z-VAD-FMK (20 μ M) was added. Lysates were made in CEB and equal protein concentrations were separated by SDS-PAGE and transferred onto PVDF using electrotransfer. Expression of different isoforms of RAGE was analysed using an anti-flag antibody.

and lysates were made after the indicated time intervals. Asynchronously growing Ba/F3 cells displayed four isoforms, the pattern showed strong similarities with the pattern seen in late G_1 (Figure 9.9.A), but a fourth spot was also detected, situated between the most acidic isoform and the duplet of more basic isoforms. After 1 and 3 hours the second most acidic isoform vanishes and the most basic spot shifts towards the more basic side. After 7 hours of IL-3 starvation only three isoforms remain and the observed pattern strongly resembles the pattern observed in late G_1 .

DISCUSSION

Activation of RAGE is not completely understood and knowledge is only fragmented. RAGE expression is high during development, repressed in mature tissues, and upregulated again in the course of several pathologies such as in different cancers, so RAGE signaling has often been involved in the regulation of cell cycle (3). The involvement of RAGE in proliferation however seems rather dualistic, since RAGE activation has been shown to be involved in proliferation (9,23,26,30,31) as well as in growth arrest and differentiation (32,36-38,57,58), two processes that are consecutive and mutually exclusive.

Three major isoforms of RAGE have been described (reviewed in (40)). However, additional splice variants and isoforms keep on being discovered, sometimes with a high

specificity for certain tissues (41,42). Hence, the outcome of RAGE signaling is most likely highly tissue-specific. Moreover, it is clear that effects mediated by RAGE are not so much the result of one specific isoform of RAGE, but of a concerted interplay of the different isoforms and the relative availability of different ligands. This prompted us to study which different isoforms are present in Ba/F3 cells and whether these were regulated in a cell cycle dependent manner. Although we did not detect a difference in total RAGE expression on the membrane, RAGE expression was indeed strongly modulated throughout the cell cycle. We were able to correlate different forms of RAGE to different phases in the cell cycle and we observed modulation of these isoforms at phase transitions.

However, given the high molecular weight of the isoforms described, the observed effects can be disputable. The high molecular weight isoforms of RAGE observed are hypothesized to be oligomers or complexes containing RAGE. Such complexation and oligomerization has been described for RAGE (43,47,59). Moreover, the specificity of the antibodies we used was demonstrated earlier (Salliau et al., submitted). Our hypothesis is further emphasized by the observation of high molecular weight bands when flag-tagged RAGE was overexpressed in Hek293 cells. Immunoblotting with an antibody raised against




Figure 9.9. RAGE is modified during the cell cycle and IL-3 withdrawal-induced G_1 arrest.

A. RAGE is posttranslationally modified during the cell cycle. Ba/F3 cells were synchronized by IL-3 starvation over night. Cell cycle was induced by readdition of IL-3. Samples were taken in several phases of the cell cycle. Cells were lysed in CEB and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with C-t-Ab. **B.** RAGE is posttranslationally modified during IL-3 withdrawal-induced G_1 arrest. Ba/F3 cells were depleted of IL-3, cells were lysed in CEB at the indicated time points and after ethanol precipitation resuspended in 2-DE lysis buffer. Protein concentrations were determined and 200 μ g of the lysates was loaded for separation by 2-dimensional gel electrophoresis (pH 4-7). Representative western blots of 2-DE gels developed with C-t-Ab.

the flag-tag also showed high molecular weight bands (data not shown).

We also demonstrated the presence of a novel low molecular weight isoform of RAGE. This isoform is strongly induced during G₁ arrest, but was present at several time points in the S phase as well. This isoform was not restricted to one specific cell type, since it could be observed in three different cell lines and two different species, mouse and human. This low molecular weight form of RAGE was most likely a product of proteolytic processing of FLRAGE since it also was formed upon overexpression from a C-terminally tagged RAGE, expressed from a cDNA, hereby excluding splicing. The precise mechanism could not be clarified yet. However, addition of MG, a cytostatic and cytotoxic compound which exerts part of its activity in a RAGE-dependent manner (60), strongly enhanced formation of this isoform.

Formation of 17 kDa RAGE upon IL-3 or serum starvation could partially be inhibited by inhibition of caspases. However, overexpression of full length RAGE could also strongly induce formation of 17 kDa RAGE, even in the presence of serum. This formation could not be inhibited by caspase inhibition, but addition of Z-VAD-FMK even seemed to enhance its formation, hereby seemingly contradicting the observations made in the depletion model system. Possibly different or redundant mechanism underlie these observations. RAGE could be processed in order to attenuate the strong signaling caused by the high levels of RAGE expression. Alternatively, the presence of the C-terminal flag-tag could interfere with the function of RAGE, leading to the breakdown of the receptor. The strong formation of 17 kDa RAGE in Hek293 cells in the presence of serum was not joined by an increase in apoptosis, suggesting that the presence of 17 kDa RAGE per se did not induce apoptosis.

The hypothesis that 17 kDa RAGE is formed by proteolysis is in line with earlier observations (43,52). Proteolytic cleavage of RAGE might be an important event in the regulation of RAGE signaling and it is conceivable that proteolytic processing of RAGE provides a negative regulation or a negative feedback loop in RAGE signaling. Either to inhibit prosurvival signaling or signaling towards proliferation, when IL-3 or serum is absent, or to downregulate RAGE signaling to prevent overactivation.

When high resolution 2-DE was used to study post translational modification of RAGE, our data further substantiate that RAGE is strongly susceptible to regulation during the progression of the cell cycle and when growth arrest is induced by IL-3 depletion. To date the only described posttranslational modification of RAGE is glycosylation (43-46). The different spots we observed could be differences in the level of glycosylation, since this would yield a different pI and deglycosylation would be able to account for the shifts observed. Glycosylation would however also alter the molecular weight of the isoforms, and deglycosylation would therefore not only induce a shift towards the basic side, but most likely also result in a faster migration in the second dimension. However, while we

were able to observe differences in pI, concomitant reductions in molecular weight could not be detected.

We observed that the shifts in pI strongly coincided with the modifications observed in GLO1 (see chapter 6 and data not shown). Furthermore, results from our lab showed that when L929 cells were metabolically labelled and stimulated with tumor necrosis factor (TNF), RAGE could be phosphorylated (Salliau, unpublished results). These observations suggest that the modifications shown are possibly phosphorylation and dephosphorylation.

In summary, the data shown here indicate a strong regulation of RAGE during the cell cycle. This regulation appears to take place at several levels. At a first level, we observed a strong correlation of certain isoforms with different cell cycle phases. This different expression could not be related to the expression of RAGE on the plasmamembrane, since the levels of membrane expression seemed to remain constant during the cell cycle. This could reflect the different requirement for signaling during different phases or a difference in the available ligands for RAGE. We also observed RAGE was regulated by proteolysis, a possible additional regulatory mechanism, or a negative feedback loop in RAGE signaling. Furthermore studies using 2-DE further show RAGE is posttranslationally modified, most likely phosphorylated. We therefore conclude that RAGE signaling is regulated at different levels during cell cycle and growth factor withdrawal-induced G₁ arrest and consequent apoptosis. This stresses the importance of RAGE signaling in the regulation of cell cycle and cell death and further emphasizes the complex nature of RAGE signaling. Our observations could also partially provide a basis for the dualistic nature role of RAGE signaling in proliferation and differentiation, and in survival and the induction of cell death.

MATERIALS AND METHODS

Cell Lines and Cultures

The IL-3 dependent pro-B cell line Ba/F3 (61) was purchased from DSMZ. Ba/F3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) with glutamax (Invitrogen) supplemented with heat-inactivated FBS (10% v/v) (Cambrex), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) and 5% conditioned medium from WEHI-3B cells as a source of mouse IL-3. Ba/F3 cells were cultured at 37°C in a humidified incubator under an 5% CO₂ atmosphere.

Hek293 cells, were cultured in DMEM with glutamax (Invitrogen) supplemented with heat-inactivated FBS (10% v/v), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) at 37°C in a humidified incubator under an 8% CO₂ atmosphere.

Cell stimulations

Ba/F3 cells were synchronized by washing exponentially growing cells 3 times in culture medium without conditioned WEHI-3B medium and resuspended in culture medium without conditioned WEHI-3B medium for 8 hours or over night. At the end of the synchronisation conditioned WEHI-3B medium (5% v/v) was readded. This time point was designated 0 hours.

For IL-3 depletion experiments, IL-3 was removed by washing exponentially growing Ba/F3 cells 3 times in culture medium without conditioned WEHI-3B medium and resuspended in culture medium without conditioned WEHI-3B medium.

Hek293 cells were depleted by washing them 3 times with culture medium without FBS and subsequently cultured in culture medium in the absence of FBS.

Hek293 cells were transiently transfected by Ca-phosphate-precipitation. Vectors used are pCAGGS vectors containing full length RAGE (FLRAGE), N-terminally flag tagged full length RAGE (N-flag-RAGE), or C-terminally tagged full length RAGE (RAGE-C-flag). The vectors were kindly provided by Stefanie Salliau (VIB department of Medical Protein Research, Molecular and Metabolic Signaling Unit, Flanders Institute for Biotechnology, Faculty of Medicine and Health Sciences, Ghent University).

Chemicals and inhibitors used are nocodazole (0.8 μ M; Sigma aaldrich) and Z-VAD-FMK (promega). Duration and concentrations used as described.

DNA distribution analysis

Cells were stained with propidium iodide (PI, Sigma) containing staining solution (62) and analyzed by a FACSCalibur flow cytometer (488_{Ex}/590_{Em}). Cell Quest software was used to analyze cell cycle distribution (FACS Calibur, Becton Dickinson) (62,63). Ten thousand cells were routinely analyzed.

Electrophoresis and immunoblotting

The cells were washed 3 times with ice-cold PBS buffer and lysed with cytosol extraction buffer (10 mM Tris-HCl pH7.4, 50 mM EDTA pH8.0, 25 mM NaCl, 0,7% TritonX-100, 100 mM PMSF, 1 tablet Complete Protease Inhibitor Cocktail (Roche)/50 ml cytosol extraction buffer) or modified radioimmune precipitation assay buffer (200 mM NaCl, 50 mM Tris-HCl (pH 8), 0.05% SDS, 2 mM EDTA, 1% Nonidet P-40, 0.5% deoxycholic acid, 1 mM Na₃VO₄, 1 mM NaF, 20 mM β -glycerophosphate, and Complete Protease Inhibitor Cocktail). Cell lysates were cleared by centrifugation (14,000 x g). Protein concentrations

were determined using Bradford method. Proteins were separated by SDS-PAGE and transferred onto a PVDF membrane (Hybond-P, Amersham Pharmacia Biotech). The blots were incubated with the desired antibodies followed by ECL-based detection (Amersham Pharmacia Biotech). Quantification of protein bands was performed using Genetools software (Westburg).

Preparation of cytosolic and nuclear fractions

Nuclear and cytoplasmic extracts were prepared as previously described with small modifications (64). In brief, 10×10^6 Ba/F3 cells were synchronized and after indicated time intervals collected. After harvesting and washing the cells 3 times with ice-cold PBS, the cells were lysed in 200 μ l buffer A (10 mM HEPES, pH 8.0, 1.5 mM $MgCl_2$, 10 mM KCl, 0.5 mM dithiothreitol, 200 mM sucrose, 1 tablet Complete Protease Inhibitor Cocktail/50 ml buffer A, 0.5% NP-40). The lysates were incubated 15 min on ice and pelleted at $11,000 \times g$ for 10 min and the cytoplasmic supernatant was transferred to a new tube. The remaining nuclear pellets were washed twice with buffer A and resuspended in 150 μ L buffer B (20 mM HEPES, pH 8.0, 20% (vol/vol) glycerol, 0.1 M KCl, 0.2 mM EDTA, 0.5 mM dithiothreitol, 1 tablet Complete Protease Inhibitor Cocktail/50 ml, 1% NP-40). These extracts were subsequently passed 1 time through a 18 G needle and 3 to 4 times through a 27 G needle and centrifuged 30 sec at $14,000 \times g$. The pelleted debris was discarded, while the nuclear extract and cytosolic supernatants were used for immunoblot analysis.

2-Dimensional Gel Electrophoresis

Isoelectric focusing was carried out on 18 cm IPG strips, pH 4-7 (GE Healthcare) according to the manufacturer's instructions. For the second dimension, proteins were separated by SDS-polyacrylamide gel electrophoresis (PAGE, 12%).

Surface staining of RAGE

1×10^6 Ba/F3 cells were washed 3 times in ice-cold FACS buffer (PBS, 1% FBS, 0.5 mM EDTA) and incubated with a polyclonal antibody raised against the N-terminus of RAGE (H-300, Santa Cruz Biotechnology) for 20 min. Subsequently cells were washed twice in FACS buffer and incubated with a secondary Alexa R488 conjugated anti-rabbit antibody (Molecular Probes) for 20 min. Cells were then washed and immediately measured and analysed on FACS.

Reference List

1. **Schmidt, A. M., M. Vianna, M. Gerlach, J. Brett, J. Ryan, J. Kao, C. Esposito, H. Hegarty, W. Hurley, M. Clauss, F. Wang, Y. C. E. Pan, T. C. Tsang, and D. Stern.** 1992. Isolation and Characterization of 2 Binding-Proteins for Advanced Glycosylation End-Products from Bovine Lung Which Are Present on the Endothelial-Cell Surface. *Journal of Biological Chemistry* **267**:14987-14997.
2. **Neeper, M., A. M. Schmidt, J. Brett, S. D. Yan, F. Wang, Y. C. E. Pan, K. Elliston, D. Stern, and A. Shaw.** 1992. Cloning and Expression of A Cell-Surface Receptor for Advanced Glycosylation End-Products of Proteins. *Journal of Biological Chemistry* **267**:14998-15004.
3. **Schmidt, A. M., S. D. Yan, S. F. Yan, and D. M. Stern.** 2000. The biology of the receptor for advanced glycation end products and its ligands. *Biochim.Biophys.Acta* **1498**:99-111.
4. **Hori, O., J. Brett, T. Slaterry, R. Cao, J. H. Zhang, J. X. Chen, M. Nagashima, E. R. Lundh, S. Vijay, D. Nitecki, J. Morser, D. Stern, and A. M. Schmidt.** 1995. The Receptor for Advanced Glycation End-Products (Rage) Is A Cellular-Binding Site for Amphoterin - Mediation of Neurite Outgrowth and Coexpression of Rage and Amphoterin in the Developing Nervous-System. *Journal of Biological Chemistry* **270**:25752-25761.
5. **Hofmann, M. A., S. Drury, C. F. Fu, W. Qu, A. Taguchi, Y. Lu, C. Avila, N. Kambham, A. Bierhaus, P. Nawroth, M. F. Neurath, T. Slaterry, D. Beach, J. McClary, M. Nagashima, J. Morser, D. Stern, and A. M. Schmidt.** 1999. RAGE mediates a novel proinflammatory axis: A central cell surface receptor for S100/calgranulin polypeptides. *Cell* **97**:889-901.
6. **Schmidt, A. M., S. Du Yan, S. F. Yan, and D. M. Stern.** 2001. The multiligand receptor RAGE as a progression factor amplifying immune and inflammatory responses. *Journal of Clinical Investigation* **108**:949-955.
7. **Yan, S. D., X. Chen, J. Fu, M. Chen, H. J. Zhu, A. Roher, T. Slaterry, L. Zhao, M. Nagashima, J. Morser, A. Migheli, P. Nawroth, D. Stern, and A. M. Schmidt.** 1996. RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* **382**:685-691.
8. **Yan, S. D., H. J. Zhu, A. P. Zhu, A. Golabek, H. Du, A. Roher, J. Yu, C. Soto, A. M. Schmidt, D. Stern, and M. Kindy.** 2000. Receptor-dependent cell stress and amyloid accumulation in systemic amyloidosis. *Nature Medicine* **6**:643-651.
9. **Cortizo, A. M., M. G. Lettieri, D. A. Barrio, N. Mercer, S. B. Etcheverry, and A. D. McCarthy.** 2003. Advanced glycation end-products (AGEs) induce concerted changes in the osteoblastic expression of their receptor RAGE and in the activation of extracellular signal-regulated kinases (ERK). *Molecular and Cellular Biochemistry* **250**: 1-10.
10. **Lander, H. M., J. M. Tauras, J. S. Ogiste, O. Hori, R. A. Moss, and A. M. Schmidt.** 1997. Activation of the receptor for advanced glycation end products triggers a p21(ras)-dependent mitogen-activated protein kinase pathway regulated by oxidant stress. *Journal of Biological Chemistry* **272**:17810-17814.
11. **Sorci, G., F. Riuzzi, A. L. Agneletti, C. Marchetti, and R. Donato.** 2004. S100B causes apoptosis in a myoblast cell line in a RAGE-independent manner. *Journal of Cellular Physiology* **199**:274-283.
12. **Wautier, M. P., O. Chappey, S. Corda, D. M. Stern, A. M. Schimidt, and J. L. Wautier.** 2001. Activation of NADPH oxidase by AGE links oxidant stress to altered gene expression via RAGE. *American Journal of Physiology-Endocrinology and Metabolism* **280**:E685-E694.
13. **Taguchi, A., D. C. Blood, G. del Toro, A. Canet, D. C. Lee, W. Qu, N. Tanji, Y. Lu, E. Lalla, C. F. Fu, M. A. Hofmann, T. Kislinger, M. Ingram, A. Lu, H. Tanaka, O. Hori, S. Ogawa, D. M. Stern, and A. M. Schmidt.** 2000. Blockade of RAGE-amphoterin signalling suppresses tumour growth and metastases. *Nature* **405**:354-360.
14. **Yeh, C. H., L. Sturgis, J. Haidacher, X. N. Zhang, S. J. Sherwood, R. J. Bjercke, O. Juhasz, M. T. Crow, R. G. Tilton, and L. Denner.** 2001. Requirement for p38 and p44/p42 mitogen-activated protein kinases in RAGE-mediated nuclear factor-kappa B transcriptional activation and cytokine secretion. *Diabetes* **50**:1495-1504.
15. **Huang, J. S., J. Y. Guh, H. C. Chen, W. C. Hung, Y. H. Lai, and L. Y. Chuang.** 2001. Role of receptor for advanced glycation end-product (RAGE) and the JAK/STAT-signaling pathway in AGE-induced collagen production in NRK-49F cells. *Journal of Cellular Biochemistry* **81**:102-113.
16. **Sakaguchi, T., S. F. Yan, S. Du Yan, D. Belov, L. L. Rong, M. Sousa, M. Andrassy, S. P. Marso, S. Duda, B. Arnold, B. Liliensiek, P. P. Nawroth, D. M. Stern, A. M. Schmidt, and Y. Naka.** 2003. Central role of RAGE-dependent neointimal expansion in arterial restenosis. *Journal of Clinical Investigation* **111**:959-972.
17. **Huttunen, H. J., C. Fages, and H. Rauvala.** 1999. Receptor for advanced glycation end products (RAGE)-mediated neurite outgrowth and activation of NF-kappa B require the cytoplasmic domain of the receptor but different downstream signaling pathways. *Journal of Biological Chemistry* **274**:19919-19924.
18. **Dukic-Stefanovic, S., J. Gasic-Milenkovic, W. Deuther-Conrad, and G. Munch.** 2003. Signal transduction pathways in mouse microglia N-11 cells activated by advanced glycation endproducts (AGEs). *Journal of Neurochemistry* **87**:44-55.
19. **Leclerc, E., G. Fritz, M. Weibel, C. W. Heizmann, and A. Galichet.** 2007. S100B and S100A6 differentially modulate cell survival by interacting with distinct RAGE (receptor for advanced glycation end products) immunoglobulin domains. *Journal of Biological Chemistry* **282**:31317-31331.
20. **Arumugam, T., D. M. Simeone, K. Van Golen, and C. D. Logsdon.** 2005. S100P promotes pancreatic cancer growth, survival, and invasion. *Clinical Cancer Research* **11**:5356-5364.
21. **Arumugam, T., V. Ramachandran, and C. D. Logsdon.** 2006. Effect of cromolyn on S100P interactions with RAGE and pancreatic cancer growth and invasion in mouse models. *Journal of the National Cancer Institute* **98**: 1806-1818.
22. **Fuentes, M. K., S. S. Nigavekar, T. Arumugam, C. D. Logsdon, A. M. Schmidt, J. C. Park, and E. H. Huang.**

2007. RAGE activation by S100P in colon cancer stimulates growth, migration, and cell signaling pathways. *Diseases of the Colon & Rectum* **50**:1230-1240.
23. **Geoffroy, K., N. Wiernsperger, M. Lagarde, and S. El Bawab.** 2004. Bimodal effect of advanced glycation end products on mesangial cell proliferation is mediated by neutral ceramidase regulation and endogenous sphingolipids. *Journal of Biological Chemistry* **279**:34343-34352.
24. **Kuniyasu, H., Y. Chihara, and H. Kondo.** 2003. Differential effects between amphotericin and advanced glycation end products on colon cancer cells. *International Journal of Cancer* **104**:722-727.
25. **Sasahira, T., T. Kirita, U. K. Bhawal, K. Yamamoto, H. Ohmori, K. Fujii, and H. Kuniyasu.** 2007. Receptor for advanced glycation end products (RAGE) is important in the prediction of recurrence in human oral squamous cell carcinoma. *Histopathology* **51**:166-172.
26. **Satoh, H., M. Togo, M. Hara, T. Miyata, K. Han, H. Maekawa, N. Ohno, Y. Hashimoto, K. Kurokawa, and T. Watanabe.** 1997. Advanced glycation endproducts stimulate mitogen-activated protein kinase and proliferation in rabbit vascular smooth muscle cells. *Biochemical and Biophysical Research Communications* **239**:111-115.
27. **Takada, M., T. Koizumi, H. Toyama, Y. Suzuki, and Y. Kuroda.** 2001. Differential expression of RAGE in human pancreatic carcinoma cells. *Hepato-Gastroenterology* **48**:1577-1578.
28. **Takada, M., K. Hirata, T. Ajiki, Y. Suzuki, and Y. Kuroda.** 2004. Expression of receptor for advanced glycation end products (RAGE) and MMP-9 in human pancreatic cancer cells. *Hepato-Gastroenterology* **51**:928-930.
29. **Arumugam, T., D. M. Simeone, A. M. Schmidt, and C. D. Logsdon.** 2004. S100P stimulates cell proliferation and survival via receptor for activated glycation end products (RAGE). *Journal of Biological Chemistry* **279**:5059-5065.
30. **Munch, G., J. Gasic-Milenkovic, and T. Arendt.** 2003. Effect of advanced glycation endproducts on cell cycle and their relevance for Alzheimer's disease. *Journal of Neural Transmission-Supplement* **63**:71.
31. **Schmidt, A., B. Kuhla, K. Bigl, G. Munch, and T. Arendt.** 2007. Cell cycle related signaling in neuro2a cells proceeds via the receptor for advanced glycation end products. *Journal of Neural Transmission* **114**:1413-1424.
32. **Huttunen, H. J., C. Fages, J. Kuja-Panula, A. J. Ridley, and H. Rauvala.** 2002. Receptor for advanced glycation end products-binding COOH-terminal motif of amphotericin inhibits invasive migration and metastasis. *Cancer Research* **62**:4805-4811.
33. **Bartling, B., H. S. Hofmann, B. Weigle, R. E. Silber, and A. Simm.** 2005. Down-regulation of the receptor for advanced glycation end-products (RAGE) supports non-small cell lung carcinoma. *Carcinogenesis* **26**:293-301.
34. **Bartling, B., N. Demling, R. E. Silber, and A. Simm.** 2006. Proliferative stimulus of lung fibroblasts on lung cancer cells is impaired by the receptor for advanced glycation end-products. *American Journal of Respiratory Cell and Molecular Biology* **34**:83-91.
35. **Hofmann, H. S., G. Hansen, S. Burdach, B. Bartling, R. E. Silber, and A. Simm.** 2004. Discrimination of human lung neoplasm from normal lung by two target genes. *American Journal of Respiratory and Critical Care Medicine* **170**:516-519.
36. **Riuzzi, F., G. Sorci, and R. Donato.** 2007. RAGE expression in rhabdomyosarcoma cells results in myogenic differentiation and reduced proliferation, migration, invasiveness, and tumor growth. *American Journal of Pathology* **171**:947-961.
37. **Brizzi, M. F., P. Dentelli, A. Rosso, C. Calvi, R. Gambino, M. Cassader, G. Salvidio, G. Deferrari, G. Camussi, L. Pegoraro, G. Pagano, and P. Cavallo-Perin.** 2004. RAGE- and TGF-beta receptor-mediated signals converge on STAT5 and P21(waf) to control cell-cycle progression of mesangial cells: a possible role in the development and progression of diabetic nephropathy. *FASEB Journal* **18**:1249-+.
38. **Dumitriu, I. E., P. Baruah, B. Valentinis, R. E. Voll, M. Herrmann, P. P. Nawroth, B. Arnold, M. E. Bianchi, A. A. Manfredi, and P. Rovere-Querini.** 2005. Release of high mobility group box 1 by dendritic cells controls T cell activation via the receptor for advanced glycation end products. *Journal of Immunology* **174**:7506-7515.
39. **Ge, J. B., Q. Z. Jia, C. Liang, Y. K. Luo, D. Huang, A. J. Sun, K. Q. Wang, Y. Z. Zou, and H. Z. Chen.** 2005. Advanced glycosylation end products might promote atherosclerosis through inducing the immune maturation of dendritic cells. *Arteriosclerosis Thrombosis and Vascular Biology* **25**:2157-2163.
40. **Ding, Q. X. and J. N. Keller.** 2005. Evaluation of RAGE isoforms, ligands, and signaling in the brain. *Biochimica et Biophysica Acta-Molecular Cell Research* **1746**:18-27.
41. **Ding, Q. X. and J. N. Keller.** 2005. Splice variants of the receptor for advanced glycosylation end products (RAGE) in human brain. *Neuroscience Letters* **373**:67-72.
42. **Schlueter, C., S. Hauke, A. M. Flohr, P. Rogalla, and J. Bullerdiek.** 2003. Tissue-specific expression patterns of the RAGE receptor and its soluble forms - a result of regulated alternative splicing? *Biochimica et Biophysica Acta-Gene Structure and Expression* **1630**:1-6.
43. **Hanford, L. E., J. J. Enghild, Z. Valnickova, S. V. Petersen, L. M. Schaefer, T. M. Schaefer, T. A. Reinhart, and T. D. Oury.** 2004. Purification and characterization of mouse soluble receptor for advanced glycation end products (sRAGE). *Journal of Biological Chemistry* **279**:50019-50024.
44. **Yonekura, H., Y. Yamamoto, S. Sakurai, R. G. Petrova, M. J. Abedin, H. Li, K. Yasui, M. Takeuchi, Z. Makita, S. Takasawa, H. Okamoto, T. Watanabe, and H. Yamamoto.** 2003. Novel splice variants of the receptor for advanced glycation end-products expressed in human vascular endothelial cells and pericytes, and their putative roles in diabetes-induced vascular injury. *Biochemical Journal* **370**:1097-1109.
45. **Harashima, A., Y. Yamamoto, C. M. Cheng, K. Tsuneyama, K. M. Myint, A. Takeuchi, K. Yoshimura, H. Li, T. Watanabe, S. Takasawa, H. Okamoto, H. Yonekura, and H. Yamamoto.** 2006. Identification of mouse orthologue of endogenous secretory receptor for advanced glycation end-products: structure, function and

- expression. *Biochemical Journal* **396**:109-115.
46. **Lizotte, P. P., L. E. Hanford, J. J. Enghild, E. Nozik-Grayck, B. L. Giles, and T. D. Oury.** 2007. Developmental expression of the receptor for advanced glycation end-products (RAGE) and its response to hyperoxia in the neonatal rat lung. *Bmc Developmental Biology* **7**.
47. **Dattilo, B. M., G. Fritz, E. Leclerc, C. W. Vander Kooi, C. W. Heizmann, and W. J. Chazin.** 2007. The extracellular region of the receptor for advanced glycation end products is composed of two independent structural units. *Biochemistry* **46**:6957-6970.
48. **Ahmed, N., U. Ahmed, P. J. Thornalley, K. Hager, G. Fleischer, and G. Munch.** 2005. Protein glycation, oxidation and nitration adduct residues and free adducts of cerebrospinal fluid in Alzheimer's disease and link to cognitive impairment. *Journal of Neurochemistry* **92**:255-263.
49. **Shinohara, M., P. J. Thornalley, I. Giardino, P. Beisswenger, S. R. Thorpe, J. Onorato, and M. Brownlee.** 1998. Overexpression of glyoxalase-I in bovine endothelial cells inhibits intracellular advanced glycation endproduct formation and prevents hyperglycemia-induced increases in macromolecular endocytosis. *Journal of Clinical Investigation* **101**:1142-1147.
50. **Li, J. F. and A. M. Schmidt.** 1997. Characterization and functional analysis of the promoter of RAGE, the receptor for advanced glycation end products. *Journal of Biological Chemistry* **272**:16498-16506.
51. **Tanaka, N., H. Yonekura, S. Yamagishi, H. Fujimori, Y. Yamamoto, and H. Yamamoto.** 2000. The receptor for advanced glycation end products is induced by the glycation products themselves and tumor necrosis factor- α through nuclear Factor-kappa B, and by 17 beta-estradiol through Sp-1 in human vascular endothelial cells. *Journal of Biological Chemistry* **275**:25781-25790.
52. **Devaux, Y., R. M. Senior, and P. Ray.** 2004. RAGE: A new target for MMP-9 in the regulation of inflammatory response in the lung during oxidative stress. *Faseb Journal* **18**:A1147.
53. **Eagle, H. and E. M. Levine.** 1967. Growth Regulatory Effects of Cellular Interaction. *Nature* **213**:1102-&.
54. **Gradl, G., D. Faust, F. Oesch, and R. J. Wieser.** 1995. Density-Dependent Regulation of Cell-Growth by Contactinhibin and the Contactinhibin Receptor. *Current Biology* **5**:526-535.
55. **Levine, E. M., Y. Becker, C. W. Boone, and H. Eagle.** 1965. Contact Inhibition Macromolecular Synthesis and Polyribosomes in Cultured Human Diploid Fibroblasts. *Proceedings of the National Academy of Sciences of the United States of America* **53**:350-&.
56. **Wieser, R. J., R. Heck, and F. Oesch.** 1985. Involvement of Plasma-Membrane Glycoproteins in the Contact-Dependent Inhibition of Growth of Human-Fibroblasts. *Experimental Cell Research* **158**:493-499.
57. **Huttunen, H. J., J. Kuja-Panula, G. Sorci, A. L. Agnelli, R. Donato, and H. Rauvala.** 2000. Coregulation of neurite outgrowth and cell survival by amphoterin and S100 proteins through receptor for advanced glycation end products (RAGE) activation. *Journal of Biological Chemistry* **275**:40096-40105.
58. **Riuzzi, F., G. Sorci, and R. Donato.** 2006. The amphoterin (HMGB1)/receptor for advanced glycation end products (RAGE) pair modulates myoblast proliferation, apoptosis, adhesiveness, migration, and invasiveness - Functional inactivation of rage in L6 myoblasts results in tumor formation in vivo. *Journal of Biological Chemistry* **281**:8242-8253.
59. **Schmidt, A. M., R. Mora, R. Cao, S. D. Yan, J. Brett, R. Ramakrishnan, T. C. Tsang, M. Simionescu, and D. Stern.** 1994. The Endothelial-Cell Binding-Site for Advanced Glycation End-Products Consists of A Complex - An Integral Membrane-Protein and A Lactoferrin-Like Polypeptide. *Journal of Biological Chemistry* **269**:9882-9888.
60. **Kalapos, M. P.** 1999. Methylglyoxal in living organisms - Chemistry, biochemistry, toxicology and biological implications. *Toxicology Letters* **110**:145-175.
61. **Palacios, R. and M. Steinmetz.** 1985. IL3-Dependent Mouse Clones That Express B-220 Surface-Antigen, Contain Ig Genes in Germ-Line Configuration, and Generate Lymphocytes-B In vivo. *Cell* **41**:727-734.
62. **Krishan, A.** 1990. Rapid DNA content analysis by the propidium iodide-hypotonic citrate method. *Methods Cell Biol.* **33**:121-125.
63. **Nicoletti, I., G. Migliorati, M. C. Pagliacci, F. Grignani, and C. Riccardi.** 1991. A Rapid and Simple Method for Measuring Thymocyte Apoptosis by Propidium Iodide Staining and Flow-Cytometry. *Journal of Immunological Methods* **139**:271-279.
64. **Ryczyn, M. A. and C. V. Clevenger.** 2000. Role of cyclophilins in somatolactogenic action. *Neuroimmunomodulation* **917**:514-521.

Part III:

General Conclusions

General conclusions

The ubiquitous and highly conserved nature of the glyoxalase system suggests a fundamental role in life. Indeed, the glyoxalase system has been linked to several cellular processes including cell proliferation and growth, the assembly of microtubules and vesicle transport (1,2). Still, the glyoxalase is a rather complex enzyme system for merely a detoxification reaction. Moreover, several other enzyme systems are also capable of detoxifying α -oxoaldehydes. These observations indicate the glyoxalase system may have a more profound physiological function. In the 1960s Nobel prize laureate Albert Szent-Györgyi hypothesized that the glyoxalase system exerts a regulatory role in cell division. Ever since then, several investigators reported a strong correlation between high levels of expression and high activity of glyoxalase I, and strong cell division. However, the exact mechanistic link has yet to be established (3).

In chapters 6 and 8, we describe a potential involvement of phosphorylated glyoxalase I in growth factor-depletion induced G_1 arrest and consequent apoptosis, and the regulation of the cell cycle. Previous research in our group showed for the first time that glyoxalase I can be phosphorylated in mammalian cells (4). Earlier findings in yeast by Inoue and his colleagues around 1990 showed that treatment with mating factor induces phosphorylation of glyoxalase I and alters glyoxalase I activity (5,6). Mating factors induce several responses in yeast, and one thereof is a cell cycle arrest in G_1 (7). As described in chapter 6, in the hematopoietic cell line Ba/F3 the induction of G_1 arrest by IL-3 depletion strongly induced multiple phosphorylation of glyoxalase I. Also during the cell cycle, glyoxalase was transiently phosphorylated. In the course of our study it was also reported that treatment of rice leafs with gibberellin, a regulator of plant growth, induced phosphorylation of glyoxalase I (8). Taken together, our data and the observations made by others in yeast and rice leafs, strongly suggest either an involvement of phosphorylated glyoxalase I in the regulation of the cell cycle or at least the necessity for cells to phosphorylate glyoxalase I during the cell cycle. However, the precise biological function of phosphorylated glyoxalase I remains to be determined. Although in yeast (5) and plants (9) phosphorylation of glyoxalase I has been reported to be accompanied by an increase in glyoxalase activity, this does not seem to be the case in mammalian cells ((4) and chapters 6 and 8). This might be explained in part by the fact that some plants express a novel plant-specific type of glyoxalase (10,11). As a consequence, our data rule out the possibility that, in mammalian cells, phosphorylation of glyoxalase I occurs in order to allow the cells to cope with higher levels of methylglyoxal, caused by a cell cycle-dependent increase of glycolysis and/or the increased oxidative stress. Together with cell division, growth factors dynamically regulate the glycolysis. Growth factor withdrawal on the other hand, leads to a decrease in glycolysis. It is therefore conceivable that growth factors link the regulation of

the glyoxalase system to the regulation of metabolism.

Studies with kinase inhibitors and *in vitro* kinase assays revealed the involvement of several kinases in pathways leading to the phosphorylation of glyoxalase I. Studies using kinase inhibitors showed the involvement of GSK-3 β in phosphorylation of glyoxalase I upon IL-3 starvation and the involvement of the PI3K/Akt/mTOR pathway in cell cycle dependent phosphorylation of glyoxalase I. On the other hand, *in vitro* studies pointed out that CaMKII can directly phosphorylate glyoxalase I. However, unlike inhibition of GSK-3 β , inhibition of CaMKII, could not inhibit G₁ arrest and apoptosis upon IL-3 depletion in Ba/F3 cells. On the other hand, in other cell signaling pathways, phosphorylation of GLO1 could be inhibited by pharmacological inhibition of CaMKII (Rondas et al., in preparation). These data indicate that glyoxalase I is a direct physiological target of CaMKII. Interestingly, multiple phosphorylations can be induced on glyoxalase I and several kinases and signaling pathways have been shown to be involved in the phosphorylation of glyoxalase I. It is therefore conceivable that multiple kinases can directly phosphorylate glyoxalase I *in vivo*, or that kinases can substitute for one another in the phosphorylation of glyoxalase I, in case the physiological kinase is impaired. Such mechanisms have been described e.g. for the phosphorylation of GSK-3 β (12,13). Alternatively, several pools of glyoxalase I might exist, and different pools might be regulated by various signaling pathways and might be involved in different cellular processes. This might explain how phosphorylation does not affect the detoxification of methylglyoxal. It might be also possible that depending on the site in glyoxalase I that is phosphorylated that effect on the enzyme and thus also the biological response can be different.

Concomitant with the phosphorylation of glyoxalase I, the formation of specific methylglyoxal-derived advanced glycation endproducts (MG-AGEs) could be observed. Currently we do not know the exact relationship between the phosphorylation of glyoxalase I, and the formation of MG-AGEs. One could speculate that formation of MG-AGEs is an indirect consequence of a phosphorylation-induced inhibition of glyoxalase I. However, this was shown not to be the case. Furthermore, phosphorylation of glyoxalase I could not be correlated with alterations in glyoxalase I activity. In yeast and plants, however, where phosphorylation was accompanied by an increase in activity of glyoxalase I (5,9). Quite interestingly, when phosphorylation of glyoxalase I was inhibited, also formation of specific MG-AGEs decreased, similar to the events in TNF-induced necrosis. These observations suggest that also in our cell system phosphorylated glyoxalase I and formation of MG-AGEs are on the same pathway.

The specific targets for methylglyoxal-modification are currently unknown. Although methylglyoxal-modification has been considered to be involved in development of several pathologies, such as vascular complications in diabetes, it's becoming more apparent that

methylglyoxal-modification of proteins could be of physiological importance as well (14,15). The presence of methylglyoxal is proportional to the rate of glycolytic flux. Since many cellular processes and signaling pathways influence glycolysis, methylglyoxal levels and methylglyoxal-modification of specific proteins could function as a sentinel for the metabolic state of cells. It is tempting to speculate that phosphorylated glyoxalase I is involved in the formation of these specific MG-AGEs.

Several mechanisms have been described of how MG-modification on proteins exerts its effect. Reaction of methylglyoxal with amino acids generates reactive oxygen species (16), which can subsequently activate several signaling pathways. Furthermore, methylglyoxal has been reported to covalently modify specific residues of several proteins, hereby regulating their function (14,15,17,18). Another possibility is through the activation of the receptor for advanced glycation endproducts (RAGE). Many signaling pathways have been reported to emanate from RAGE (see chapter 3.3). Although RAGE is expressed in every cell line studied so far (19), under physiological conditions expression levels are low, and RAGE is strongly upregulated under pathological conditions (20). When RAGE was studied throughout the cell cycle, we found a strong regulation of the different isoforms. These differences could be related to specific cell cycle phases and alterations in the expression levels coincided with transitions between phases. In spite of the strong regulation of the different isoforms of RAGE, no alteration in expression of RAGE at the plasmamembrane was measured. We also discovered a new proteolytic cleaved isoform of RAGE, 17 kDa RAGE, which was induced upon IL-3 depletion-induced G_1 arrest. We therefore hypothesize that RAGE signaling could be involved in proliferation or survival. The proteolytic processing of RAGE could thus provide a mechanism for cells to attenuate RAGE signaling. Alternatively, the 17 kDa RAGE itself could exert another function in G_1 arrest or the induction of apoptosis.

Reference List

1. **Thornalley, P. J.** 1990. The Glyoxalase System - New Developments Towards Functional-Characterization of A Metabolic Pathway Fundamental to Biological Life. *Biochemical Journal* **269**:1-11.
2. **Thornalley, P. J.** 2003. Glyoxalase I - structure, function and a critical role in the enzymatic defence against glycation. *Biochemical Society Transactions* **31**:1343-1348.
3. **Kalapos, M. P.** 1999. On the promine/retine theory of cell division: now and then. *Biochimica et Biophysica Acta-General Subjects* **1426**:1-16.
4. **Van Herreweghe, F., J. Q. Mao, F. W. R. Chaplen, J. Grooten, K. Gevaert, J. Vandekerckhove, and K. Vancompernelle.** 2002. Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methyl glyoxal-derived AGE. *Proceedings of the National Academy of Sciences of the United States of America* **99**:949-954.
5. **Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura.** 1989. Sexual-Response in *Saccharomyces-Cerevisiae* - Alteration of Enzyme-Activity in the Glyoxalase System by Mating Factor. *Biochemical and Biophysical Research Communications* **165**:1091-1095.
6. **Inoue, Y., B. Y. Choi, K. Murata, and A. Kimura.** 1990. Sexual-Response of *Saccharomyces-Cerevisiae* - Phosphorylation of Yeast Glyoxalase-I by A Cell Extract of Mating Factor Treated-Cells. *Journal of Biochemistry* **108**:4-6.
7. **Bucking-Throm, E., W. Duntze, L. H. Hartwell, and T. R. Manney.** 1973. Reversible arrest of haploid yeast cells in the initiation of DNA synthesis by a diffusible sex factor. *Exp.Cell Res.* **76**:99-110.
8. **Khan, M. M. K., A. Jan, H. Karibe, and S. Komatsu.** 2005. Identification of phosphoproteins regulated by gibberellin in rice leaf sheath. *Plant Molecular Biology* **58**:27-40.
9. **Shin, R., S. Alvarez, A. Y. Burch, J. M. Jez, and D. P. Schachtman.** 2007. Phosphoproteomic identification of targets of the Arabidopsis sucrose nonfermenting-like kinase SnRK2.8 reveals a connection to metabolic processes. *Proceedings of the National Academy of Sciences of the United States of America* **104**:6460-6465.
10. **Deswal, R. and S. K. Sopory.** 1999. Glyoxalase I from *Brassica juncea* is a calmodulin stimulated protein. *Biochimica et Biophysica Acta-Molecular Cell Research* **1450**:460-467.
11. **Ghosh, S., J. Sen, S. Kalia, and S. Guha-Mukherjee.** 1999. Establishment of synchronization in carrot cell suspension culture and studies on stage specific activation of glyoxalase I. *Methods Cell Sci.* **21**:141-148.
12. **Usui, Y., M. Nakase, H. Hotta, A. Urisu, N. Aoki, K. Kitajima, and T. Matsuda.** 2001. A 33-kDa allergen from rice (*Oryza sativa* L. Japonica) - cDNA cloning, expression, and identification as a novel glyoxalase I. *Journal of Biological Chemistry* **276**:11376-11381.
13. **Fang, X. J., S. X. Yu, J. L. Tanyi, Y. L. Lu, J. R. Woodgett, and G. B. Mills.** 2002. Convergence of multiple signaling cascades at glycogen synthase kinase 3: Edg receptor-mediated phosphorylation and inactivation by lysophosphatidic acid through a protein kinase C-dependent intracellular pathway. *Molecular and Cellular Biology* **22**:2099-2110.
14. **Zhang, H. H., A. I. Lipovsky, C. C. Dibble, M. Sahin, and B. D. Manning.** 2006. S6K1 regulates GSK3 under conditions of mTOR-dependent feedback inhibition of Akt. *Molecular Cell* **24**:185-197.
15. **Maeta, K., S. Izawa, S. Okazaki, S. Kuge, and Y. Inoue.** 2004. Activity of the Yap1 transcription factor in *Saccharomyces cerevisiae* is modulated by methylglyoxal, a metabolite derived from glycolysis. *Molecular and Cellular Biology* **24**:8753-8764.
16. **Yao, D., T. Taguchi, T. Matsumura, R. Pestell, D. Edelstein, I. Giardino, G. Suske, N. Rabbani, P. J. Thornalley, V. P. Sarthy, H. P. Hammes, and M. Brownlee.** 2007. High glucose increases angiotensin-2 transcription in microvascular endothelial cells through methylglyoxal modification of mSin3A. *Journal of Biological Chemistry* **282**:31038-31045.
17. **Yim, H. S., S. O. Kang, Y. C. Hah, P. B. Chock, and M. B. Yim.** 1995. Free-Radicals Generated During the Glycation Reaction of Amino-Acids by Methylglyoxal - A Model Study of Protein-Cross-Linked Free-Radicals. *Journal of Biological Chemistry* **270**:28228-28233.
18. **Sakamoto, H., T. Mashima, K. Yamamoto, and T. Tsuruo.** 2002. Modulation of heat-shock protein 27 (Hsp27) anti-apoptotic activity by methylglyoxal modification. *Journal of Biological Chemistry* **277**:45770-45775.
19. **Speer, O., S. Morkunaite-Haimi, J. Liobikas, M. Franck, L. Hensbo, M. D. Linder, P. K. J. Kinnunen, T. Wallimann, and O. Eriksson.** 2003. Rapid suppression of mitochondrial permeability transition by methylglyoxal - Role of reversible arginine modification. *Journal of Biological Chemistry* **278**:34757-34763.
20. **Brett, J., A. M. Schmidt, S. D. Yan, Y. S. Zou, E. Weidman, D. Pinsky, R. Nowygrod, M. Neeper, C. Przysiecki, A. Shaw, A. Migheli, and D. Stern.** 1993. Survey of the Distribution of A Newly Characterized Receptor for Advanced Glycation End-Products in Tissues. *American Journal of Pathology* **143**:1699-1712.
21. **Schmidt, A. M., S. D. Yan, S. F. Yan, and D. M. Stern.** 2000. The biology of the receptor for advanced glycation end products and its ligands. *Biochim.Biophys.Acta* **1498**:99-111.

Part IV: Acknowledgements

Dankuwel...

... en dan nu het stukje dat iedereen leest. Het dankwoord. Op zich misschien goed dat iedereen het leest, want het is belangrijk. Ik heb ondertussen ontdekt hoe moeilijk doctoreren is en hoeveel een mens bijleert in die periode...en dan is er nog het wetenschappelijke luik ook. Ik besef dan ook dat onderstaande personen bijzonder belangrijk zijn geweest bij het tot stand komen van dit werkje en dat deze woordjes niet toerijkend zullen zijn.

Eerst en vooral wil ik Prof Katia Vancompernelle bedanken. Dankuwel Katia. Ik heb van jou de kans gekregen om dit onderzoek te starten en gaandeweg heb je me ook op andere vlakken kansen gegeven en dingen aangereikt. Op de werkvloer verbaasden je enthousiasme en de manier waarop je positief tegen verschillende uitdagingen aan bleef kijken me keer op keer.

Ook vakgroepvoorzitter Prof. Joël Vanderkerckhove zou ik hard willen bedanken voor de steun tijdens het doctoraat en voor de kans om op het toch wel mooie instituut Rommelaere te werken.

Ook naar mijn co-promotor Prof. Guy Haegeman gaat mijn dank uit. Zonder hem zou dit doctoraat heel wat moeilijker tot stand zijn gekomen.

Een persoon die het mogelijks niet verwacht om bedankt te worden (maar ze zal daar nog wel achter komen) is mevr. Lisa Duchâtelet. Mijn interesse in wetenschap werd voor een zeer groot stuk door haar bepaald, en ze heeft er geen idee van (werkelijk niet, maar ook daar zal ze nog wel achter komen) hoe hard ze me beïnvloed heeft.

Ik was niet de enige die voor Katia werkte. Ik heb gedurende de jaren enkele schitterende collega's gehad. De eerste die ik daar zag, en meteen ook degene die in hetzelfde bootje zit als ik, Stef. U was een fijne collega. Het was ook aangenaam om samen met jou ook deze laatste dingen te doorworstelen. Samen schrijven, elkaar onze nieuwste ontdekkingen van coreldraw aanleren, dat soort dingen....zelfs samen verdedigen binnenkort. Het was plezant. Vervolgens Virginie. Collega's als jij, dat ga ik niet meer vinden, denk ik. Bedankt voor de vele wetenschappelijke en menselijke gesprekken. Ik wens je nog veel succes bij het afronden van je phd en ik weet zeker dat de dingen die je erna gaat doen bijzonder spannend gaan zijn. Dieter, op het eind van de rit naast mezelf de enige man. Bedankt voor de vele babbels, de voetbalpraatjes, def jam fight for NY, je vermogen om te relativeren en gewoon een geweldig toffe collega te zijn. Nog veel succes met je doctoraat. Anneleen, ook jij was een ware sensatie om mee samen te werken. Inmiddels ben je ook met veel belangrijkere dingen bezig dan doctoreren. Ik hoop dat je bevalling binnenkort goed verloopt, en dat je een gezond kindje krijgt waar je veel van kan genieten. Mascha, de buitenlandse post-doc, ook jij bedankt voor de geweldige inbreng in

het labo...vooral de pindakaas met cashewnoten, ik kan je er niet genoeg voor bedanken. Ik wens je verder veel moed en sterkte momenteel. Ook Sigrid, van harte bedankt. Je harde, goede en verzorgde werk was ongelooflijk nuttig. Veel succes voor je nieuwe job. Het kan niet anders dan dat men daar content gaat zijn van jou. Ook de culinaire babbeltjes en dat soort dingen, het was plezant. Mathias, bedankt voor de vele keren dat je geholpen hebt wanneer we samen projecten moesten indienen, wanneer ik weer problemen had met computers...dat soort dingen. Ook Franky ben ik dankbaar.

Ik heb ook het ongelooflijke voorrecht gehad om een studente te mogen begeleiden tijdens een thesis. Een ongelooflijke ervaring. Niet alleen heeft ze bijgedragen tot een mooi deel van deze scriptie, ook de vele gesprekken over 'andromorphoseren van katten', 'de relatie tussen de lengte van de passen en de frequentie van het klikken van de hielen', en nog zoveel meer...een mens moet het een keer meemaken...veel succes nog tijdens jouw verdere wetenschappelijke loopbaan, maar vooral, geniet met volle teugen van het moederschap.

Werken op het Rommelaere werd vooral ook mogelijk gemaakt door Johan en Serge. Jullie leveren schitterend werk. Niet te onderschatten. Ook de mensen van het secretariaat Dirk en Veronique, en in een recent verleden Kristien, super hard bedankt. Ook jullie doen in alle bescheidenheid geweldig werk en het was fijn hoe hard jullie je best deden.

Eervolle vermeldingen gaan meteen ook naar Julie, bedankt voor de aanmoedigingen tijdens de laatste maanden en naar al die mensen die oprecht interesse toonden (Frank, Sofie, Prof. Broekaert, ...) vaak kleine dingen, maar ze zijn wel belangrijk.

Deze scriptie zou niet tot stand zijn gekomen zonder Sylvie. Ik ben je voor veel dankbaar. Voor je steun op bepaalde zware momenten en je vriendschap, voor het nalezen van de scriptie en nog zoveel meer. Jij ook nog succes tijdens het laatste jaartje.

Een hartelijke dankuwel zeker ook voor Erika. Dank voor het nalezen van delen van de thesis en mijn gebrekkig Engels bij te sturen. Jij en Tim (en Alister) hartelijk bedankt ook om gewoon vrienden te zijn.

Ik ben ook dankbaar voor de geweldige vrienden die ik heb. Ze hebben een grenzeloos geduld getoond en zijn nog steeds mijn vrienden, ook al heb ik niet veel tijd voor hen gehad de laatste maanden. Ik kan ze niet allen bedanken, maar enkele wil ik toch bij naam noemen. Tom, je bent me ongelooflijk dierbaar. Ik heb veel steun van je gekregen. Wim en Birgit, zo hard bedankt, voor de aanmoedigingen en zo, en voor de vriendschap. Wim, ook ongelooflijk bedankt dat je iets later dit jaar ook mijn getuige wil zijn. Jan en Joke, vooral de regelmatige smsjes met aanmoedigingen de laatste maanden van Joke. Maarten en Hanna, voor het geduld. A special thanks goes to my dear friend Maete. If anything good came out of this phd, it's my friendship with you. Thanks for the support. Thanks

for taking me to Spain. Thanks for so much more. Cindy, wanneer beginnen we met die cocktailboot?

Een speciale vermelding ook voor een speciaal iemand...Gawein. Ook al zit je nu aan't andere eind van deze mooie planeet, ik heb veel aan je gehad en zal blij zijn wanneer je weer terug in't land bent.

Ik ben nu ongetwijfeld belangrijke mensen aan het vergeten...sorry daarvoor, dat wil niet zeggen dat ik minder dankbaar ben. Ik ben wel zeker dat ik de beste vriendenkring aller tijden heb.

Papa en Brigitte, Mama en Staf, ook naar jullie gaat een ongelooflijke dankuwel uit. Jullie hebben mij altijd gesteund, voor me gebeden, jullie hebben altijd interesse getoond, zijn er altijd voor mij geweest en hebben jezelf vaak weggecijferd voor mij. Zonder jullie zou ik dit niet gekund hebben. Had ik dit niet uitgehouden. Ik hou veel van jullie. Ook dank u aan Luc en Nelly (ook papa en mama). Bedankt voor de interesse en steun.

Ook mijn zussen en broers ongelooflijk veel dank voor de steun, de interesse en het begrip.

Ten slotte Ann. Bedankt voor je liefde en geduld. Bedankt dat je er bent, en dank je dat je mijn vrouw wil zijn. Ik hou van je.

*Nochtans zal ik juichen in de Here,
Jubelen in de God van mijn heil.*